

# Constancy of Expression of the Protein Kinase A Regulatory Subunit R1 $\alpha$ in Hepatoma Cell Lines of Different Phenotypes<sup>1</sup>

Daniela M. Faust, Michael Boshart,<sup>2</sup>  
Tereza Imaizumi-Scherrer, Günther Schütz, and  
Mary C. Weiss<sup>3</sup>

Unité de Génétique de la Différenciation, URA Centre National de la Recherche Scientifique 1149, Institut Pasteur, 28, rue du Dr. Roux, 75724 Paris, France [D. M. F., T. I.-S., M. C. W.], and German Cancer Research Center, Division Molecular Biology of the Cell I, Im Neuenheimer Feld 280, 69120 Heidelberg 1, Germany [M. B., G. S.]

## Abstract

In somatic hybrids between fibroblast microcells and rat hepatoma cells, tissue-specific extinguisher 1 (TSE1), localized to mouse chromosome 11, extinguishes the expression of tyrosine aminotransferase and phosphoenolpyruvate carboxykinase. Recently, it was demonstrated that TSE1 corresponds to R1 $\alpha$ , a regulatory subunit of protein kinase A. Here, we have analyzed whether R1 $\alpha$  could play a role in differentiation of the hepatocyte. It is known that the TSE1/R1 $\alpha$  target genes belong to the group of neonatal functions that are repressed until birth. High expression of R1 $\alpha$  characterizes fetal-type BW1J hepatoma cells in which the neonatal target genes are silent. This R1 $\alpha$  is active in *trans* to extinguish these genes in hybrids between BW1J and Fao adult-type rat hepatoma cells. Reexpression of the target genes is correlated with loss of R1 $\alpha$  and/or overexpression of the mRNA for the hepatocyte-enriched transcription factors HNF4 and HNF3 $\alpha$ . Phenylalanine hydroxylase is shown to be another function negatively regulated by R1 $\alpha$ . In BW cells in which expression of phenylalanine hydroxylase has been activated (after either 5-aza-cytidine treatment or transfection with genomic DNA from adult-type hepatoma cells), no down-regulation of R1 $\alpha$  expression occurs: an independent mechanism overcomes R1 $\alpha$  repression. Finally, dedifferentiated derivatives of the adult-type rat hepatoma cells express neither the R1 $\alpha$  target genes nor the R1 $\alpha$  gene itself. Thus, in three different situations in which modulation of R1 $\alpha$  expression could be anticipated, it fails to occur. It is concluded that somatic cells in culture show *cis*-heritability of the R1 $\alpha$  gene, even under conditions where its effect is selected against.

Received 6/25/93; revised 10/1/93; accepted 10/26/93.

<sup>1</sup> Work in the Heidelberg laboratory was supported by the Deutsche Forschungsgemeinschaft through Grant SFB229, the Leibniz-Programm, and Fonds der Chemischen Industrie, and the Paris laboratory was supported by grants from l'Association pour la Recherche sur le Cancer and La Ligue Nationale contre le Cancer and by a contract from the BRIDGE Programme of the European Economic Community.

<sup>2</sup> Present address: Max Planck Institute for Biochemistry, GeneCenter, 82152 Martinsried, Germany.

<sup>3</sup> To whom requests for reprints should be addressed.

## Introduction

Somatic hybrids obtained from the fusion of cells of different histotypes generally show extinction of expression of the differentiated functions of the parents (1). This phenomenon of negative gene regulation has been studied particularly in hybrids derived from crosses involving hepatoma cells, for which it has been shown that the multiple hepatocyte functions are all extinguished and that reexpression of these functions occurs independently, correlated with loss of chromosomes of the extinguishing parent (2). Whereas these observations implied a genetic basis for the extinction phenomenon, proof was provided by identification of TSEs<sup>4</sup> localized to specific chromosomes (3). The identification of TSE1 was achieved by the study of microcell hybrids, derived from fibroblast microcells fused with rat hepatoma cells and selected for retention of the fibroblast-derived thymidine kinase-bearing chromosome, autosomes 11 and 17, for mouse and human, respectively. In such microcell hybrids, the liver-specific hormone-inducible enzyme TAT was selectively extinguished and reexpressed upon loss of the thymidine kinase carrying chromosome (4).

It was subsequently shown that TSE1 has several target genes, all of them having in common the property of inducibility by cAMP (5, 6). By a thorough analysis of the mechanism of action of TSE1 as well as by positional cloning, it was demonstrated that TSE1 corresponds to the R1 $\alpha$  regulatory subunit of PKA (7-9). Once TSE1 was identified as R1 $\alpha$ , it became clear that well differentiated hepatoma cells have very low levels of R1 $\alpha$ , whereas fibroblasts, and most other cells as well, produce large amounts of the mRNA. In microcell hybrids, the R1 $\alpha$  gene of the fibroblast-derived chromosome remains transcriptionally active, and its product titrates out the hepatoma-derived PKA catalytic subunits in the form of inactive tetramers of catalytic and regulatory subunits. In order to maintain basal transcription of TSE1 target genes in the absence of elevated cAMP, catalytic subunits of PKA must be available to phosphorylate CREB, thereby enhancing its transcription-stimulating activity. Extinction mediated by TSE1/R1 $\alpha$  is entirely reversible upon treatment with cAMP, since cAMP activates PKA by dissociation of its subunits (discussed in Ref. 10).

One of the major assumptions underlying the somatic cell genetic analysis of liver cell differentiation has been that elucidation of the mechanisms responsible for the extinction phenomenon will shed light on mechanisms operating during normal development and differentiation (1). Following this line of reasoning, the study of TSE1 may have revealed

<sup>4</sup> The abbreviations used are: TSE, tissue-specific extinguisher; TAT, tyrosine aminotransferase; cAMP, cyclic AMP; PKA, protein kinase A; CREB, cAMP regulatory element-binding protein; PEPCK, phosphoenolpyruvate carboxykinase; PAH, phenylalanine hydroxylase; G-, glucose-free; kb, kilobase(s); GAPDH, glyceraldehyde-3-phosphate dehydrogenase; cDNA, complementary DNA.

a new level of activity of R1 $\alpha$ , involving it in a regulatory circuit (and/or developmental switch). To approach this possibility, we have analyzed whether R1 $\alpha$ /TSE1 could play a role during ontogeny of the liver. It is known, on the one hand, that TSE1/R1 $\alpha$  prevents transcription of several of the genes encoding gluconeogenic enzymes and, on the other hand, that these genes are not expressed in fetal liver. The neonatal functions are activated only after birth, subsequent to decreases in circulating insulin and increases in glucocorticoids and glucagon, and consequently of cAMP. The possibility that a regulatory subunit of PKA is involved in these processes is an attractive one.


Although much work and speculation have been devoted to the mechanisms involved in the activation of expression of the liver-specific metabolic enzymes (for a review, see Ref. 11), less attention has been given to the aspect of prevention of their precocious expression. Here, we examine whether R1 $\alpha$ /TSE1 could be involved in preventing the precocious expression of genes coding for "neonatal" functions such as gluconeogenic enzymes, and by extension, in specifying the developmental stage of the hepatocyte. The data suggest developmentally programmed heritable down-regulation of R1 $\alpha$ /TSE1 expression as one step for activation of basal expression of the genes encoding neonatal hepatic functions. We show, in addition, that the level of expression of the R1 $\alpha$  gene is an extremely stable trait in cell culture. Thus, regulation of the R1 $\alpha$  gene might be restricted to a "determination" event during ontogenesis.

## Results

**R1 $\alpha$ /TSE1 Contributes to Extinction in Fetal  $\times$  Adult Hepatoma Hybrids.** The hypothesis that R1 $\alpha$ /TSE1 acts in fetal liver to prevent precocious expression of the hormone-inducible neonatal liver-specific enzymes (7, 8) is consistent with the results of genetic analysis of hepatoma cells of fetal and neonatal phenotype. Mouse hepatoma cells of fetal phenotype of the cell line BW1J and its subclones (12) do not show significant expression of the genes repressed by R1 $\alpha$ /TSE1, such as TAT, PEPCK, and PAH (see below). Moreover, this set of genes is expressed in rat hepatoma cells of adult phenotype, such as Fao, and is extinguished upon fusion of Fao cells with BW1J (13). Finally, reexpression of these genes in Fao  $\times$  BW1J hybrids is frequently correlated with loss of mouse chromosome 11 (12), as would be expected if R1 $\alpha$ /TSE1 were responsible for extinction. Moreover, rat hepatoma microcell hybrids containing mouse chromosome 11 from the fetal-like hepatoma cells of line Hepa1a show extinction of expression of the TSE1 target functions (14), implying that R1 $\alpha$  is expressed in Hepa1a cells.

We have compared R1 $\alpha$  mRNA levels in BW1J, Fao, and hybrid clones (FoB) as well as in reexpressing subclones derived from these hybrids. Fig. 1 shows the filiation of the cell lines involved and a Northern blot. TAT, PEPCK, and PAH mRNAs are absent from BW1J but present in Fao cells, whereas R1 $\alpha$  is abundant in BW1J cells but barely detectable for Fao. Consequently, the anticipated exclusivity in expression of R1 $\alpha$  and its target genes applies to hepatoma cells of fetal as well as adult phenotype.

As shown in the family tree of Fig. 1, two generations of hybrids are considered: FoB4 and FoB7 are primary hybrid clones which show extinction of expression of the "neonatal" functions. Reexpressing progeny subclones (4-3, 4-7, 7-b, and 7-IV) were derived by selection in glucose-free medium, where growth depends upon activity of the gluconeogenic pathway, of which PEPCK is a key enzyme; consequently, only cells which reexpress PEPCK can form



**Fig. 1.** *A*, filiation of the cell lines used. BW1J, mouse hepatoma cells of fetal phenotype; Fao, rat hepatoma cells of adult phenotype. FoB4 and FoB7, two different primary hybrid clones obtained by fusion of Fao and BW1J. FoB4-3, FoB4-7, FoB7-b, and FoB7-IV, clones obtained by selection in glucose-free (G<sup>-</sup>) medium, where only cells expressing the gluconeogenic enzymes can survive. *B*, composite Northern blot of the cell lines. *Upper three panels*, probed with cDNA for the R1 $\alpha$ /TSE1 target genes TAT, PEPCK, and PAH (see below). *Middle panel*, probed for R1 $\alpha$ , using an oligonucleotide probe that is complementary to both rat and mouse R1 $\alpha$  mRNA. *Lower panels*, hybridized with probes for liver-enriched transcription factors. Note that mRNA for HNF1 is of two sizes in the rat, at 3.6 and 3.2 kb, owing to the existence of alternate polyadenylation signals; in the mouse, there is only one mRNA of 3.2 kb. For both species, the mRNAs for HNF3 are of three sizes, corresponding to HNF3 $\alpha$  (3.4 kb),  $\beta$  (2.2 kb), and  $\gamma$  (2 kb). Three parallel blots were used: each was hybridized with a rRNA probe (only one example is shown). Lines at right, the specific hybridizing bands, which are of the following sizes: TAT (2.4 kb); PEPCK (2.8 kb); PAH (2.4 and 2.0 kb); R1 $\alpha$  (3.2, 2.9, and 1.7 kb); HNF4 (4.5 kb); HNF1 and HNF3 (see above).

**Table 1** Expression of R1 $\alpha$  and hepatic functions and factors: hybrid clones grouped by phenotype

The values given are specific activity for PKA (pmol/min/mg), and relative values from quantification of Southern blots (R1 $\alpha$  DNA) or RNA blots (all other functions) that are shown in Fig. 1. For the relative mRNA values (R1 $\alpha$  RNA, R1 $\alpha$  target genes, and hepatic transcription factors), the more positive parent was set at 100. Quantitation from the blots (PhosphorImager) includes degraded hybridizing material (where present). Mouse R1 $\alpha$  DNA was quantitated from the 1.9-kb mouse-specific band in Fig. 2, setting BW1 to 100. As background, the region corresponding to the 1.9-kb band from the Fao lane was used. ND, not determined.

	PKA (specific activity)		R1 $\alpha$		R1 $\alpha$ target genes			Hepatic transcription factors	
	Basal	cAMP:fold induction	DNA (mouse)	RNA	TAT	PEPCK	PAH	HNF1	HNF4
Fao	250	12		8	100	100	100	100	100
BW1	8	190	100	100	6	2	0	24	18
FoB4	16	200	27	45	0	4	3	50	51
FoB4-7	20	130	43	64	50	10	23	41	210
FoB7	ND	ND	30	39	0	3	2	10	15
FoB4-3	306	7	<1	11	1570	34	360	34	310
FoB7-b	280	12	12	28	6400	200	680	24	340
FoB7-IV	206	10	9	16	1200	94	280	20	180

colonies (12). The composite Northern blot of Fig. 1 shows that FoB4 and FoB7 cells show little or no signal for the three target genes *TAT*, *PEPCK*, and *PAH*; however, each of these functions is reexpressed, but at variable levels, in the subclones derived in G<sup>-</sup> medium. R1 $\alpha$  expression is inversely correlated with that of the target genes: significant levels of the RNA are observed for the primary hybrid clones, and for FoB4-7 cells, which show the weakest reexpression of *TAT*, *PEPCK*, and *PAH*. Thus, extinction of expression of the hepatic enzymes is correlated with significant levels of the R1 $\alpha$  transcript, whereas their reexpression to parental levels parallels loss of R1 $\alpha$ .

The level of R1 $\alpha$  protein affects the basal activity of PKA (see Ref. 8 and references therein). When assayed in the absence of cAMP, only the free catalytic subunits contribute to kinase activity, whereas in the presence of cAMP, all catalytic subunits are liberated from their regulatory subunits. As shown in Table 1, all of the cell lines analyzed fall into one of two classes: high basal PKA activity and weak cAMP induction, or low basal activity and a 100–200-fold induction. These results are in perfect agreement with R1 $\alpha$  mRNA expression. Thus, all of the clones showing significant basal expression of the three R1 $\alpha$ /TSE1 target genes, *TAT*, *PEPCK*, and *PAH*, are characterized by high basal PKA activity that is correlated with low R1 $\alpha$  mRNA. In contrast, the cell lines characterized by low basal PKA activity and high R1 $\alpha$  expression all show, with one exception, extinction of expression of the target genes. Since basal PKA activity is essential for expression of *TAT* and *PEPCK* (8) and *PAH* (see below) in adult-type hepatoma cells, it appears clear that R1 $\alpha$  expression contributes to extinction of *TAT*, *PEPCK*, and *PAH* in fetal  $\times$  adult hepatoma hybrids.

The one exceptional line, FoB4-7, has been selected as were the others for the reexpression of *PEPCK*. In these cells, there is high R1 $\alpha$  expression and low basal PKA; the three target genes are reexpressed, albeit at relatively low levels (10–50% of the Fao values). We therefore investigated by what mechanism expression of the target genes occurs, in spite of the presence of high levels of R1 $\alpha$ . Overexpression of PKA catalytic subunits cannot be proposed as a mechanism, since total PKA activity in the presence of cAMP is very similar in all cell lines (Table 1). We then analyzed mRNA for some of the liver-enriched transcription factors which either have been demonstrated (see "Discussion") or are likely to be involved in expression of the liver-specific target

genes, to investigate the basis for quantitative variations in the levels of reexpression after selection in G<sup>-</sup> medium. As shown in Fig. 1 and Table 1, cells of the Fao and BW1 parental lines both produce HNF4, HNF1, and HNF3 mRNA. Among the different forms of HNF3,  $\alpha$  is particularly abundant in Fao, and  $\beta$  and  $\gamma$  are the most prominent in BW1 cells.

In the primary hybrid clones that had not been subjected to selection for *PEPCK* reexpression, the mRNAs of HNF4, HNF1, and HNF3 are intermediate between the parental values or are somewhat lower. Among the hybrid clones selected for reexpression of *PEPCK*, all show overexpression of HNF4 RNA, at levels that exceed the Fao value by a factor of 2 or 3: this could account for the strong basal expression of *TAT* and *PAH*. Concerning HNF1 and HNF3, no systematic change is observed. However, for the exceptional clone FoB4-7, which reexpresses the TSE1/R1 $\alpha$  target genes despite high expression of R1 $\alpha$ , mRNAs for both HNF4 and HNF3 show enhanced expression. Moreover, for HNF3, the abundance of form  $\alpha$  compared to  $\beta$  and  $\gamma$  is greatly increased. It can be proposed that enhanced expression of HNF4 and HNF3 $\alpha$  overrides TSE1/R1 $\alpha$ -mediated repression of *TAT*, *PEPCK*, and *PAH* in FoB4-7 cells.

These results clearly implicate R1 $\alpha$  expression as one factor in heritability of the fetal hepatocyte phenotype. Furthermore, high R1 $\alpha$  mRNA expression is dominant in BW1  $\times$  Fao hybrids. In other words, the mouse R1 $\alpha$  gene of BW1 seems not to be affected in *trans* by any mechanism restricting expression of the rat R1 $\alpha$  gene in Fao. Therefore, we determined the number of gene copies of mouse R1 $\alpha$  retained on average in the hybrids and subclones selected in G<sup>-</sup> medium (some chromosomal heterogeneity cannot be excluded). Fig. 2 shows a Southern blot hybridized with an R1 $\alpha$ -specific probe which detects fragments of different size representing the rat and mouse R1 $\alpha$  alleles, respectively. The quantification of the hybridization signals reveals a perfect correlation of R1 $\alpha$  expression with gene dosage of the mouse R1 $\alpha$  allele, namely the number of mouse chromosomes 11 retained on average in the respective cell line (Fig. 2; Table 1). Furthermore, we find R1 $\alpha$  mRNA expressed from the mouse allele but not from the rat allele in the hybrid clones shown in Fig. 1, as concluded from slight differences in electrophoretic mobility of the rat and mouse R1 $\alpha$  mRNA species (data not shown). Thus, no modification in expression of either the rat or mouse R1 $\alpha$  gene ensues from the confron-




Fig. 2. Southern blot with genomic DNA (EcoRI digested) from the cell lines analyzed in Fig. 1 and probed with the same R1 $\alpha$  oligonucleotide. The unique mouse- and rat-specific bands are of 1.9 and 6.5 kb, respectively.

tation of active and inactive genes in a single nucleus: the mouse gene remains active, and the rat gene remains silent.

**Selection of Fetal Hepatoma Cells for Phenylalanine Hydroxylase Expression Does Not Result in R1 $\alpha$  Down-Regulation.** Perhaps the most striking result obtained from this analysis is the invariability in expression of the R1 $\alpha$  gene. We attempted to challenge this invariability. Selection for reexpression of the PEPCK gene in the context of hybrid cells (see previous section) may be facilitated by the possibility of chromosome segregation and loss of the active R1 $\alpha$  gene. Therefore, we turned to another system to select for expression of an R1 $\alpha$ /TSE1 target function in mouse hepatoma cells with a fetal phenotype and high R1 $\alpha$  expression.

The gene encoding phenylalanine hydroxylase, another of the adult hepatic functions, is known to be inducible by glucocorticoids and cAMP, just like TAT and PEPCK. It therefore was reasonable to postulate that it would be sensitive to regulation by R1 $\alpha$ /TSE1. As anticipated, in several clones of rat hepatoma cells stably transfected by an inducible R1 $\alpha$ -encoding plasmid (8), the level of PAH is inversely correlated with the R1 $\alpha$  expression levels (Fig. 3). These results, along with those already presented in Fig. 1 and Table 1, lead us to conclude that PAH belongs to the class of R1 $\alpha$ /TSE1 target genes.

Consequently, we have analyzed R1 $\alpha$  levels in cells where PAH has been activated. The cell lines all derive from BWTG3. A first series, MIRZA cells, was obtained by 5-azacytidine treatment and selection in tyrosine-deficient medium where only PAH<sup>+</sup> cells can proliferate. A second series of cell lines (MIRO) derives from genomic DNA transfection and was selected in tyrosine-free medium (15). Both the MIRZA and MIRO cells were characterized for basal expression of PAH mRNA: in all cases, a significant signal was observed (Fig. 4). However, in no case was a signal observed for PEPCK (data not shown), and only one of the MIRZA lines showed TAT expression. These genes are clearly not co-regulated in BWTG3 cells selected for activation of the PAH gene. Furthermore, R1 $\alpha$  mRNA was equally expressed, at BWTG3 levels, in MIRO and MIRZA lines (Fig. 4).

As shown above for FoB4-7 cells, significant basal PAH activity can be obtained even without loss of R1 $\alpha$  expression. However, examination of the mRNAs for the liver-enriched transcription factors HNF1, 3, and 4 failed to reveal overexpression of one of them that might compensate for the continued presence of R1 $\alpha$  (Fig. 4), as suggested for reexpression in FoB4-7 hybrids. Further experiments will be required to elucidate the mechanism involved in PAH activation. Nevertheless, the question posed has been clearly answered: even when selection is carried out under condi-




Fig. 3. PAH expression in FTO2B cells and clones transfected with R1 $\alpha$ (mut). The R1 $\alpha$ (mut) plasmid used for the transfection as well as the selection of clones is described in Ref. 8. R1 $\alpha$ (mut) corresponds to a cDNA mutated at the cAMP binding site; its expression is inducible by Zn (added as ZnCl<sub>2</sub> at 100 mM concentration to the culture medium for 18 h). The strong PAH signal (for mRNA sizes, see Fig. 1) obtained with RNA from FTO2B cells is not modified by the addition of ZnCl<sub>2</sub>. The transfector clones 12, 17, and 21 display a reduced signal, which is further decreased in the presence of Zn. The same filter was hybridized with the R1 $\alpha$  probe (not shown). Relative PAH mRNA values were calculated as detailed in the introduction to Table 1; R1 $\alpha$  was determined by scintillation counting of the excised bands.

tions in which the active R1 $\alpha$  gene cannot be lost, the cells use some mechanism other than regulation of R1 $\alpha$  expression to respond to the selective pressure.

**Distinct Mechanisms Are Involved in Extinction and De-differentiation.** In earlier work, it has been shown that de-differentiated variants derived from H4II or Fao cells fail to express the group of adult liver-specific genes that have been identified as targets of negative regulation by R1 $\alpha$ . In addition, in most cases, the change is dominant negative: in hybrids between dedifferentiated variants and their differentiated parent, the dedifferentiated state is epistatic (2). To investigate the possibility that activation of R1 $\alpha$  might be correlated with dedifferentiation, a panel of variants was examined for expression of the R1 $\alpha$ /TSE1 target genes and for R1 $\alpha$  mRNA (Fig. 5). H4II and Fao rat hepatoma cells, three independent variant lines (H5, C2, p4), one revertant of a variant (C2Rev7), and mouse fibroblasts (Cl1D L cells) were analyzed. The distribution of RNAs for TAT, PEPCK, and PAH was as expected: present and, where tested, dexamethasone inducible, in H4II, Fao, and C2Rev7 revertant cells, whereas the corresponding RNAs were entirely absent from the variants and from fibroblasts. In none of the hepatoma-derived variants was R1 $\alpha$  activated. Finally, as already noted for fibroblast-hepatoma hybrid cells showing extinction and reexpression of hepatic functions (16), the distribution of RNA for the hepatocyte-enriched transcription factors HNF4 and HNF1 exactly parallels the differentiation state of the cells. R1 $\alpha$  is clearly not a negative regulator that comes into play during the dedifferentiation of H4II or Fao cells, nor is its expression a consequence of dedifferentiation. We conclude that extinction in hepatoma hybrids and dedifferentiation of hepatoma cells in culture are mediated by distinct mechanisms.

## Discussion

The results presented here are relevant to three problems of hepatic differentiation: the first concerns a possible role for




Fig. 4. Northern blot of BW TG3 subclones obtained by selection for expression of the *PAH* gene. For derivations of the lines, see "Materials and Methods." The primary MIRO transfectants are designated by a number, and the secondary transfectants by a number and a letter. The sizes of the specific mRNAs are given in the legend to Fig. 1. GAPDH (1.4 kb) was used for normalization.

*R1α* in the fetal to adult transition; the second, the combinatorial action of regulatory factors specifying the hepatic phenotype; and the third, the *cis*-heritability of the expression state of the *R1α* gene.

Now that the first tissue-specific extinguisher has been cloned, the supposition that the mechanisms involved in the extinction of differentiated functions in intertypic hybrids are operative during development (1) can be tested directly. It would be predicted that *R1α/TSE1* expression is low in the adult liver: this has been observed to be true<sup>5</sup> (17). Moreover, if *R1α/TSE1* expression is implicated in hepatocyte ontogeny, it should be high in the fetal liver, where most of the *R1α/TSE1* target genes remain silent until birth. However, this second prediction is more difficult to test than would be expected, because the fetal liver is primarily a hematopoietic tissue, and blood cells are particularly rich in *R1α*. Consequently, tests of fetal liver extracts or RNA are not informative, and *in situ* hybridization of fetal liver sections is difficult to interpret owing to the high density of heavily labeled blood cells (data not shown).




Fig. 5. Composite Northern blot of differentiated rat hepatoma cells of adult phenotype and of dedifferentiated variant derivatives. Fao and H4II, cells treated (+) or not (-) with 1 mM dexamethasone for 24 h. For origins of the variants, see text. Clone 1D (cl1D) is a fibroblastic L cell, included as a positive control for the *R1α* hybridization. For mRNA sizes, see Figs. 1 and 4.

Alternatively, this prediction can be tested using hepatoma cells of fetal phenotype. BW cells present the expression profile (13) anticipated of fetal hepatocytes: production of  $\alpha$ -fetoprotein, albumin, and other serum proteins, and absence of expression of the group of enzymes implicated in liver-specific metabolism, whose expression first occurs at the time of birth (11). BW cells proved to express *R1α* abundantly: high levels of the mRNA are observed on Northern blots, and enough regulatory subunits are present to inactivate all available catalytic subunits as concluded from the absence of basal PKA activity. Although it cannot be affirmed that *R1α* alone is responsible in the BW cells for inactivity of the adult liver functions, we do know that *R1α/TSE1* expressed by these cells can act in *trans* to extinguish the expression of a set of liver-specific genes, including *TAT* and *PEPCK* (this work, and Ref. 14). However, additional factors must prevent expression of the *R1α/TSE1* target genes in BW cells, since the addition of cAMP does not lead to their expression (data not shown). Moreover, *R1α* expression is not the unique mechanism underlying the epistasis of the fetal phenotype in BW  $\times$  Fao hybrids, for the entire group of neonatal functions, including genes not affected by *R1α/TSE1*, is extinguished (13). Consequently, additional *trans*-acting mechanisms must be involved in determining the phenotypic profile of the fetal-type hepatoma cells. *R1α*

<sup>5</sup> M. Boshart and S. Ruppert, unpublished observations.

would then represent only one of several factors that act to inhibit the precocious expression of a set of adult metabolic functions of hepatocytes.

In earlier work, it was observed that loss from BW  $\times$  Fao hybrids of chromosome 11, and by extension TSE1/R1 $\alpha$ , is not essential to observe reexpression under selective pressure of the gluconeogenic enzymes (12). The hybrid FoB4-7 is an example. In these cells, abundant R1 $\alpha$  expression is maintained but apparently is compensated for by enhanced expression of the RNAs encoding HNF4 and HNF3. It is known that the enhancer of the *TAT* gene at -3.6 kb contains closely juxtaposed binding sites for CREB and for HNF4. Using a transient expression assay, it has been shown that either element alone is unable to activate a heterologous promoter, but that the two elements are interdependent, resulting in a synergistic response (7, 18, 19). In FoB4-7 cells, the reduced availability of phosphorylated CREB can apparently be compensated for by an increased level of HNF4, implying that the element interdependence deduced from transfection studies reflects the situation occurring *in vivo*. Thus, analysis of fetal  $\times$  adult hepatoma hybrids supports the concept that complete extinction results from the combined action of multiple genetic loci. For example, Nitsch *et al.* (18) have demonstrated that the total extinction of the *TAT* gene in hepatoma  $\times$  fibroblast hybrids is due to a deficiency in several *trans*-acting factors implicated in regulation of each of the three enhancers controlling *TAT* transcription.

Dedifferentiated variants of the rat hepatoma provide an independent case of an apparently coordinate shutdown of expression of the entire group of adult hepatic functions, including the gluconeogenic functions such as TAT. Here, there is no change in expression of the R1 $\alpha$  gene: it remains silent, just as in the parental (adult phenotype) hepatoma line. As has been observed by others (20-24), dedifferentiation is accompanied by loss of HNF4 and HNF1 mRNAs. Consequently, loss of expression of the same group of functions in fetal  $\times$  adult hepatoma cell hybrids and in dedifferentiated hepatoma variants clearly is due to distinct mechanisms, which in the case of the *TAT* gene both converge to the same regulatory element, the enhancer at -3.6 kb. That HNF4 plays an important role in both regulatory situations supports the suggestion of HNF4 being a "master regulator" of liver development (see Ref. 21).

A second aspect of R1 $\alpha$  that emerges from this analysis is the remarkable stability of its expression. Although *cis*-heritability of R1 $\alpha$  expression in the context of a fibroblast genome was apparent from the microcell hybrid analysis, there was no *a priori* reason to expect that this would be a general feature of the R1 $\alpha$  gene. Nevertheless, it is very clear that R1 $\alpha$  expression is fixed in hepatoma cells of fetal phenotype, and in spite of selection for progeny that reexpress R1 $\alpha$ -repressed functions, including PEPCK in rat hepatoma-mouse hepatoma hybrids and PAH in mouse hepatoma cells, the cells use mechanisms other than down-regulation of R1 $\alpha$  to establish the phenotype that is selected for.

This leads us to a new problem concerning R1 $\alpha$  and its role during ontogeny. These results, together with the dominant extinction effect elicited by the TSE1/R1 $\alpha$ -carrying chromosome in monochromosomal hybrids, implies that the R1 $\alpha$  gene is not subject to regulation by *trans*-acting factors differentially produced in somatic cells of diverse tissue origin or developmental stage. If this is true, it is difficult to imagine how down-regulation of R1 $\alpha$  can be achieved in hepatocytes at birth. The analysis of hybrids presented here has shown that hepatocytes of adult phenotype do not produce *trans*-acting factors capable of down-regulating R1 $\alpha$ ,

nor do the "fetal" cells appear to be capable of activating the silent R1 $\alpha$  gene of the "adult" parent: inheritance of the pattern of R1 $\alpha$  expression of each genome is *cis*-autonomous. This is unusual behavior for a gene involved in cell differentiation, for essentially all liver-specific functions, and even liver-enriched transcription factors (2, 3, 16), are subject to extinction, reexpression, and activation in hybrid cells: in other words, they are regulated by mechanisms acting in *trans*. Consequently, the change in R1 $\alpha$  expression during the fetal to adult transition must occur in a context that cannot be reproduced in cell lines. Hormonal changes (25) which occur around birth could lead to a cascade of modifications at the chromatin level that become "irreversible." These modifications would be characterized by an initiation mechanism, followed by a maintenance phase: somatic cell lines could present their remarkable stability of R1 $\alpha$  expression because of the maintenance mechanism. Formally, this situation of *cis*-heritability presents parallels with imprinting, with X chromosome inactivation, and with the apparently irreversible event of determination during the course of embryogenesis (26).

## Materials and Methods

**Cell Lines and Culture.** BW7G3 and BW1J are mouse hepatoma lines (13, 27) of fetal phenotype, derived, like Hepa1a cells, from the mouse BW7756 transplantable hepatoma. Hybrids of Fao rat hepatoma cells and BW1J have been described (13). Transfectant clones designated MIRO were isolated following cotransfection of BW7G3 with pSVneo and genomic DNA from H4II cells, and selection first in G418 and then in tyrosine-free medium (detailed in Ref. 15). MIRZA cells were selected in tyrosine-free medium (see Ref. 15) from BW7G3 cells treated for 48 h with 10  $\mu$ M 5-azacytidine; a 2-week expression time preceded selection. Rat hepatoma lines H4II and Fao as well as dedifferentiated (H5, p4, and C2) and revertant (C2Rev7) cells have been described (28-30). FTO2B cells (31) were transfected with a plasmid encoding mutant R1 $\alpha$ . Isolation and properties of clones 17 and 21 are detailed (8); clone 12, used here, shows properties similar to those of clone 17 (8).

**Preparation of Extracts and PKA Assay.** Cell extracts were prepared from confluent cultures, and assays for PKA activity were performed as described (8). Assays were carried out at 30°C in the presence and absence of cAMP and the protein kinase inhibitor (PKI) (1-31) peptide. Basal PKA specific activity is the PKI-inhibitable activity in the absence of cAMP. PKA specific activity is expressed in pmol/min/mg protein.

**Preparation of RNA and DNA.** RNA was prepared from the cytoplasmic supernatant of lysed cells (see Ref. 15) using cultures parallel to those for the preparation of extracts. The nuclei were used to isolate genomic DNA (32), which was digested with EcoRI as indicated by the supplier.

**Northern and Southern Blot Analysis.** Digested DNA or RNA was separated by electrophoresis in agarose gels and transferred to Nylon membranes (Hybond N; Amersham) using a Vacu-blot transfer system (LKB).

The following hybridization probes were used. R1 $\alpha$ : the oligonucleotide probe described in the legend to Fig. 2C of Ref. 8 under the conditions indicated there. HNF3: a riboprobe transcribed from a pBluescript vector containing a 300-base pair polymerase chain reaction fragment encompassing the fork head domain of the mouse HNF3 $\alpha$  gene (Ref. 33; kindly provided by H. Hiemisch, German Cancer Research Center). HNF4: a riboprobe transcribed from a pBluescript vector (kindly provided by S. Taraviras, German

Cancer Research Center) containing the rat HNF4 cDNA (Ref. 34; a kind gift of J. Darnell, Rockefeller University). TAT, PEPCK, PAH, HNF1, GAPDH, and 28S ribosomal RNA random primed probes (kit from Boehringer Mannheim) were derived from the following plasmids: pTAT1 (35), pCK10 (36), pmPAH20 (15), pRSV-HNF1 (37), pGAPDH (38), and p28S (39).

Riboprobes were hybridized at 70°C in the solution given in Ref. 8 and washed off at 70–75°C for 3 × 10 min in 0.1× standard saline citrate-1% sodium dodecyl sulfate. Random primed probes were hybridized as described (40).

The filters were directly scanned, and the signals were quantified with a PhosphorImager (Molecular Dynamics).

### Acknowledgments

We are grateful to Doris Cassio and Catherine Fougère-Descharette for giving us the hybrid lines and to Solange Papelard for preparation of the manuscript.

### References

1. Ephrussi, B. Hybridization of Somatic Cells. Princeton, NJ: Princeton University Press, 1972.
2. Weiss, M. C., Cassio, D., and Sellem, C. H. Cell differentiation: contributions of somatic cell genetics. *Cancer Surv.*, 7: 295–302, 1988.
3. Gourdeau, H., and Fournier, R. E. K. Genetic analysis of mammalian cell differentiation. *Annu. Rev. Cell Biol.*, 6: 69–94, 1990.
4. Killary, A. M., and Fournier, R. E. K. A genetic analysis of extinction: *trans*-dominant loci regulate expression of liver-specific traits in hepatoma hybrid cells. *Cell*, 38: 523–534, 1984.
5. Thayer, M. J., and Fournier, R. E. K. Hormonal regulation of *TSE1*-repressed genes: evidence for multiple genetic controls in extinction. *Mol. Cell. Biol.*, 9: 2837–2846, 1989.
6. Ruppert, S., Boshart, M., Bosch, F. X., Schmid, W., Fournier, R. E. K., and Schütz, G. Two genetically defined *trans*-acting loci coordinately regulate overlapping sets of liver-specific genes. *Cell*, 61: 895–904, 1990.
7. Boshart, M., Weih, F., Schmidt, A., Fournier, R. E. K., and Schütz, G. A cyclic AMP response element mediates repression of tyrosine aminotransferase gene transcription by the tissue-specific extinguisher locus *Tse-1*. *Cell*, 61: 905–916, 1990.
8. Boshart, M., Weih, F., Nichols, M., and Schütz, G. The tissue-specific extinguisher locus *TSE1* encodes a regulatory subunit of cAMP-dependent protein kinase. *Cell*, 66: 849–859, 1991.
9. Jones, K. W., Shapero, M. H., Chevrette, M., and Fournier, R. E. K. Subtractive hybridization cloning of a tissue-specific extinguisher: *TSE1* encodes a regulatory subunit of protein kinase A. *Cell*, 66: 861–872, 1991.
10. Boshart, M., Nitsch, D., and Schütz, G. Extinction of gene expression in somatic cell hybrids—a reflection of important regulatory mechanisms. *Trends Genet.*, 9: 240–245, 1993.
11. Greengard, O. Enzymatic differentiation in mammalian tissues. *Essays Biochem.*, 7: 159–205, 1971.
12. Fougère-Descharette, C., and Weiss, M. C. Interaction of *trans*-acting factors and re-expression of liver functions in hepatoma hybrid cells. *Mol. Biol. Med.*, 7: 97–103, 1990.
13. Cassio, D., and Weiss, M. C. Expression of fetal and neonatal hepatic functions by mouse hepatoma-rat hepatoma hybrids. *Somat. Cell Genet.*, 5: 719–738, 1979.
14. Peterson, T. C., and Fournier, R. E. K. Genetic activity of a *trans*-regulatory locus in hepatoma hybrid cells. *Mol. Biol. Med.*, 7: 109–116, 1990.
15. Faust, D. M., Imaizumi-Scherrer, T., Fulchignoni-Lataud, M-C., Catherin, A-M., Iost, I., and Weiss, M. C. Activation of phenylalanine hydroxylase expression following genomic DNA transfection of hepatoma cells. *Differentiation*, 44: 74–79, 1990.
16. Griffio, G., Hamon-Benais, C., Angrand, P. O., Fox, M., West, L., Lecoq, O., Povey, S., Cassio, D., and Weiss, M. C. HNF4 and HNF1 as well as a panel of hepatic functions are extinguished and reexpressed in parallel in chromosomally reduced rat hepatoma-human fibroblast hybrids. *J. Cell Biol.*, 121: 887–898, 1993.
17. Cadd, G., and McKnight, G. S. Distinct patterns of cAMP-dependent protein kinase gene expression in mouse brain. *Neuron*, 3: 71–79, 1989.
18. Nitsch, D., Boshart, M., and Schütz, G. Extinction of tyrosine aminotransferase gene activity in somatic cell hybrids involves modification and loss of several transcriptional activators. *Genes & Dev.*, 7: 308–319, 1993.
19. Nitsch, D., Boshart, M., and Schütz, G. Activation of the tyrosine aminotransferase gene is dependent on synergy between liver-specific and hormone-responsive elements. *Proc. Natl. Acad. Sci. USA*, 90: 5479–5483, 1993.
20. Cereghini, S., Yaniv, M., and Cortese, R. Hepatocyte dedifferentiation and extinction is accompanied by a block in the synthesis of mRNA coding for the transcription factor HNF1/LFB1. *EMBO J.*, 9: 2257–2263, 1990.
21. Kuo, C. J., Conley, P. B., Chen, L., Sladek, F. M., Darnell, J. E., Jr., and Crabtree, G. R. A transcriptional hierarchy involved in mammalian cell-type specification. *Nature (Lond.)*, 355: 457–461, 1992.
22. Angrand, P. O., Rousset, J. P., and Weiss, M. C. Cell phenotype, binding affinity, and promoter structure modulate transactivation by HNF1 and LAP. *J. Cell Sci.*, 103: 1083–1092, 1992.
23. Angrand, P. O., and Weiss, M. C. HNF1 and HNF4 protein abundance correlates with the degree of differentiation of rat hepatoma cells. In: *Model Systems of Hepatocyte Culture and Gene Regulation*. Paris: INSERM/John Libbey Eurotext Eds., in press, 1993.
24. Herbst, R. S., Nielsch, U., Sladek, F., Lai, E., Babiss, L. E., and Darnell, J. E., Jr. Differential regulation of hepatocyte-enriched transcription factors explains changes in albumin and transthyretin gene expression among hepatoma cells. *New Biol.*, 3: 289–296, 1991.
25. Girard, J. La gluconéogénèse: une voie métabolique essentielle au maintien de l'homéostasie glucidique du nouveau-né. *Med. Sci.*, 9: 297–306, 1993.
26. Riggs, A. D., and Pfeifer, G. P. X-chromosome inactivation and cell memory. *Trends Genet.*, 8: 169–174, 1992.
27. Szpirer, C., and Szpirer, J. A mouse hepatoma cell line which secretes several serum proteins including albumin and  $\alpha$ -fetoprotein. *Differentiation*, 4: 85–91, 1975.
28. Pitot, H. C., Peraino, C., Morse, P. A., and Potter, V. A. Hepatoma in tissue culture compared with adapting liver *in vitro*. *Natl. Cancer Inst. Monogr.*, 13: 229–242, 1964.
29. Descharette, J., and Weiss, M. C. Characterization of differentiated and dedifferentiated clones from a rat hepatoma. *Biochimie*, 56: 1603–1611, 1974.
30. Descharette, J., Moore, E. E., Dubois, M., and Weiss, M. C. Dedifferentiated variants of a rat hepatoma: reversion analysis. *Cell*, 19: 1043–1051, 1980.
31. Killary, A. M., Lugo, T. G., and Fournier, R. E. K. Isolation of thymidine kinase-deficient rat hepatoma cells by selection with bromodeoxyuridine, Hoechst 33258, and visible light. *Biochem. Genet.*, 22: 201–213, 1984.
32. Pellicier, A., Wigler, M., and Axel, R. The transfer and stable integration of the HSV thymidine kinase gene into mouse cells. *Cell*, 14: 133–141, 1978.
33. Lai, E., Prezioso, V. R., Tao, W., Chen, W. S., and Darnell, J. E., Jr. Hepatocyte nuclear factor 3 $\alpha$  belongs to a gene family in mammals that is homologous to the *Drosophila* homeotic gene fork head. *Genes & Dev.*, 5: 416–427, 1991.
34. Sladek, F. M., Zhong, W., Lai, E., and Darnell, J. E., Jr. Liver-enriched transcription factor HNF-4 is a novel member of the steroid hormone receptor superfamily. *Genes & Dev.*, 4: 2353–2365, 1990.
35. Scherer, G., Schmid, W., Strang, C., Röwekamp, W., and Schütz, G. Isolation of cDNA clones coding for rat tyrosine aminotransferase. *Proc. Natl. Acad. Sci. USA*, 79: 7205–7208, 1982.
36. Yoo-Warren, H., Monahan, J. E., Short, J., Short, H., Bruzel, A., Wynshaw-Boris, A., Meissner, H. M., Samols, D., and Hanson, R. W. Isolation and characterization of the gene coding for cytosolic phosphoenolpyruvate carboxykinase (GTP) from the rat. *Proc. Natl. Acad. Sci. USA*, 80: 3656–3660, 1983.
37. Rey-Campos, J., Chouard, T., Yaniv, M., and Cereghini, S. vHNF1 is a homeoprotein that activates transcription and forms heterodimers with HNF1. *EMBO J.*, 10: 1445–1457, 1991.
38. Piechaczyk, M., Blanchard, J. M., Marty, L., Dani, C., Panabieres, F., El Sabouty, S., Fost, P., and Jeanteur, P. Post-transcriptional regulation of glyceraldehyde-3-phosphate-dehydrogenase gene expression in rat tissues. *Nucleic Acids Res.*, 12: 6951–6963, 1984.
39. Tiemeier, D. C., Tilghman, S. M., and Leder, P. Purification and cloning of a mouse ribosomal gene fragment in coliphage lambda. *Gene (Amst.)*, 2: 173–191, 1977.
40. Church, G., and Gilbert, W. Genomic sequencing. *Proc. Natl. Acad. Sci. USA*, 81: 1991–1995, 1984.