

Chemische Berichte

Gesellschaft
Deutscher Chemiker

Founded 1868

January 1994 · Volume 127 · Issue 1 · Pages 1 – 274

| | | | |
|-------------------------|---|--|---|
| Editorial Board: | Wolfgang Beck Rolf Gleiter Klaus Hafner Henning Hopf | Martin Jansen Klaus Müllen Heinrich Nöth Werner Schroth | Helmut Schwarz Werner Tochtermann Heinrich Vahrenkamp |
|-------------------------|---|--|---|

| | | | |
|---------------------------|------------------------------------|------------------------|----------------|
| Managing Editor: | Robert Temme | Administration: | Hannelore Holm |
| Associate Editors: | Udo Eberhardt Johannes Strehlow | Production: | Olga Olivecká |

| | | | |
|--------------------------|--|-------------------|---|
| Editorial Office: | Chemische Berichte P.O. Box 101161 D-69451 Weinheim Federal Republic of Germany Telephone (internat.) +49 6201 606-255 Telefax (internat.) +49 6201 606-328 | Publisher: | VCH Verlagsgesellschaft mbH P.O. Box 101161 D-69451 Weinheim Federal Republic of Germany Telephone (internat.) +49 6201 606-0 Telefax (internat.) +49 6201 606-328 Telex 465516 vchwh d |
|--------------------------|--|-------------------|---|

Die Chemischen Berichte erscheinen monatlich. Die Bezugsbedingungen für Mitglieder der Gesellschaft Deutscher Chemiker werden auf Anfrage von der Geschäftsstelle, Carl-Bosch-Haus, Varrentrappstraße 40–42, Postfach 900440, D-60444 Frankfurt, mitgeteilt. Bestellungen richten Sie bitte an Ihre Fachbuchhandlung oder unmittelbar an den Verlag. Abbestellungen nur bis spätestens 3 Monate vor Ablauf des Kalenderjahres. Die Lieferung erfolgt auf Rechnung und Gefahr des Empfängers. Gerichtsstand und Erfüllungsort: Weinheim.

Subscription rates (1994) including postage and handling:

Annual subscription: DM 1590.– (Germany), ÖS 11468.– (Austria), SFr 1468.– (Switzerland), GB£ 647.– (Great Britain), US\$ 1030.– (USA), DM 1617.– (elsewhere); single issue: DM 152.– (Germany; elsewhere on request).

Back volumes of Chemische Berichte are available on microfilm (further information and prices on request).

For the USA and Canada: Chemische Berichte (ISSN 0009–2940) is published monthly by VCH Publishers, Inc., 303 N.W. 12th Avenue, Deerfield Beach FL 33442-1788; Telefax (305) 428-8201; Telephone (305) 428-5566 or (800) 422-8824. Second-class postage paid at Deerfield Beach FL 33441. Annual subscription price: US \$ 1030.00, single issue: US \$ 107.00 including postage and handling charges. Reduced rate for individual members of the American Chemical Society on request. – Printed in the Federal Republic of Germany.

U.S. POSTMASTER: Send address changes to Chemische Berichte, c/o VCH Publishers, Inc., 303 N.W. 12th Avenue, Deerfield Beach FL 33442-1788.



Autorenregister / Author Index

- Abele, H., 145
Amrein, W., 97
Anders, C., 231
Bauer, I., 173
Binger, P., 39
Bläser, D., 73
Bock, H., 55
Boese, R., 73, 173, 191
Boldt, P., 219
Born, D., 173
Braunschweig, H., 91
Buchweitz, J., 23
Bussenius, J., 247
Butenschön, H., 137
Carell, T., 223
Cremer, D., 173
Dau-Schmidt, J.-P., 205, 213
de Meijere, A., 237
Döhling, A., 223
Döring, D., 219
Eberbach, W., 247
Englert, U., 87, 93
Enk, M., 127
Feucht, C., 113
Fiegenbaum, P., 127
Frey, R., 101
Gabor, B., 39
Gompper, R., 23
Haas, A., 145
Hanack, M., 113
Hansmersmann, U., 127
Haske, S., 73
Heidel, H., 271
Helmchen, G., 271
Herrmann, W. A., 47
Hintze, F., 267
Hohla, M., 119
Hünig, S., 165
Huttner, G., 271
Jeitschko, W., 15
Jones, P. G., 219
Jutzi, P., 107
Kalinowski, H.-O., 191, 201
Karraß, S., 261
Kaschube, W., 127
Kauffmann, T., 127
Klaunzer, N., 165
Kleine, M., 47
Köhler, J. E. H., 119
König, W. A. 119
Köpfer, H., 219
Krieger, C., 231
Krüger, C., 39
Krynitz, U., 55
Kuck, D., 151
Kühn, F. E., 47
Laber, N., 247
Langhauser, F., 39
Lieb, M., 145
Lindermeier, T., 107
Linßen, T., 113
Linti, G., 101
Maier, G., 173, 191, 201
Mayr, H., 205, 213
Meuret, J., 55
Meyer, F., 93
Mink, J., 47
Mix, A., 107
Müller, T., 247
Mynott, R., 39
Näther, C., 55
Neumann, B., 107
Neumann, E., 151
Neumüller, B., 67
Niediek, K., 67
Nöth, H., 81
Nürnberg, O., 27
Paetzold, P., 87, 91, 93
Papenberg, M., 127
Polborn, K., 23, 101
Raabe, N., 261
Richters, M., 119
Robl, C., 23, 97
Romão, C. C., 47
Rückemann, A., 231
Sailer, M.-T., 23
Sander, W., 267
Schäfer, M., 27
Schmid, G., 73
Schneider, L., 87
Schuster, A., 151
Schwarz, H., 261
Sitzmann, H., 3
Spaniol, T. P., 91
Staab, H. A., 223, 231
Stammler, H.-G., 107
Stolpmann, H., 81
Thomann, M., 81
Toliopoulos, E., 127
Tremel, W., 11, 15
Trog, R.-S., 219
Untiedt, S., 237
Vogel, R., 271
Vomhof, T., 15
Walsh, R., 237
Wedemann, P., 39
Weigand, W., 23, 97
Welke, S., 127
Wenner, H., 165
Werner, H., 27
Wolf, J., 27
Wolf, R., 173, 191, 201
Wolmershäuser, G., 3
Wortmann, U., 15
Wünsch, R., 97
Zaika, D., 73
Zhou, P., 3
Zwingenberger, J., 145


Bemerkungen der Redaktion

- Die Chemischen Berichte (zu zitieren als Chem. Ber.) setzen seit 1946 die Berichte der Deutschen Chemischen Gesellschaft (zu zitieren als Ber. Dtsch. Chem. Ges.) fort. Die „Berichte“ enthalten Arbeiten aus dem Bereich der Anorganischen und Elementorganischen Chemie (Teil A) sowie der Organischen und Physikalisch-Organischen Chemie (Teil B). Bei der Organischen Chemie liegt der Schwerpunkt auf allgemeinen synthetischen Methoden und mechanistischen Problemen.
- Die „Berichte“ erscheinen monatlich; ein Register beschließt jeden Jahrgang.
- Die Verantwortung für ihre Mitteilungen tragen die Verfasser selbst. – Der Korrespondenzautor ist durch einen hochgestellten Stern hervorgehoben. Die Zugehörigkeit der Autoren zu den Instituten ist durch hochgestellte Kleinbuchstaben markiert.
- Es werden grundsätzlich nur Arbeiten aufgenommen, die vorher weder im Inland noch im Ausland veröffentlicht worden sind.
- Eine Anweisung zur Abfassung von Manuskripten für die „Berichte“ wird auf Wunsch zugestellt (siehe auch Heft 1, S. AI).
- Manuskripte (in dreifacher Ausfertigung) sind zu senden an Redaktion der Chemischen Berichte, Dr. R. Temme, Postfach 101161, D-69451 Weinheim; Telefon (06201) 606-255.
- Der Eingang der Abhandlungen wird den Autoren am Tage der Registrierung angezeigt.
- Es werden nur Manuskripte in deutscher oder englischer Sprache aufgenommen. Allen Beiträgen ist eine knappe Zusammenfassung in Englisch voranzustellen.
- Der Autor muß das alleinige Urheberrecht besitzen. Mit der Annahme des Manuskriptes durch die Redaktion überträgt er der VCH Verlagsgesellschaft das ausschließliche Nutzungsrecht, insbesondere das Recht der Vervielfältigung wie Fotokopie, Mikrofilm – oder mit irgendeinem anderen Verfahren – oder das Manuskript in eine von Maschinen, insbesondere von Datenverarbeitungsmaschinen, verwendbare Sprache zu übergeben oder zu übersetzen (auch in fremde Sprachen).
- Den Autoren werden 25 Sonderdrucke unentgeltlich portofrei zugesandt. Wünscht ein Autor mehr als 25 Abzüge, so ist dies auf dem Manuskript oder spätestens bei Rücksendung der Korrektur auf dieser zu vermerken. Den Autoren werden nur die Selbstkosten für die Zahl der die Freixemplare überschreitenden Sonderabzüge berechnet.
- Anfragen nach dem Verbleib nicht eingetreffener Berichte-Hefte oder Sonderdrucke sind zu richten an: VCH Verlagsgesellschaft mbH, Postfach 101161, D-69451 Weinheim, Telefon (06201) 606-0.


Notice to Authors

- Chemische Berichte (abbreviated Chem. Ber.; until 1946 Berichte der Deutschen Chemischen Gesellschaft, abbreviated Ber. Dtsch. Chem. Ges.) publishes articles on studies in inorganic and organometallic chemistry (Part A) as well as in organic and physical organic chemistry (Part B). In organic chemistry, emphasis is given to general synthetic methods and mechanistic problems.
- Chemische Berichte is published monthly; the annual volume concludes with a List of Authors and a List of Contents.
- Authors are solely responsible for their contributions. – The author to whom correspondence should be addressed should be indicated by an asterisk. Different addresses of coauthors should be given in lower-case letters.
- The contents of a manuscript submitted to Chemische Berichte should not have been published previously, nor submitted to any other journal, except in the form of brief preliminary communication.
- For the preparation of manuscripts, a copy of the “Instructions for Authors” is available on request (see also issue 1, p. AI).
- Manuscripts should be submitted in triplicate to: Redaktion Chemische Berichte, Dr. R. Temme, P.O. Box 101161, D-69451 Weinheim (FRG); Telephone (internat.) +49 6201 606-255.
- Acknowledgement of receipt of the manuscript will be confirmed on the day of registration.
- Manuscripts may be submitted in German or in English. All contributions must be accompanied by a summary in English.
- It is assumed that the authors concerned have the necessary authority for publication. Upon acceptance for publication, VCH Verlagsgesellschaft acquires all publishing rights including those of reprinting, translating or reproducing and distributing the manuscript in other forms (e.g. as photocopy, microfilm or in machine-readable form). Moreover, the provisions of the copyright law of the Federal Republic of Germany apply.
- The author will receive one copy of the galley proofs, together with the edited manuscript and a reprint order form; the author will receive 25 reprints free of charge and may order additional copies at cost price.
- All inquiries concerning journals or reprints should be addressed to: VCH Verlagsgesellschaft mbH, P.O. Box 101161, D-69451 Weinheim (FRG). Telephone (internat.) +49 6201 606-0.


- 3 **Sitzmann*, H., Zhou, P., Wolmershäuser, G.**
 ▶ Titan-, Zirconium- und Hafniumkomplexe mit 1,2,4-Tri-*tert*-butylcyclopentadienyl-Liganden
 Titanium, Zirconium and Hafnium Complexes with 1,2,4-Tri-*tert*-butylcyclopentadienyl Ligands




- 11 **Tremel, W.**
 $\text{Nb}_4\text{Te}_{17}\text{I}_4$, ein neues pseudo-eindimensionales Festkörper-Polytellurid
 ▶ $\text{Nb}_4\text{Te}_{17}\text{I}_4$, a New Pseudo One-Dimensional Solid-State Polytelluride




- 15 **Tremel*, W., Wortmann, U., Vomhof, T., Jeitschko, W.**
 ▶ Synthese, Struktur und elektronische Eigenschaften von $\text{Cu}_{0.69}\text{NbTe}_2$
 Synthesis, Structure and Electronic Properties of $\text{Cu}_{0.69}\text{NbTe}_2$



- 23 **Buchweitz, J., Gompper*, R., Polborn, K., Robl, C., Sailer, M.-T., Weigand*, W.**
 Funktionalisierte 1,1-Ethendithiolate als Liganden, III. – Palladium(II)- und Platin(II)-Komplexe mit Ferrocenyl-substituierten 1,1-Ethendithiolat-Liganden. Kristallstrukturanalysen von *cis*-(Ph_3P)₂M[S₂C=CH-C(O)-(η⁵-C₅H₄)Fe(η⁵-C₅H₅)] (M = Pd, Pt)
 ▶ Functionalized 1,1-Ethenedithiolates as Ligands, III. – Palladium(II) and Platinum(II) Complexes with Ferrocenyl-Substituted 1,1-Ethenedithiolate Ligands. Crystal Structure Analyses of *cis*-(Ph_3P)₂M[S₂C=CH-C(O)-(η⁵-C₅H₄)Fe(η⁵-C₅H₅)] (M = Pd, Pt)




- 27 **Werner*, H., Schäfer, M., Nürnberg, O., Wolf, J.**
 ▶ η³-Allyl- und η³-Benzyl-Rhodiumkomplexe: Synthese, Strukturdynamik und Reaktionen mit Carbonsäuren
 η³-Allyl- and η³-Benzyl Rhodium Complexes: Synthesis, Structure Dynamic, and Reactions with Carboxylic Acids




Binger*, P., Langhauser, F., Wedemann, P., Gabor, B., Mynott, R., Krüger, C.

- ▶ Allenkomplexe des Titanocens und Zirkonocens: Darstellung und Reaktivität
Allene Complexes of Titanocene and Zirconocene: Synthesis and Reactivity



Herrmann*, W. A., Kühn, F. E., Romão, C. C., Kleine, M., Mink, J.


- ▶ Mehrfachbindungen zwischen Hauptgruppenelementen und Übergangsmetallen, CXXIX. – Chlorotrioxorhenium. Neue Synthesen, Reaktionen und Derivate
Multiple Bonds between Main Group Elements and Transition Metals, CXXIX. – Chlorotrioxorhenium. Novel Syntheses, Reactions, and Derivatives



Bock*, H., Meuret, J., Näther, C., Krynitz, U.

- ▶ Strukturen sterisch überfüllter und ladungsgestörter Moleküle, 38. – Die verschiedenartigen Konformationen *N*-Trimethylsilyl-substituierter *p*-Phenyldiaminderivate: Untersuchungen an Einkristallen sowie in der Gasphase


Structures of Sterically Overcrowded and Charge-Perturbed Molecules, 38. – The Different Conformations of *N*-Trimethylsilyl-Substituted *p*-Phenylenediamine Derivatives: Investigation of Single Crystals and in the Gas Phase



Niediek, K., Neumüller*, B.

Synthese und Charakterisierung von Gallium-Phosphor-Heterocubanen


- ▶ Synthesis and Characterization of Gallium-Phosphorus Heterocubanes



73 Schmid*, G., Haske, S., Zaika, D., Boese, R., Bläser, D.

1,2-Azaboroly-Komplexe, XXIX. – Synthese und Eigenschaften von 1,2-Azaborolyboranen


- 1,2-Azaboroly Complexes, XXIX. – Synthesis and Properties of 1,2-Azaborolyboranes



81 Nöth*, H., Stolpmann, H., Thomann, M.

- Beiträge zur Chemie des Bors, 221. – Kinetische und thermodynamische Produktkontrolle bei der Addition von HX-Verbindungen an ein Amino-phosphanyl-imino-boran $R_2N-B\equiv N-PR'_2$


Contributions to the Chemistry of Boron, 221. – Kinetic and Thermodynamic Product Control in the Addition of HX Compounds to an Amino Phosphanyl-imino Borane $R_2N-B\equiv N-PR'_2$



87 Schneider, L., Englert, U., Paetzold*, P.

- Ein neuer Zugang zum ikosaedrischen *closo*-NB₁₁-Gerüst

A Novel Access to the Icosahedral *closo* NB₁₁ Skeleton




Notizen / Notes

91 Braunschweig, H., Spaniol, T., Paetzold*, P.

- Zur Bildung eines Fe₂(CO)₆-Komplexes mit (Boryloxy)allyl-Brücke


On the Formation of a Fe₂(CO)₆ Complex with a (Boryloxy)allyl Bridge



93 Meyer, F., Paetzold*, P., Englert, U.


- Reaktion von Decaboran mit dem Phosphaalkin PC*t*Bu

Reaction of Decaborane with the Phosphaalkyne PC*t*Bu



Weigand*, W., Wunsch, R., Robl, C., Amrein, W.
Metall-Komplexe von funktionalisierten schwefelhaltigen Liganden, VII. – Synthese von Platin(II)-Alkan- und -Arenthiosulfinato-Komplexen – Kristallstrukturanalyse von *N*-[(Benzylsulfinyl)thio]phthalimid

- Metal Complexes of Functionalized Sulfur-Containing Ligands, VII. – Synthesis of Platinum(II) Alkane- and Arenethiosulfinato – Complexes Crystal Structure Analysis of *N*-[(Benzylsulfinyl)thio]phthalimide




a: R = C₃H₇ b: R = C₄H₉
c: R = CH₂C₆H₅ d: R = *p*-CH₃-C₆H₄

Frey, R., Linti*, G., Polborn, K.

- Zur Chemie der Galliumverbindungen, II. – Darstellung und Struktur von Bis(2,2,6,6-tetramethylpiperidino)[tris(trimethylsilyl)silyl]gallium – ein erstes Beispiel für ein monomeres Bis(amino)gallan mit Gallium-Silicium-Bindung


On the Chemistry of Gallium Compounds, II. – Preparation and Structure of Bis(2,2,6,6-tetramethylpiperidino)[tris(trimethylsilyl)silyl]gallium – First Example for a Monomeric Bis(amino)gallane with Gallium – Silicon Bond



B Teil B: Organische Chemie / Part B: Organic Chemistry


Jutzi*, P., Mix, A., Lindermeier, T., Stammler, H.-G., Neumann, B.

- Bis(pentamethylcyclopentadienyl)methanol: Synthese, Struktur und Derivatisierung
Bis(pentamethylcyclopentadienyl)methanol: Synthesis, Structure, and Derivatization



Feucht, C., Linßen, T., Hanack*, M.


- Synthese dienophiler und enophiler Phthalocyanine
Synthesis of Dienophilic and Enophilic Phthalocyanines



Köhler*, J. E. H., Hohla, M., Richters, M., König, W. A.

Eine moleküldynamische Simulation der Komplexbildung zwischen (*R*)/(*S*)-2-Chlorpropionsäure-methylester und Heptakis(3-*O*-acetyl-2,6-di-*O*-pentyl)- β -cyclodextrin


- A Molecular-Dynamics Simulation of the Complex Formation between Methyl (*R*)/(*S*)-2-Chloropropionate and Heptakis(3-*O*-acetyl-2,6-di-*O*-pentyl)- β -cyclodextrin



127 **Kauffmann*, T., Enk, M., Fiegenbaum, P., Hansmersmann, U., Kaschube, W., Papenberg, M., Toliopoulos, E., Welke, S.**

- ▶ Organomolybdän- und Organowolfram-Reagenzien, VI. – Zur Kenntnis der carbonylmethylenierenden Molybdän-Aluminium- und Wolfram-Aluminium- μ -methylene-Komplexe


Organomolybdenum and Organotungsten Reagents, VI. – On the Carbonyl-Methylenating Molybdenum-Aluminium and Tungsten-Aluminium μ -Methylene Complexes



137 **Butenschön, H.**

Synthese von 5-Methoxy- und 5-(Dialkylamino)bicyclo[3.2.0]hept-2-en-6-on-Derivaten durch *cine*-Substitution mit Methoxid-Anionen und Dialkylaminen


- ▶ Synthesis of 5-Methoxy- and 5-(Dialkylamino)bicyclo[3.2.0]hept-2-en-6-one Derivatives by *cine* Substitution with Methoxide Anions and Dialkylamines



145 **Abele, H., Haas*, A., Lieb, M., Zwingenberger, J.**

- ▶ Darstellung geeigneter Vorstufen zur Synthese von 16,16,16,17,17,17-Hexafluorretinalen


Preparation of Suitable Precursors for the Synthesis of 16,16,16,17,17,17-Hexafluorretinals



151 **Kuck*, D., Neumann, E., Schuster, A.**

Benzo-anellierte Centropolyquinane, 14. – Synthese von Tribenzotriquinacen durch stereokontrollierte Cyclisierung von Phenyl-substituierten C_x -Diindanen (4ba,9,9aa,10-Tetrahydroindeno[1,2-*a*]indenen)


- ▶ Benzoannellated Centropolyquinanes, 14. – Synthesis of Tribenzotriquinacene by Stereocontrolled Cyclization of Phenyl-Substituted C_x -Diindans (4ba,9,9aa,10-Tetrahydroindeno[1,2-*a*]indenes)



165 **Hünig*, S., Klaunzer, N., Wenner H.**

- ▶ Stereoselektive Protonierung von Carbanionen, 2. – Diastereoselektive Protonierung von Schöllkopf-Bis-lactimether-Anionen


Stereoselective Protonation of Carbanions, 2. – Diastereoselective Protonation of Schöllkopf's Bislactim Ether Anions



173 **Maier*, G., Born, D., Bauer, I., Wolf, R., Boese, R., Cremer, D.**

- ▶ Kleine Ringe, 78. – Tri-*tert*-butyl(trimethylsilyl)cyclobutadien und Tri-*tert*-butyl(trimethylsilyl)tetrahedran


Small Rings, 78. – Tri-*tert*-butyl(trimethylsilyl)cyclobutadiene and Tri-*tert*-butyl(trimethylsilyl)tetrahedrane




191 **Maier*, G., Wolf, R., Kalinowski, H.-O., Boese, R.**

- ▶ Kleine Ringe, 79. – Synthese und Eigenschaften neuer Silyl-substituierter Cyclobutadiene und Tetrahedrane


Small Rings, 79. – Synthesis and Properties of Novel Silyl-Substituted Cyclobutadienes and Tetrahedranes




- 201 **Maier*, G., Wolf, R., Kalinowski, H.-O.**
 ▶ Kleine Ringe, 80. – Tri-*tert*-butyl(trimethylgermyl)cyclobutadien und Tri-*tert*-butyl(trimethylgermyl)tetrahedran
 Small Rings, 80. – Tri-*tert*-butyl(trimethylgermyl)cyclobutadiene and Tri-*tert*-butyl(trimethylgermyl)tetrahedrane




- 205 **Dau-Schmidt, J.-P., Mayr*, H.**
 Relative Reaktivitäten von Alkylchloriden unter Friedel-Crafts-Reaktionsbedingungen
 ▶ Relative Reactivities of Alkyl Chlorides under Friedel-Crafts Conditions




- 213 **Mayr*, H., Dau-Schmidt, J.-P.**
 Relative Reaktivitäten von Acetalen und Ethern unter Friedel-Crafts-Reaktionsbedingungen
 ▶ Relative Reactivities of Acetals and Ethers under Friedel-Crafts Conditions




- 219 **Boldt*, P., Döring, D., Jones, P. G., Köpper, H., Trog, R.-S.**
 ▶ (1 α ,6 α ,7 α ,8 α ,12 α ,17 α ,18 α ,19 α)-Heptacyclo[17.3.1.1^{8,12}.0^{5,23}.0^{6,18}.0^{7,17}.0^{16,24}]tetracosa-5(23),16(24)-dien, ein hyperstabiles Alken
 (1 α ,6 α ,7 α ,8 α ,12 α ,17 α ,18 α ,19 α)-Heptacyclo[17.3.1.1^{8,12}.0^{5,23}.0^{6,18}.0^{7,17}.0^{16,24}]tetracosa-5(23),16(24)-diene, a Hyperstable Alkene






- 223 **Staab*, H. A., Carell, T., Döhling, A.**
 Photoinduzierter Elektronen-Transfer in Porphyrin-Chinon-Cyclophanen, 9. – Synthese und Eigenschaften neuer Chinon-verbrückter Diphenyl- und Tetraphenylporphyrine
 ▶ Photoinduced Electron Transfer in Porphyrin-Quinone Cyclophanes, 9. – Syntheses and Properties of New Quinone-Bridged Diphenyl- and Tetraphenylporphyrins





- 231 **Staab*, H. A., Krieger, C., Anders, C., Rückemann, A.**
 Photoinduzierter Elektronen-Transfer in Porphyrin-Chinon-Cyclophanen, 10. – Zink-Komplexe von Porphyrin-Chinon-Cyclophanen: Synthesen, Strukturen und Elektronen-Transfer-bezogene Eigenschaften
 ▶ Photoinduced Electron Transfer in Porphyrin-Quinone Cyclophanes, 10. – Zinc Complexes of Porphyrin-Quinone Cyclophanes: Syntheses, Structures and Electron-Transfer-Related Properties





Kurzmitteilungen / Short Communications



Relative Reactivities of Acetals and Ethers under Friedel-Crafts Conditions

Herbert Mayr^{*a} and Jan-Peter Dau-Schmidt^b

Institut für Organische Chemie, Technische Hochschule Darmstadt^a,
Petersenstraße 22, 64287 Darmstadt, F.R.G.

Institut für Chemie, Medizinische Universität zu Lübeck^b,
Ratzeburger Allee 160, 23538 Lübeck, F.R.G.


Received August 2, 1993

Key Words: Alkylation / Alkoxyalkylation / Carboxonium ions / Acetals


Competition experiments have been performed to determine the relative reactivities of acetals and ethers toward allyltrimethylsilane in the presence of catalytic amounts of $\text{BF}_3 \cdot \text{OEt}_2$. It is found that acetals $\text{R}-\text{CH}(\text{OMe})_2$ and their phenyl-

ogous *p*-anisyl ethers $\text{R}-\text{CH}(p\text{-MeOC}_6\text{H}_4)(\text{OMe})$ show very little differences in reactivity. The reactivity scales are employed to rationalize the results of Lewis acid-catalyzed additions of acetals and ethers to CC double bonds.

In the preceding article, we have shown that the electrophilicities of alkyl chlorides in the presence of catalytic amounts of Lewis acids are proportional to their solvolysis rates in ethanol^[1]. This correlation represents a quantitative basis for our working hypothesis that alkylations according to eq. (1) can only give 1:1 adducts if the reactants RCl solvolyze faster than the products. Since solvolysis rate constants for most types of alkyl chlorides are known, the synthetic potential of reaction (1) can easily be predicted.





An analogous analysis of the outcome of acetal and ortho ester additions is not possible, as their solvolyses require acid catalysis, and the rate constants for the proton-catalyzed hydrolysis are not only a function of the Lewis acidity of the intermediate carboxonium ions. We had, therefore, investigated the relative reactivities of acetals and ortho esters towards methyl vinyl ether in the presence of $\text{BF}_3 \cdot \text{OEt}_2$ by competition experiments^[2]. The reactivity scale, thus obtained, allowed the rationalization of the large manifold of alkoxyalkylations of vinyl ethers (eq. 2)^[2].



Since this reactivity series was restricted to acetals and ortho esters and did not include ethers, it did not allow predictions whether Lewis acid-catalyzed reactions of acetals with arenes yield benzyl ethers or 2:1 products arising from reactions with two arene molecules (eq. 3) and

whether it was possible to produce acetals by reactions of ordinary ethers with enol ethers (eq. 4).




Results

The acetals and ethers listed in Table 1 react with allyltrimethylsilane in the presence of $\text{BF}_3 \cdot \text{OEt}_2$ to give 51–88% (isolated yield) of the expected $\text{S}_{\text{E}}2'$ products, some of which have been described in ref.^[1,3] Competition experiments, as described for alkyl chlorides in the preceding article^[1], have been used to determine the relative reactivities of acetals and ethers in presence of $\text{BF}_3 \cdot \text{OEt}_2$ (Scheme 1).

Table 1 shows that aromatic acetals and benzhydryl ethers possess comparable reactivities. Since not all competition constants given in Table 1 are linked with each other, they can not be combined to give a single reactivity scale. This goal can be achieved by relating these data to the rela-

Scheme 1



tive reactivities of alkyl chlorides from ref.^[1] as described below.

Table 1. Competition constants for the $\text{BF}_3 \cdot \text{OEt}_2$ - catalyzed reactions of acetals and ethers with allyltrimethylsilane in CH_2Cl_2 at -70°C

| Reactant A | Reactant B | $\kappa_{\text{A/B}}$ |
|---|---|---------------------------|
| 1 (p-MeOC ₆ H ₄)CH(OMe) ₂ | 2 (p-MeOC ₆ H ₄) ₂ CHOMe | 1.5 ± 0.1 |
| 1 (p-MeOC ₆ H ₄)CH(OMe) ₂ | 3 (p-MeC ₆ H ₄)CH(OMe) ₂ | 3.4 ± 0.1 |
| 2 (p-MeOC ₆ H ₄) ₂ CHOMe | 3 (p-MeC ₆ H ₄)CH(OMe) ₂ | 2.3 ± 0.1 |
| 3 (p-MeC ₆ H ₄)CH(OMe) ₂ | 4 C ₆ H ₅ CH(OMe) ₂ | 4.3 ± 0.3 |
| 3 (p-MeC ₆ H ₄)CH(OMe) ₂ | 4 C ₆ H ₅ CH(OMe) ₂ | 3.1 ± 0.2 ^[a] |
| 5 (p-ClC ₆ H ₄)CH(OMe) ₂ | 6 (p-MeC ₆ H ₄) ₂ CHOMe | 9.5 ± 0.3 |
| 6 (p-MeC ₆ H ₄) ₂ CHOMe | 9 (p-MeC ₆ H ₄)(C ₆ H ₅)CHOMe | 12.0 ± 0.6 ^[a] |
| 7 n-C ₃ H ₇ -CH(OMe) ₂ | 8 (CH ₂) ₂ -CH=CH-CHOMe | 5.6 ± 0.1 |

^[a] Lewis acid $\text{ZnCl}_2 \cdot \text{OEt}_2$ instead of $\text{BF}_3 \cdot \text{OEt}_2$.

Let us assume that the reactions described in Scheme 1 involve a rapid preequilibrium^[5]. As discussed in ref.^[1], the competition constant is then given by eq. (5), where equilibrium and rate constants have the meaning defined in Scheme 1.

$$\kappa_{\text{AB}} = \frac{K_{1\text{A}}k_{2\text{A}}}{K_{1\text{B}}k_{2\text{B}}} \quad (5)$$

Since in several examples the rate constants for the reactions of carbenium ions with π nucleophiles ($k_{2\text{A}}$, $k_{2\text{B}}$) have been found to be independent of the counterion^[4], $K_{1\text{A}}$ and $K_{1\text{B}}$ are the only quantities in eq. (5) that are leaving group-dependent. As the ratios of two ionization constants ($K_{1\text{A}}/K_{1\text{B}}$) have been shown to be leaving group-independent if variable steric or anomeric effects are absent^[7], κ_{AB} will be approximately equal for corresponding pairs of alkyl chlorides and methyl ethers. This conclusion is corroborated by comparison of the entry before last in Table 1 ($\kappa_{\text{AB}} = 12.0$) with the relative reactivity of the corresponding benzhydryl chlorides ($\kappa_{\text{AB}} = 14.2$ ^[1]). By assuming the relative reactivities of (p-MeOC₆H₄)₂CHOMe/(p-MeC₆H₄)₂CHOMe and (p-MeC₆H₄)₂CHOMe/3-methoxycyclopentene to be 126 and 22.1, respectively, as determined for the corresponding

Scheme 2. Relative reactivities of acetals and ethers towards allyltrimethylsilane (CH_2Cl_2 , -70°C)

| Compound | k_{rel} |
|---|------------------------|
| 1 (p-MeOC ₆ H ₄)CH(OMe) ₂ | 4.19 × 10 ³ |
| 2 (p-MeOC ₆ H ₄) ₂ CHOMe | 2.79 × 10 ³ |
| 3 (p-MeC ₆ H ₄)CH(OMe) ₂ | 1.27 × 10 ³ |
| 4 C ₆ H ₅ CH(OMe) ₂ | 3.02 × 10 ² |
| 5 (p-ClC ₆ H ₄)CH(OMe) ₂ | 2.08 × 10 ² |
| 6 (p-MeC ₆ H ₄) ₂ CHOMe | 2.21 × 10 ¹ |
| 7 n-C ₃ H ₇ -CH(OMe) ₂ | 5.60 |
| 8 (CH ₂) ₂ -CH=CH-CHOMe | 1 |

^[a] Relative reactivities of the corresponding chloro compounds; see text.




Figure 1. Correlation of the relative reactivities of acetals towards allyltrimethylsilane and methyl vinyl ether ($\text{BF}_3 \cdot \text{OEt}_2$, CH_2Cl_2 , -70°C)

chloro compounds^[1], the data of Table 1 can be connected to give the reactivity scale of Scheme 2.

The relative reactivities of the acetals listed in Scheme 2 can be compared with the relative reactivities of these compounds toward methyl vinyl ether as reference nucleophile^[2]. Figure 1 shows a fairly good correlation between the two sets of data, and the correlation equation (6) indicates that variation of the acetal has a greater effect on the reactivity toward methyl vinyl ether than on the reactivity toward allyltrimethylsilane.

$$\log k_{\text{rel}} (\text{allyltrimethylsilane}) = 0.687 \log k_{\text{rel}} (\text{methyl vinyl ether}) + 0.49 \quad (6)$$

Equation (6) now allows to adjust von der Brüggen's reactivity scale for acetals toward methyl vinyl ether^[2] to the

scale of Scheme 2. Combination with some additional k_{rel} values for alkyl chlorides^[1] as discussed above yields the electrophilicity scale of Figure 2. Because of the approximations necessary for its construction, this scale is less precise than that presented for alkyl chlorides in the preceding paper^[1]. It may be more valuable for the synthetic chemist, however, because this reactivity order cannot be derived from readily available data (as solvolysis rates in the case of alkyl chlorides).




Figure 2. Relative reactivities of ethers and acetals towards allyltrimethylsilane (CH_2Cl_2 , -70°C)

Consequences for Organic Synthesis

The relative reactivities of acetals in Figure 2 have already been used to explain, why the reactions of benzaldehyde acetals and of α,β -unsaturated acetals with alkyl vinyl ethers give better yields of 1:1 products than the corresponding reactions of saturated acetals^[2]. Similarly, we have rationalized why only methoxy-substituted benzaldehyde acetals and α,β -unsaturated acetals had been reported to give 1:1 products with 1-alkoxy-1,3-butadienes^[2].

The reactions of benzaldehyde dimethyl acetal with isobutene or styrene give products which incorporate a tert. alkyl ether and/or an α -phenyl alkyl ether substructure (Scheme 3). The isolation of such products is possible^[8]

since tert. alkyl ethers and α -phenyl alkyl ethers are even less reactive than the least reactive compounds depicted in Figure 2^[1]. For the same reason, homoallylic ethers are produced in high yields from acetals and allylsilanes by the Hosomi-Sakurai reaction^[9].

Scheme 3





Figure 2 suggests dianisylmethyl- and ditolylmethyl methyl ether to be more reactive than saturated acetals. Consequently, we have now succeeded to combine these ethers with methyl vinyl ether to produce the acetals as shown in Scheme 4.

Scheme 4



In contrast, the corresponding reaction of methyl phenyl-*p*-tolylmethyl ether with methyl vinyl ether gave polymeric vinyl ether predominantly (^1H NMR). In accord with Figure 2, the resulting saturated acetals are more reactive than methyl phenyl-*p*-tolylmethyl ether, and the 1:1 products are, therefore, not isolable.

The reaction of *p*-anisaldehyde acetal with anisole gives a benzhydryl ether which is predicted by Figure 2 to possess a similar reactivity as the reactant acetal. The corresponding reaction gave only tris(*p*-anisyl)methane, and we have not been able to isolate the intermediate bis(*p*-anisyl)methyl methyl ether. On the other hand, Figure 2 predicts that a benzhydryl ether should be isolable from the reaction of benzaldehyde acetal with toluene. Our attempts to realize this reaction failed, however; disproportionation of the acetal turned out to be faster than the electrophilic aromatic substitution, probably because of the low nucleophilicity of toluene.

Conclusion

The reactivity scale shown in Figure 2 may serve as a guide for rationalizing and designing acetal and ether additions to CC double-bonded compounds. It should be noted, however, that it is not applicable to reactions with silylated enol ethers since in these cases rapid desilylation of the 1:1 adducts prevents sequential reactions and the formation of higher adducts.

We thank Dr. B. Dogan for critical comments and Dipl.-Ing. M. Hartnagel for experimental assistance. Financial support by the Deutsche Forschungsgemeinschaft and the Fonds der Chemischen Industrie is gratefully acknowledged.

Experimental

General: See ref.^[1]. — **Reaction products:** The products obtained by reaction of diarylmethoxymethanes and allyltrimethylsilane are identical to those formed by the analogous reactions with diarylchloromethanes and have been described in ref.^[1]

For the reactions of aromatic acetals with allyltrimethylsilane, the procedure described in ref.^[3] has been used: A solution of the acetal (2.00 g) in CH₂Cl₂ (4.5 ml) was added dropwise (15 min) to a cooled (−78°C, N₂ protection) mixture of allyltrimethylsilane (1.5 equivalents), CH₂Cl₂ (1 ml), and trimethylsilyl triflate (0.15 equivalents). After completion of the reaction the mixture was washed with an equal volume of concd. aqueous ammonia. The phases were separated, the organic layer was dried with CaCl₂ and evaporated.

Reaction with benzaldehyde dimethyl acetal (4): See ref.^[3,10].

p-Anisaldehyde Dimethyl Acetal (1) (2.00 g, 11.0 mmol) and allyltrimethylsilane (1.88 g, 16.5 mmol) reacted within 2 h under the conditions described above to give 1.38 g (65%) of 4-methoxy-4-(4-methoxyphenyl)-1-butene with b.p. 58°C (bath)/0.15–0.2 mbar. — ¹H NMR (90 MHz, CDCl₃): δ = 2.47 (mc, 2H, 3-H), 3.20 (s, 3H, 4-OCH₃), 3.82 (s, 3H, Ar-OCH₃), 4.12 (t, J = 8 Hz, 1H, 4-H), 4.88–5.18 (m, 2H, =CH₂), 5.51–6.03 (m, 1H, =CH–), 6.86, 7.22 (AA'BB' system, J_{AB} = 8 Hz, 4H, aromatic H). — The ¹H-NMR spectrum is in accord with the data in ref.^[10].

p-Tolylaldehyde Dimethyl Acetal (3) (2.00 g, 12.0 mmol) and allyltrimethylsilane (2.05 g, 18.0 mmol) reacted within 2 h under the conditions described above to give 1.70 g (80%) of 4-methoxy-4-(4-methylphenyl)-1-butene with b.p. 60–63°C (bath)/1 mbar. — ¹H NMR (90 MHz, CCl₄): δ = 2.30 (s, 3H, CH₃), 2.37 (mc, 2H, 3-H, superimposed by the singlet at 2.30), 3.12 (s, 3H, 4-OCH₃), 4.00 (t, J = 8 Hz, 1H, 4-H), 4.78–5.12 (m, 2H, =CH₂), 5.43–5.96 (m, 1H, =CH–), 7.05 (s, 4H, aromatic H).

4-Chlorobenzaldehyde Dimethyl Acetal (5) (2.00 g, 10.7 mmol) and allyltrimethylsilane (1.83 g, 16.0 mmol) reacted within 3 h under the conditions described above to give 1.84 g (88%) of 4-(4-chlorophenyl)-4-methoxy-1-butene with b.p. 76°C (bath)/1.5 mbar. — ¹H NMR (90 MHz, CDCl₃): δ = 2.48 (mc, 2H, 3-H), 3.25 (s, 3H, 4-OCH₃), 4.18 (t, J = 8 Hz, 1H, 4-H), 4.90–5.20 (m, 2H, =CH₂), 5.51–6.03 (m, 1H, =CH–), 7.28 (mc, 4H, aromatic H).

4-Methoxy-1-heptene was synthesized from butyraldehyde dimethyl acetal (7) and allyltripropylsilane (instead of allyltrimethylsilane) to facilitate distillative workup of the reaction mixture (see ref.^[1]): After dropwise addition (15 min) of 7 (1.14 g, 9.65 mmol) to a cooled (−78°C, N₂ protection) solution of TiCl₄ (1.07 ml, 9.76 mmol) in CH₂Cl₂ (30.0 ml), a mixture of allyltripropylsilane (2.12 g, 10.7 mmol) and CH₂Cl₂ (15 ml) was added dropwise within 1 h. The mixture was allowed to react for 3 h and was then washed with

5 N HCl (60 ml). The phases were separated, and the organic layer was dried with CaCl₂/K₂CO₃ and evaporated. Twofold distillation yielded 680 mg (55%) of 4-methoxy-1-heptene with b.p. 65°C (bath)/91 mbar. — ¹H-NMR (200 MHz, CDCl₃): δ = 0.86–0.99 (m, 3H, 7-H), 1.26–1.57 (m, 4H, 5-H, 6-H), 2.22–2.34 (m, 2H, 3-H), 3.14–3.31 (m, 1H, 4-H), 3.35 (s, 3H, OCH₃), 5.02–5.13 (m, 2H, =CH₂), 5.72–5.94 (m, 1H, =CH–). — ¹³C NMR (50 MHz, CDCl₃): δ = 14.21 (q, C-7), 18.52 (t, C-6), 35.63, 37.78 (2 t, C-3, C-5), 56.55 (q, OCH₃), 80.27 (d, C-4), 116.73 (t, =CH₂), 135.00 (d, =CH–). — MS (70 eV), m/z (%) = 87 (56), 85 (19), 55 (30), 45 (100), 41 (29). The M⁺ peak could not be detected.

Methoxybis(4-methoxyphenyl)methane (2) and Methyl Vinyl Ether: Compound 2 (260 mg, 1.01 mmol) was dissolved in anhydrous CH₂Cl₂ (10 ml) and the solution cooled at −78°C (N₂ protection). After addition of BF₃ · OEt₂ (50 μl, 0.4 mmol), a solution of methyl vinyl ether (92.8 mg, 1.65 mmol) in CH₂Cl₂ (25 ml) was added dropwise within 30 min. After 24 h the mixture was washed with an equal volume of dil. aqueous ammonia. The phases were separated, the organic layer was dried with CaCl₂ and the solvent evaporated in vacuo. Filtration over Al₂O₃ (neutral) with hexane/ether (80:20) and removal of solvents (0.1 mbar) yielded 160 mg (51%) of 3,3-bis(4-methoxyphenyl)propionaldehyde dimethyl acetal as a viscous oil. — ¹H NMR (300 MHz, CDCl₃): δ = 2.27 (dd, J₁₂ = 5.9, J₂₃ = 8.0 Hz, 2H, 2-H), 3.27 (s, 6H, 1-OCH₃), 3.72 (s, 6H, aryl-OCH₃), 3.99 (t, J₂₃ = 8.0 Hz, 1H, 3-H), 4.15 (t, J₁₂ = 5.9 Hz, 1H, 1-H), 6.80, 7.14 (AA'BB' system, J_{AB} = 8.7 Hz, 8H, aromatic H). — ¹³C NMR (75 MHz, CDCl₃): δ = 37.61 (t, C-2), 44.02 (d, C-3), 51.67 (q, OCH₃), 54.13 (q, aryl-OCH₃), 101.84 (d, C-1), 112.88 (d, C_m), 127.59 (d, C_o), 135.93 (s, C_i), 156.96 (s, C_p). — C₁₉H₂₄O₄ (316.4): calcd. C 72.13, H 7.65; found C 71.75, H 7.58.

Methoxybis(4-methylphenyl)methane (6) (230 mg, 1.02 mmol) and methyl vinyl ether (92.8 mg, 1.60 mmol) were combined analogously to yield 180 mg (62%) of 3,3-bis(4-methylphenyl)propionaldehyde dimethyl acetal as a viscous oil. — ¹H NMR (300 MHz, CDCl₃): δ = 2.27 (s, 6H, CH₃), 2.30 (dd, partially masked, J₁₂ = 5.9, J₂₃ = 8.0 Hz, 2H, 2-H), 3.27 (s, 6H, OCH₃), 4.02 (t, J₂₃ = 8.0 Hz, 1H, 3-H), 4.17 (t, J₁₂ = 5.9 Hz, 1H, 1-H), 7.06, 7.13 (AA'BB'

Table 2. Determination of the competition constants in CH₂Cl₂ (10 ml) at −70°C

| R _A Ome / mg | R _B Ome / mg | AS ^[a] / mg | [LA]/[R _A Ome] ^[b] (molar ratio) | [P _A]/[P _B] (molar ratio) | Time | Standard ^[c] | κ _{A/B} |
|-------------------------|-------------------------|------------------------|--|---|--------|-------------------------|------------------|
| 1 35.9 | 2 24.0 | 57 | 0.60 | 3.00 | 61 min | PPR | 1.4 |
| | 51.6 | 57 | 0.60 | 1.02 | 61 min | PPR | 1.6 |
| 1 21.3 | 3 19.2 | 57 | 0.84 | 3.75 | 61 min | PPR | 3.45 |
| | 39.6 | 57 | 0.84 | 1.34 | 61 min | PPR | 3.35 |
| 2 22.2 | 3 74.5 | 57 | 1.46 | 0.40 | 61 min | PPR | 2.3 |
| | 168.6 | 57 | 1.46 | 0.18 | 61 min | PPR | 2.2 |
| 3 23.6 | 4 104.1 | 57 | 1.12 | 0.65 | 61 min | PPT | 4.0 |
| | 161.3 | 57 | 1.12 | 0.51 | 61 min | PPT | 4.5 |
| 3 21.6 | 4 118.6 | 5.7 | 1.60 ^[d,e] | 0.46 | 63 h | PPT | 2.9 |
| | 204.1 | 5.7 | 1.60 ^[d,e] | 0.32 | 63 h | PPT | 3.3 |
| 5 21.2 | 6 22.0 | 57 | 1.22 | 8.30 | 61 min | PPT | 9.2 |
| | 84.1 | 57 | 1.22 | 2.35 | 61 min | PPT | 10.0 |
| | 167.8 | 57 | 1.22 | 1.42 | 61 min | PPT | 9.2 |
| 6 23.6 | 9 106.8 | 5.7 | 1.60 ^[d,e] | — | 90 h | PO | 11.4 |
| | 195.1 | 5.7 | 1.60 ^[d,e] | — | 90 h | PO | 12.6 |
| 7 17.1 | 8 15.9 | 358 | 5.50 | 5.02 | 60 min | C | 5.8 |
| | 35.0 | 358 | 5.50 | 2.50 | 60 min | C | 5.5 |
| | 61.6 | 358 | 5.50 | 1.16 | 60 min | C | 5.5 |

[a] AS = allyltrimethylsilane. — [b] Lewis acid (LA) is BF₃ · OEt₂. — [c] HPLC analysis apart from the 7/8 couple which was analyzed by GC. Standards: C = cumene, PO = 1-phenyloctane, PPR = 1-phenylpropane, PPT = 1-phenylpentane. — [d] 5 ml of CH₂Cl₂. — [e] Lewis acid is ZnCl₂ · (OEt₂)_{1.03}.

system, $J_{AB} = 8.1$ Hz, 8H, aromatic H). – ^{13}C NMR (75 MHz, CDCl_3): $\delta = 21.11$ (q, CH_3), 38.45 (t, C-2), 46.05 (d, C-3), 52.81 (q, OCH_3), 102.92 (d, C-1), 127.85 (d, C_o), 129.35 (d, C_m), 135.78 (s, C_p), 141.81 (s, C_i). – $\text{C}_{19}\text{H}_{24}\text{O}_2$ (284.4): calcd. C 80.24, H 8.51; found C 80.12, H 8.63.

Competition Experiments have been performed as described in ref.^[1] by using $\text{BF}_3 \cdot \text{OEt}_2$ instead of ZnCl_2 .

- [1] J.-P. Dau-Schmidt, H. Mayr, *Chem. Ber.* **1993**, *126*, 205–212, preceding paper.
 [2] U. von der Brüggen, R. Lammers, H. Mayr, *J. Org. Chem.* **1988**, *53*, 2920–2925.
 [3] R. Noyori, S. Murata, M. Suzuki, *Tetrahedron* **1981**, *37*, 3899–3910.
 [4] ^[4a] H. Mayr, R. Schneider, C. Schade, J. Bartl, R. Bederke, *J.*

- Am. Chem. Soc.* **1990**, *112*, 4446–4454. – ^[4b] G. Hagen, H. Mayr, *J. Am. Chem. Soc.* **1991**, *113*, 4954–4961.
 [5] An alternative mechanism, attack of the allylsilane at an acetal-Lewis acid complex, which has been observed for related reactions^[6], has not been considered for the same reasons discussed in the preceding article^[1].
 [6] ^[6a] S. E. Denmark, T. M. Willson, *J. Am. Chem. Soc.* **1989**, *111*, 3475–3476. – ^[6b] S. E. Denmark, N. G. Almstead, *J. Am. Chem. Soc.* **1991**, *113*, 8089–8110.
 [7] S. V. McKinley, J. W. Rakshys, Jr., A. E. Young, H. H. Freedman, *J. Am. Chem. Soc.* **1971**, *93*, 4715–4724.
 [8] ^[8a] O. C. Dermer, J. J. Hawkins, *J. Am. Chem. Soc.* **1952**, *74*, 4595–4597. – ^[8b] J. W. Copenhaver (General Aniline & Film Corp.), U.S. Pat. 2,768,212, **1956**; *Chem. Abstr.* **1958**, *52*, 433a. – ^[8c] R. Paul, S. Tchelitcheff, *Bull. Soc. Chim. Fr.* **1951**, 125–129.
 [9] A. Hosomi, M. Endo, H. Sakurai, *Chem. Lett.* **1976**, 941–942.
 [10] S. Torii, T. Inokuchi, S. Takagishi, H. Horike, H. Kuroda, K. Uneyama, *Bull. Chem. Soc. Jpn.* **1987**, *60*, 2173–2188.

[255/93]