



Staphylococcus aureus Population Structure and Genomic Profiles in Asymptomatic Carriers in Switzerland

Danai Etter^{1,2*}, Sabrina Corti¹, Simona Spirig¹, Nicole Cernela¹, Roger Stephan¹ and Sophia Johler¹

¹ Institute for Food Safety and Hygiene, Vetsuisse Faculty, University of Zurich, Zurich, Switzerland, ² Institute of Food, Nutrition and Health, Department of Health Science and Technology, ETH Zürich, Zurich, Switzerland

OPEN ACCESS

Edited by:

Mattias Collin,
Lund University, Sweden

Reviewed by:

Frieder Schaumburg,
University of Münster, Germany
Peter Kinnevey,
Dublin Dental University Hospital,
Ireland

*Correspondence:

Danai Etter
danai.etter@hest.ethz.ch

Specialty section:

This article was submitted to
Infectious Diseases,
a section of the journal
Frontiers in Microbiology

Received: 06 April 2020

Accepted: 20 May 2020

Published: 24 June 2020

Citation:

Etter D, Corti S, Spirig S,
Cernela N, Stephan R and Johler S
(2020) Staphylococcus aureus
Population Structure and Genomic
Profiles in Asymptomatic Carriers
in Switzerland.
Front. Microbiol. 11:1289.
doi: 10.3389/fmicb.2020.01289

Staphylococcus aureus is a leading cause for clinical infections and food intoxications, causing over 100,000 yearly cases of bacteremia in the United States and 434 food-borne outbreaks in the European Union. Approximately 30% of the population permanently carry *S. aureus* asymptotically in their nasal cavity. The risk of infection and transmission to food items or the environment is higher in individuals that are nasally colonized. In addition, *S. aureus* can acquire various antimicrobial resistances leading to therapeutic failure, additional medical costs, and fatalities. Methicillin-resistant *S. aureus* (MRSA) cause a considerable burden of disease in humans and animals. MRSA carriage has been associated with animal and in particular livestock contact. Extensive current data on the virulence gene profiles, as well as data on antimicrobial resistance determinants is crucial in developing effective strategies to mitigate the burden of disease. To this end, we screened the anterior nares of 160 test subjects (87 pupils and 73 members of farmer families) in Switzerland for *S. aureus* carriage. A total of 73 *S. aureus* isolates were obtained. Factors such as exposure to farm or companion animals and personal medical history were recorded using a questionnaire. Using a DNA microarray, isolates were assigned to clonal complexes (CCs), and virulence and resistance gene profiles were determined. The collected strains were assigned to 20 CCs, among others CC1, CC7, CC8, CC15, CC30, CC45, CC97, and CC398. Two MRSA strains and one multiresistant isolate carrying genes *blaZ//R*, *InuA*, *aadD*, *tetK*, and *fosB* were isolated from farmers with intensive exposure to animals. Strains carrying *pvl*, causing severe skin lesions and necrotizing pneumonia, as well as tetracycline, erythromycin, and kanamycin resistance genes were found in individuals that had taken antibiotics during the last year. A variety of superantigenic toxin genes was detected, including among others, the toxic shock syndrome toxin (*tst1*), and various enterotoxins (*sea*, *sec*, *sel*, and the *egc* cluster). Contact to chickens was identified as a significant factor contributing to *S. aureus* colonization.

Keywords: nasal colonization, MRSA, *pvl*, farmers, livestock, community, hospital, companion animals

INTRODUCTION

Staphylococcus aureus is a human pathogen, resulting in infections and foodborne intoxications. It caused over 100,000 cases of bacteremia in the United States in 2017 and 434 food-borne outbreaks in the European Union in 2015 (Tong et al., 2015; EFSA, 2016; Kourtis et al., 2019). Furthermore, it can colonize human anterior nares as a commensal bacterium. Around 20–30% of the population permanently carry *S. aureus* asymptomatically in their nose (Sakr et al., 2018), another 20–60% can be intermittent carriers (Kluytmans et al., 1997; Wertheim et al., 2005). Initial colonization may take place during the first days of life via transmission from mother to child (Maayan-Metzger et al., 2017). After birth the hands have been identified as the main source of transmission for the pathogen to colonize the nose (Wertheim et al., 2005). The nasal habitat may act as a reservoir for pathogen spread and can pose a threat to the carrier itself (Nouwen et al., 2006). An elevated risk of infection associated with persistent nasal carriage of *S. aureus* in surgical or dialysis patients emphasizes the importance of characterizing nasal colonization and associated risk factors (Kluytmans et al., 1997; Nouwen et al., 2006). In addition, antimicrobial resistance among *S. aureus* is rapidly emerging globally and rendering treatment of chronic and acute *S. aureus* infections increasingly difficult. The alarming prospect of a post-antibiotic era highlights the importance of identifying new therapeutic agents (Johler, 2017; Vestergaard et al., 2019). Specific virulence factors and/or master virulence regulators have been suggested as promising targets (Maura et al., 2016; Ansari et al., 2019). In particular, methicillin-resistant *S. aureus* (MRSA) have been studied in recent years with regard to their transmission dynamics (Christopher et al., 2011; Davis et al., 2012). Forms of transmission include hospital-acquired, community-acquired, and livestock-acquired with the MRSA in question being designated as HA-MRSA, CA-MRSA, and LA-MRSA, respectively (Deurenberg and Stobberingh, 2008; Knox et al., 2015; Smith, 2015; Kwok et al., 2018). Intensive contact to animals has been shown to increase the risk of LA-MRSA carriage among farmers, veterinarians, and animal owners (Graveland et al., 2011; Davis et al., 2012; Fluit, 2012). Several studies have investigated *S. aureus* colonization of veterinarians (Huber et al., 2011; Rosenkranz Wettstein et al., 2014), on-farm and food-chain epidemiology (Kraemer et al., 2017; Leuenberger et al., 2019; Morach et al., 2019), or colonization in the general population in Switzerland (Mégevand et al., 2010; Olearo et al., 2016). Here, we aim to provide insight into colonization patterns in asymptomatic carriers with varying animal exposure. We screened 160 individuals for nasal *S. aureus* and collected data on animal contact and medical history using a questionnaire. The obtained isolates were characterized by DNA microarray, which allows for assignment of clonal complexes (CCs), and determines the presence of resistance and virulence genes.

MATERIALS AND METHODS

Questionnaire

A questionnaire was used to collect metadata. We assessed exposure of participants to farm and companion animals, medical history including antibiotic prescriptions and hospital or doctor visits. Answers concerning exposure to animals ranged from never to daily (0, 1, 2, 3, 4, 5). Medical history was recorded to assess, whether the individual had been to a doctor/hospital and/or received antibiotics in the last year (0/1). Raw data can be examined in **Supplementary Table S2**.

Sampling, Bacterial Isolation, Species Identification, and DNA Extraction

Each volunteer (87 pupils and 73 members of farmer families) provided a written declaration of consent and took his/her own nasal swab after prior demonstration of the sampling procedure. All nasal swabs were streaked on EASY Staph[®] plates (BIOKAR Diagnostics, Beauvais, France) and incubated at 37°C for 48 h. Presumptive *S. aureus* colonies were confirmed using matrix assisted laser desorption/ionization—time of flight mass spectrometry (MALDI-TOF MS, Bruker, Fällanden, Switzerland). After cell lysis with Lysostaphin (Sigma–Aldrich, Buchs, Schweiz), DNA was extracted with the Blood & Tissue Kit (Qiagen, Hombrechtikon, Switzerland).

DNA Microarray Analysis

DNA microarray was performed using Staphytype genotyping kit 2.0 (Alere, Wädenswil, Switzerland) following the manufacturer's instructions. An ArrayMate reader (Alere) was used for signal acquisition. In addition, the similarity of the virulence and resistance gene profiles was visualized using SplitsTree⁴ (Huson and Bryant, 2006) as previously described (Wattlinger et al., 2012). Readouts can be found in **Supplementary Table S1**.

Statistical Analysis

Binomial and multinomial logistic regression analyses were performed to ascertain the effects of age, gender, profession, medical history, and exposure to farm and companion animals on the likelihood that participants are colonized with *S. aureus*, with certain CCs or harbor MRSA strains or enterotoxin gene carrying isolates (SPSS, v26.0.0.0). *P*-values < 0.05 were considered significant (**Supplementary Table S4**).

RESULTS

S. aureus Prevalence and Clonal Complexes in Sampled Populations

Of the 160 nasal swabs from asymptomatic carriers 46% ($n = 73$, confidence interval = 7.72) were positive for *S. aureus*. In the sampled 87 pupils and 73 family members

¹<http://www.splittree.org/>

of farmers colonization rates slightly differed with 44% in pupils and 48% in family members of farmers. The most common CCs for pupils were CC45 ($n = 11$) and CC15 ($n = 10$), while for family members of farmers, it were mainly CC30 ($n = 9$) and CC45 ($n = 5$) (Table 1). Some CC were exclusively found in one of the two groups, e.g., CC1, CC22, CC101, CC121, CC182, CC361, and CC509 in family members of farmers, while CC6, CC12, CC152, CC188, and CC398 were only found in pupils (Figure 1). All other strains belonged to CC5, CC7, CC8, CC59, and CC97.

Exposure to Animals and Medical History of Participants

Overall animal exposure varied greatly between the two populations. Family members of farmers had an average exposure value of 2.4, corresponding to monthly-to-weekly exposure. Pupils on the other hand only reached a value of 0.8, corresponding to never-to-yearly exposure. Almost all family members of farmers were in contact with cattle on a daily basis ($\emptyset = 4.8$). Binomial logistic regression analysis revealed that exposure to chickens correlated significantly with whether an individual was colonized with *S. aureus* or not ($p = 0.019$). No association between exposure to certain animals and the occurrence of specific CC could be found. Medical history including visits to doctors or hospitals, as well as antibiotic prescription was not significantly associated

with neither *S. aureus* carriage, nor occurrence of resistance in this study.

Similarity of Genomic Profiles

All isolates grouped according to their respective CC, with the exception of one CC8 strain that clustered closely with CC5 and CC97 strains (Figure 1). The two MRSA strains grouped very far from each other. The same applies for the two *pvl* positive strains. Genomic profiles of CC1, CC6, CC7, CC59, CC101, CC121, CC188, and CC361 all clustered closely.

Resistance Genes


A broad variety of resistance genes was found among isolates. The most common resistance genes were *blaZ/I/R* ($n = 52$) conferring β -lactam resistance and *fosB* ($n = 40$) coding for a metallothiol transferase, followed by *ermA* ($n = 3$) an erythromycin resistance gene, and *tetK* ($n = 2$) coding for tetracycline resistance. Two strains were identified as MRSA (CC8, CC30), both harboring the *mecA* cassette. Both were isolated from family members of farmers (Figure 1).

Other resistance genes included the arginine catabolic mobile element (ACME) that provides multiple immune modulating functions including resistance to polyamines (Joshi et al., 2012). A fusidic acid resistance protein *fusC*, and the genes *msr(A)* and *mph(C)* both coding for macrolide efflux pumps (Ross et al., 1995; Matsuoka et al., 2003) were observed. Genes *aadD* and *aphA3* conferring resistance to aminoglycosides such as kanamycin and

TABLE 1 | Clonal complexes found in nasal samples from farmers and pupils.

Clonal complex	Farmers	Pupils	Total	Resistance genes ^a	Enterotoxin genes ^a	Comments ^a
CC1	1		1	<i>fusC</i> , <i>blaZ/I/R</i>	<i>sea^b</i> , <i>seb</i> , <i>seh</i> , <i>sek</i> , <i>seq</i>	
CC5	1	1	2	<i>blaZ/I/R</i> , <i>fosB</i>	<i>sed</i> , <i>sej</i> , <i>ser</i> , <i>egc^c</i>	
CC6		1	1	<i>fosB</i>	<i>sea^b</i>	
CC7	2	1	3	<i>blaZ/I/R</i> (2)	<i>sea^b</i>	
CC8	4	2	6	<i>mecA</i> (1), <i>blaZ/I/R</i> (3), <i>ACME</i> (1), <i>msr(A)</i> (1), <i>mph(C)</i> (1), <i>aphA3</i> (1), <i>fosB</i>	<i>sea^b</i> (2), <i>sed</i> (1), <i>sej</i> (1), <i>sek</i> (2), <i>seq</i> (2), <i>ser</i> (1), <i>egc^c</i> (1)	MRSA (<i>mecA</i>) and <i>pvl</i> (1)
CC12		1	1	<i>blaZ/I/R</i> , <i>fosB</i>	<i>sea^b</i>	
CC15	3	10	13	<i>blaZ/I/R</i> (12), <i>lnuA</i> (1), <i>aadD</i> (1), <i>tetK</i> (1), <i>fosB</i>		
CC22	1		1	<i>blaZ/I/R</i>	<i>egc</i>	
CC30	9	4	13	<i>mecA</i> (1), <i>blaZ/I/R</i> , <i>ermA</i> (2), <i>fosB</i>	<i>tst1</i> (9), <i>sea^b</i> (5), <i>sec</i> (1), <i>sel</i> (1), <i>egc</i>	MRSA (<i>mecA</i>) (1)
CC45	5	11	16	<i>blaZ/I/R</i> (8)	<i>sec</i> (10), <i>seg</i> (1), <i>sel</i> (10), <i>egc^c</i>	
CC59	1	1	2	<i>ermA</i> (1)		
CC97	2	1	3			
CC101	1		1	<i>blaZ/I/R</i> , <i>fosB</i>		
CC121	2		2	<i>blaZ/I/R</i> , <i>fosB</i>	<i>egc^c</i>	
CC152		1	1	<i>blaZ/I/R</i> , <i>tetK</i> , <i>qacA</i>		<i>pvl</i>
CC182	1		1		<i>seh</i> , <i>egc^c</i>	
CC188		1	1	<i>blaZ/I/R</i>		
CC361	1		1	<i>blaZ/I/R</i> , <i>fosB</i>	<i>egc^c</i>	
CC398		3	3	<i>blaZ/I/R</i> (2), <i>qacC</i> (1)		
CC509	1		1	<i>blaZ/I/R</i>	<i>egc^c</i>	
Total	35	38	73			

Resistance and enterotoxin genes found by microarray are listed, grouped by CC. ^aIf not all isolates assigned to the respective CC harbored a gene, the number of positive isolates is indicated in brackets. ^bEither *sea* or allelic variant *seaN315* was found, the respective variant is indicated in Supplementary Table S3. ^cEnterotoxin gene cluster (*egc*) containing genes: *seg*, *sei*, *sem*, *sen*, *seo*, *seu*.



gentamycin were found. Additionally, *InuA* a gene responsible for lincosamide modification, and *qacA* and *caqC* encoding antiseptic resistance were discovered (Supplementary Table S3).

Enterotoxin Genes and Other Virulence Factors

Of the 73 isolates 48 isolates (66%) carried at least one enterotoxin gene (Table 1). Two isolates (CC8, CC152) carried the virulence factor panton-valentine leukocidin (*pvl*) gene that is responsible for severe skin lesions and necrotizing pneumonia (Lina et al., 1999; Kaneko and Kamio, 2004). One of the obtained *pvl* isolates was identified as an MRSA strain (Figure 1). All strains harboring the *tst1* gene ($n = 9$) were assigned to CC30 (Figure 1). The most common enterotoxin genes were the genes grouped in the

enterotoxin gene cluster *egc*: *seg*, *sei*, *sem*, *sen*, *seo*, *seu* ($n = 38$), *sea* ($n = 13$), *sec* ($n = 11$), and *sel* ($n = 11$). Other enterotoxin genes included *seb* ($n = 1$), *sed* ($n = 3$), *seh* ($n = 2$), *sej* ($n = 3$), *sek* ($n = 3$), and *seq* ($n = 3$).

DISCUSSION

The observed colonization frequency of 46% (44% in pupils and 48% in family members of farmers) is in accordance with current estimates for nasal carriage rates (Sakr et al., 2018). When we compared the genomic profiles of all isolates, there were no overlaps, meaning that all isolates were unique in their genomic profile and therefore not clonal. All big clusters of CCs with more than three isolates were found in both family

members of farmers and pupils. The large clusters of CC8, CC15, CC30, and CC45 comprised 66% of isolates and all represent community or hospital associated complexes (Pérez-Montarelo et al., 2017). One CC8 strain differed from all other detected CC8 strains in view of its enterotoxin gene profile. While other CC8 strains harbored diverse enterotoxins, the outlier strain only carried *egc*. Strains carrying *egc* were shown to be more common in asymptomatic carriers than in infections (Van Belkum et al., 2006). Similar distributions have been reported (Van Belkum et al., 2006; Mohamed et al., 2019). Surprisingly, strains of CC398 were exclusively found in pupils with very limited farm animal contact (once a year or less), although this CC has regularly been linked to livestock (Verkade and Kluytmans, 2014). This may be a first indicator for a possible shift from livestock-acquired to community-acquired transmission. Most other complexes were associated with isolates from both populations. This is consistent with findings of previous studies for various CCs including CC5, CC7, CC59, and CC97 (Nulens et al., 2008; Coombs et al., 2010; Budd et al., 2015; Wang et al., 2017; Challagundla et al., 2018).

Chickens were identified as a significant factor for whether an individual was colonized with *S. aureus* or not. No significant factors that correlated with colonization of certain CC, resistance genes, or toxin profiles could be determined. This was likely due to limitations in sample size and overall low number of strains exhibiting resistance/toxin genes. It is, however, noteworthy that one of the two MRSA strains was isolated from an individual that had taken antibiotics during the last year. The same was the case for both strains harboring *pvl*, both strains harboring *qacA/C*, and one of the strains carrying *ermA*. Increased occurrence of these genes in HA-strains has been reported (Lina et al., 1999; Schmitz et al., 2000; Mayer et al., 2001). An unusual accumulation of CC152 *pvl* strains has been reported in African refugees in Switzerland, possibly representing an emerging genotype (Jaton et al., 2016). One of two MRSA strains (CC8) that was isolated from a family member on a farm with intensive contact to cattle, pigs, and chicken possessed additional genes coding for *pvl*, ACME, *blaZ/I/R*, *msr(A)*, *mph(C)*, *aphA3*, and *fosB*. This represents a combination very similar to USA300 MRSA, a CA-MRSA strain often encountered in the US and considered a “superbug” (Monecke et al., 2011; Smith, 2015; Planet, 2017). Similar strains were found in Swiss hospital patients accounting for approximately 10% of the encountered MRSA (Seidl et al., 2014). The second MRSA strain featured genetic elements that are typical for CA- or HA-acquired MRSA of CC30, including the presence of *blaZ/I/R*, *ermA*, *fosB*, *egc*, and *tst1* (Monecke et al., 2011). Another strain belonging to CC15 was classified as multiresistant according to international standard definitions for acquired resistance (Magiorakos et al., 2012) by carrying *blaZ/I/R*, *InuA*, *aadD*, *tetK*, and *fosB*. All three multiresistant strains were isolated from individuals regularly exposed to animals. While the DNA microarray data presented in this study provide new insights into the genetic structure of the tested isolates, it should not be used to extrapolate phenotypic traits.

This study provides an overview of *S. aureus* isolates found in asymptomatic carriers with varying exposure to animals. A diverse set of strains was characterized in terms of population structure and presence of virulence and resistance determinants. Our findings suggest that exposure to chicken may increase the risk of *S. aureus* nasal colonization and that intensive animal exposure may represent a risk factor for acquisition of multiresistant strains.

DATA AVAILABILITY STATEMENT

The original contributions presented in the study are publicly available. These data can be found here: <https://www.ebi.ac.uk/arrayexpress/>. Accession number: E-MTAB-8960.

ETHICS STATEMENT

Ethical review and approval was not required for the study on human participants in accordance with the local legislation and institutional requirements. The patients/participants provided their written informed consent to participate in this study.

AUTHOR CONTRIBUTIONS

SC and RS contributed to conception and design of the study. DE analyzed the data and wrote the first draft of the manuscript. SS, SC, and NC were responsible for data acquisition. DE, SJ, and RS wrote sections of the manuscript. All authors contributed to manuscript revision and read and approved the submitted version.

ACKNOWLEDGMENTS

We acknowledge all participants for filling out the questionnaire and providing nasal samples.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fmicb.2020.01289/full#supplementary-material>

TABLE S1 | Microarray profile raw data. The outcome of every hybridization reaction is shown as either positive, negative, or ambiguous as determined by the ArrayMate reader (Alere).

TABLE S2 | Questionnaire raw data. Answers to the questionnaire are provided as raw data.

TABLE S3 | Strains. List of all isolates with corresponding CCs, resistance genes, and toxin profile.

TABLE S4 | Statistics. Binomial regression analysis output from SPSS.

REFERENCES

- Ansari, S., Jha, R. K., Mishra, S. K., Tiwari, B. R., and Asaad, A. M. (2019). Recent advances in *Staphylococcus aureus* infection: focus on vaccine development. *Infect. Drug Resist.* 12, 1243–1255. doi: 10.2147/IDR.S175014
- Budd, K. E., McCoy, F., Monecke, S., Cormican, P., Mitchell, J., and Keane, O. M. (2015). Extensive genomic diversity among bovine-adapted *Staphylococcus aureus*: evidence for a genomic rearrangement within CC97. *PLoS One* 10:e0134592. doi: 10.1371/journal.pone.0134592
- Challagundla, L., Reyes, J., Rafiqullah, I., Sordelli, D. O., Echaniz-Aviles, G., Velazquez-Meza, M. E., et al. (2018). Phylogenomic classification and the evolution of clonal complex 5 methicillin-resistant *Staphylococcus aureus* in the Western Hemisphere. *Front. Microbiol.* 9:1901. doi: 10.3389/fmicb.2018.01901
- Christopher, S., Verghis, R. M., Antonisamy, B., Sowmyanarayanan, T. V., Brahmadathan, K. N., Kang, G., et al. (2011). Transmission dynamics of methicillin-resistant *Staphylococcus aureus* in a medical intensive care unit in India. *PLoS One* 6:e020604. doi: 10.1371/journal.pone.020604
- Coombs, G. W., Monecke, S., Ehrlich, R., Slickers, P., Pearson, J. C., Tan, H. L., et al. (2010). Differentiation of clonal complex 59 community-associated methicillin-resistant *Staphylococcus aureus* in Western Australia. *Antimicrob. Agents Chemother.* 54, 1914–1921. doi: 10.1128/AAC.01287-09
- Davis, M. F., Iverson, S. A., Baron, P., Vasse, A., Silbergeld, E. K., Lautenbach, E., et al. (2012). Household transmission of methicillin-resistant *Staphylococcus aureus* and other staphylococci. *Lancet Infect. Dis.* 12, 703–716. doi: 10.1016/S1473-3099(12)70156-1
- Deurenberg, R. H., and Stobberingh, E. E. (2008). The evolution of *Staphylococcus aureus*. *Infect. Genet. Evol.* 8, 747–763. doi: 10.1016/j.meegid.2008.07.007
- EFSA (2016). The European Union summary report on trends and sources of zoonoses, zoonotic agents and food-borne outbreaks in 2015. *EFSA J.* 13:3991.
- Fluit, A. C. (2012). Livestock-associated *Staphylococcus aureus*. *Clin. Microbiol. Infect.* 18, 735–744. doi: 10.1111/j.1469-0691.2012.03846.x
- Graveland, H., Wagenaar, J. A., Bergs, K., Heesterbeek, H., and Heederik, D. (2011). Persistence of livestock associated MRSA CC398 in humans is dependent on intensity of animal contact. *PLoS One* 6:e0016830. doi: 10.1371/journal.pone.0016830
- Huber, H., Giezendanner, N., Stephan, R., and Zweifel, C. (2011). Genotypes, antibiotic resistance profiles and microarray-based characterization of methicillin-resistant *Staphylococcus aureus* strains isolated from livestock and veterinarians in Switzerland. *Zoonoses Public Health* 58, 343–349. doi: 10.1111/j.1863-2378.2010.01353.x
- Huson, D. H., and Bryant, D. (2006). Application of phylogenetic networks in evolutionary studies. *Mol. Biol. Evol.* 23, 254–267. doi: 10.1093/molbev/msj030
- Jaton, L., Pillonel, T., Jaton, K., Dory, E., Prod'homme, G., Blanc, D. S., et al. (2016). Common skin infection due to panton-valentine leukocidin-producing *Staphylococcus aureus* strains in asylum seekers from Eritrea: a genome-based investigation of a suspected outbreak. *Clin. Microbiol. Infect.* 22, 739.e5–739.e5. doi: 10.1016/j.cmi.2016.05.026
- Johler, S. (2017). A glimpse into the future—new therapeutic targets could transform the way we treat staphylococcal infections. *Virulence* 8, 1508–1510. doi: 10.1080/21505594.2017.1393138
- Joshi, G., Spontak, J., Klapper, D., and Richardson, A. (2012). ACME encoded speG abrogates the unique hypersensitivity of *Staphylococcus aureus* to exogenous polyami. *Mol. Microbiol.* 82, 9–20. doi: 10.1111/j.1365-2958.2011.07809
- Kaneko, J., and Kamio, Y. (2004). Bacterial two-component and hetero-heptameric pore-forming cytolytic toxins: structures, pore-forming mechanism, and organization of the genes. *Biosci. Biotechnol. Biochem.* 68, 981–1003. doi: 10.1271/bbb.68.981
- Kluytmans, J., Van Belkum, A., and Verbrugh, H. (1997). Nasal carriage of *Staphylococcus aureus*: epidemiology, underlying mechanisms, and associated risks. *Clin. Microbiol. Rev.* 10, 505–520. doi: 10.1128/cmr.10.3.505
- Knox, J., Uhlemann, A.-C., and Lowy, F. D. (2015). *Staphylococcus aureus* infections: transmission within households and the community. *Trends Microbiol.* 23, 437–444. doi: 10.1016/j.tim.2015.03.007
- Kourtis, A. P., Hatfield, K., Baggs, J., Mu, Y., See, I., Epton, E., et al. (2019). Vital signs: epidemiology and recent trends in methicillin-resistant and in methicillin-susceptible *Staphylococcus aureus* bloodstream infections - United States. *Morb. Mortal. Wkly. Rep.* 68, 214–219. doi: 10.15585/mmwr.mm6809e1
- Kraemer, J. G., Pires, J., Kueffer, M., Semaani, E., Endimiani, A., Hilty, M., et al. (2017). Prevalence of extended-spectrum (β -lactamase-producing *Enterobacteriaceae* and methicillin-resistant *Staphylococcus aureus* in pig farms in Switzerland. *Sci. Total Environ.* 603–604, 401–405. doi: 10.1016/j.scitotenv.2017.06.110
- Kwok, K. O., Read, J. M., Tang, A., Chen, H., Riley, S., and Kam, K. M. (2018). A systematic review of transmission dynamic studies of methicillin-resistant *Staphylococcus aureus* in non-hospital residential facilities. *BMC Infect. Dis.* 18:188. doi: 10.1186/s12879-018-3060-6
- Leuenberger, A., Sartori, C., Boss, R., Resch, G., Oechslin, F., Steiner, A., et al. (2019). Genotypes of *Staphylococcus aureus*: on-farm epidemiology and the consequences for prevention of intramammary infections. *J. Dairy Sci.* 102, 3295–3309. doi: 10.3168/jds.2018-15181
- Lina, G., Piemont, Y., Godail-Gamot, F., Bes, M., Peter, M.-O., Gauduchon, V., et al. (1999). Involvement of panton-valentine leukocidin-producing *Staphylococcus aureus* in primary skin infections and pneumonia. *Clin. Infect. Dis.* 29, 1128–1132. doi: 10.1086/313461
- Maayan-Metzger, A., Strauss, T., Rubin, C., Jaber, H., Dulitzky, M., Reiss-Mandel, A., et al. (2017). Clinical evaluation of early acquisition of *Staphylococcus aureus* carriage by newborns. *Int. J. Infect. Dis.* 64, 9–14. doi: 10.1016/j.ijid.2017.08.013
- Magiorakos, A. P., Srinivasan, A., Carey, R. B., Carmeli, Y., Falagas, M. E., Giske, C. G., et al. (2012). Multidrug-resistant, extensively drug-resistant and pandrug-resistant bacteria: an international expert proposal for interim standard definitions for acquired resistance. *Clin. Microbiol. Infect.* 18, 268–281. doi: 10.1111/j.1469-0691.2011.03570.x
- Matsuoka, M., Inoue, M., Endo, Y., and Nakajima, Y. (2003). Characteristic expression of three genes, *msr(A)*, *mph(C)* and *erm(Y)*, that confer resistance to macrolide antibiotics on *Staphylococcus aureus*. *FEMS Microbiol. Lett.* 220, 287–293. doi: 10.1016/S0378-1097(03)00134-4
- Maura, D., Ballok, A. E., and Rahme, L. G. (2016). Considerations and caveats in anti-virulence drug development. *Curr. Opin. Microbiol.* 33, 41–46. doi: 10.1016/j.mib.2016.06.001
- Mayer, S., Boos, M., Beyer, A., Fluid, A. C., and Schmitz, F.-J. (2001). Distribution of the antiseptic resistance genes. *J. Antimicrob. Chemother.* 47, 896–897.
- Mégevand, C., Gervaix, A., Heininger, U., Berger, C., Aebi, C., Vaudaux, B., et al. (2010). Molecular epidemiology of the nasal colonization by methicillin-susceptible *Staphylococcus aureus* in Swiss children. *Clin. Microbiol. Infect.* 16, 1414–1420. doi: 10.1111/j.1469-0691.2009.03090.x
- Mohamed, N., Timofeyeva, Y., Jamroz, D., Rojas, E., Hao, L., Silmon de Monerri, N. C., et al. (2019). Molecular epidemiology and expression of capsular polysaccharides in *Staphylococcus aureus* clinical isolates in the United States. *PLoS One* 14:e0208356. doi: 10.1371/journal.pone.0208356
- Monecke, S., Coombs, G., Shore, A. C., Coleman, D. C., Akpaka, P., Borg, M., et al. (2011). A field guide to pandemic, epidemic and sporadic clones of methicillin-resistant *Staphylococcus aureus*. *PLoS One* 6:e17936. doi: 10.1371/journal.pone.0017936
- Morach, M., Käppeli, N., Hochreutener, M., Johler, S., Julmi, J., Stephan, R., et al. (2019). Microarray based genetic profiling of *Staphylococcus aureus* isolated from abattoir byproducts of pork origin. *PLoS One* 14:e0222036. doi: 10.1371/journal.pone.0222036
- Nouwen, J., Schouten, J., Schneebergen, P., Snijders, S., Maaskant, J., Koolen, M., et al. (2006). *Staphylococcus aureus* carriage patterns and the risk of infections associated with continuous peritoneal dialysis. *J. Clin. Microbiol.* 44, 2233–2236. doi: 10.1128/JCM.02083-05
- Nulens, E., Stobberingh, E. E., Van Dessel, H., Sebastian, S., Van Tiel, F. H., Beisser, P. S., et al. (2008). Molecular characterization of *Staphylococcus aureus* bloodstream isolates collected in a Dutch university hospital between 1999 and 2006. *J. Clin. Microbiol.* 46, 2438–2441. doi: 10.1128/JCM.00808-08
- Olearo, F., Albrich, W. C., Vernaz, N., Harbarth, S., and Kronenberg, A. (2016). *Staphylococcus aureus* and methicillin resistance in Switzerland: regional differences and trends from 2004 to 2014. *Swiss Med. Wkly.* 146:w14339. doi: 10.4414/sm.w.2016.14339
- Pérez-Montarelo, D., Viedma, E., Murcia, M., Muñoz-Gallego, I., Larrosa, N., Brañas, P., et al. (2017). Pathogenic characteristics of *Staphylococcus aureus* endovascular infection isolates from different clonal complexes. *Front. Microbiol.* 8:917. doi: 10.3389/fmicb.2017.00917
- Planet, P. J. (2017). Life after USA300: the rise and fall of a superbug. *J. Infect. Dis.* 215, S71–S77. doi: 10.1093/infdis/jiw444

- Rosenkranz Wettstein, K., Rothenanger, E., Brodard, I., Collaud, A., Overesch, G., Bigler, B., et al. (2014). Nasal carriage of methicillin-resistant *Staphylococcus aureus* (MRSA) among Swiss veterinary health care providers: detection of livestock- and healthcare-associated clones. *Schweiz. Arch. Tierheilkd.* 156, 317–325. doi: 10.1024/0036-7281
- Ross, J. I., Eady, E. A., Cove, J. H., and Baumberg, S. (1995). Identification of a chromosomally encoded ABC-transport system with which the staphylococcal erythromycin exporter MsrA may interact. *Gene* 153, 93–98. doi: 10.1016/0378-1119(94)00833-E
- Sakr, A., Brégeon, F., Mège, J. L., Rolain, J. M., and Blin, O. (2018). *Staphylococcus aureus* nasal colonization: an update on mechanisms, epidemiology, risk factors, and subsequent infections. *Front. Microbiol.* 9:2419. doi: 10.3389/fmicb.2018.02419
- Schmitz, F. J., Sadurski, R., Kray, A., Boos, M., Geisel, R., Koehler, K., et al. (2000). Prevalence of macrolide-resistance genes in *Staphylococcus aureus* and *Enterococcus faecium* isolates from 24 European university hospitals. *J. Antimicrob. Chemother.* 45, 891–894. doi: 10.1093/jac/45.6.891
- Seidl, K., Leimer, N., Palheiros Marques, M., Furrer, A., Senn, G., Holzmann-Bürgel, A., et al. (2014). USA300 methicillin-resistant *Staphylococcus aureus* in Zurich, Switzerland between 2001 and 2013. *Int. J. Med. Microbiol.* 304, 1118–1122. doi: 10.1016/j.ijmm.2014.08.005
- Smith, T. C. (2015). Livestock-associated *Staphylococcus aureus*: the United States experience. *PLoS Pathog.* 11:e1004564. doi: 10.1371/journal.ppat.1004564
- Tong, S. Y. C., Davis, J. S., Eichenberger, E., Holland, T. L., and Fowler, V. G. (2015). *Staphylococcus aureus* infections: epidemiology, pathophysiology, clinical manifestations, and management. *Clin. Microbiol. Rev.* 28, 603–661. doi: 10.1128/CMR.00134-14
- Van Belkum, A., Melles, D. C., Snijders, S. V., Van Leeuwen, W. B., Wertheim, H. F. L., Nouwen, J. L., et al. (2006). Clonal distribution and differential occurrence of the enterotoxin gene cluster, *egc*, in carriage- versus bacteremia-associated isolates of *Staphylococcus aureus*. *J. Clin. Microbiol.* 44, 1555–1557. doi: 10.1128/JCM.44.4.1555-1557.2006
- Verkade, E., and Kluytmans, J. (2014). Livestock-associated *Staphylococcus aureus* CC398: animal reservoirs and human infections. *Infect. Genet. Evol.* 21, 523–530. doi: 10.1016/j.meegid.2013.02.013
- Vestergaard, M., Frees, D., and Ingmer, H. (2019). Antibiotic resistance and the MRSA problem. *Microbiol. Spectr.* 7, 1–23. doi: 10.1128/microbiolspec.gpp3-0057-2018
- Wang, X. L., Li, L., Li, S. M., Huang, J. Y., Fan, Y. P., Yao, Z. J., et al. (2017). Phenotypic and molecular characteristics of *Staphylococcus aureus* and methicillin-resistant *Staphylococcus aureus* in slaughterhouse pig-related workers and control workers in Guangdong Province, China. *Epidemiol. Infect.* 145, 1843–1851. doi: 10.1017/S095026881700085
- Wattlinger, L., Stephan, R., Layer, F., and Jöhler, S. (2012). Comparison of *Staphylococcus aureus* isolates associated with food intoxication with isolates from human nasal carriers and human infections. *Eur. J. Clin. Microbiol. Infect. Dis.* 31, 455–464. doi: 10.1007/s10096-011-1330-y
- Wertheim, H. F. L., Melles, D. C., Vos, M. C., Van Leeuwen, W., Van Belkum, A., Verbrugh, H. A., et al. (2005). The role of nasal carriage in *Staphylococcus aureus* infections. *Lancet Infect. Dis.* 5, 751–762. doi: 10.1016/S1473-3099(05)70295-4

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2020 Etter, Corti, Spirig, Cernela, Stephan and Jöhler. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.