

Received 11 October 2024 Accepted 30 November 2024

Edited by T. Ohhara, J-PARC Center, Japan Atomic Energy Agency, Japan

**Keywords:** crystal structure; superacidic system; ester; protonation; Raman spectroscopy.

**CCDC references:** 2407012; 2407013; 2407014; 2407015; 2407016

**Supporting information:** this article has supporting information at journals.iucr.org/c





# The crystal structures of methyl prop-2-ynoate, dimethyl fumarate and their protonated species

#### Dirk Hollenwäger,\* Valentin Bockmair and Andreas J. Kornath

Department Chemie, Ludwig-Maximilians Universität, Butenandtstrasse 5-13 (Haus D), D-81377 München, Germany. \*Correspondence e-mail: dirk.hollenwaeger@cup.uni-muenchen.de

Methyl prop-2-ynoate  $(C_4H_4O_2)$  was investigated in the binary superacidic system HF/MF<sub>5</sub> (M = Sb or As) and dimethyl fumarate  $(C_6H_8O_4)$  in the superacidic system HF/SbF<sub>5</sub>, as well as HF/BF<sub>3</sub>. The starting materials methyl prop-2ynoate and dimethyl fumarate were crystallized, the former for the first time. The protonated species of these esters, namely, (1-methoxyprop-2-yn-1-ylidene) oxidanium hexafluoroarsenate,  $C_4H_5O_2^+$  AsF<sub>6</sub><sup>-</sup>, 1,4-dimethoxy-4-oxobut-2-en-1ylidene]oxidanium tetrafluoroborate bis(hydrogen fluoride),  $C_6H_9O_4^+$  BF<sub>4</sub><sup>-</sup> 2HF, and hemi{[1,4-dimethoxy-4-oxidaniumylidenebut-2-en-1-ylidene]oxidanium} undecafluorodiantimonate,  $0.5C_6H_{10}O_4^{2+}$  Sb<sub>2</sub>F<sub>11</sub><sup>-</sup>, were characterized by single-crystal X-ray diffraction and Raman spectroscopy. The protonated species were recrystallized from anhydrous hydrogen fluoride. In the solid state of the monoprotonated species of methyl prop-2-ynoate and the diprotonated species of dimethyl fumarate, strong intramolecular  $O-H \cdots F$  hydrogen bonds build a three-dimensional network. The monoprotonated species of dimethyl fumarate builds chains by strong  $O-H \cdots O$  hydrogen bonds between the cations.

#### 1. Introduction

Protonated esters have occur in two conformations, namely, *syn-anti* and *syn-syn* (Hogeveen, 1967; Olah *et al.*, 1967, 2009). The *syn-anti* conformation is more stable and is therefore consistent with the protonation of carboxylic acids (Olah *et al.*, 2009; Hollenwäger *et al.*, 2024*b*; Hogeveen, 1968). The two conformers were observed in solution by NMR spectroscopy (Olah *et al.*, 2009). It has not yet been possible to crystallize the *syn-syn* conformer of protonated esters in the solid state; the example of prop-2-ynoic acid (propiolic acid) shows that this could be achieved with a H/D exchange and solid-state effects through a larger anion (Hollenwäger *et al.*, 2024*b*). In magic acid (FSO<sub>3</sub>H/SbF<sub>5</sub>), the esters show the unimolecular cleavage of methanol by warming to 20 °C (Olah *et al.*, 2009). An exception was observed with glycine methyl ester, which is still stable in magic acid even at 93 °C (Hollenwäger, *et al.*, 2024*a*).

The isolation of protonated esters enables the characterization of an important intermediate that is present in solution in every acid-catalyzed esterification process. The selected esters also offer the possibility of further functionalization steps due to the double and triple bonds present. This prompted us to investigate methyl prop-2-ynoate and dimethyl fumarate in the binary superacidic media  $HF/MF_5$ .

#### 2. Experimental

2.1. Synthesis and crystallization

2.1.1.  $[C_4H_5O_4][MF_6]$  (*M* = Sb or As)

The Lewis acids (SbF<sub>5</sub>: 433 mg, 2 mmol; AsF<sub>5</sub>: 340 mg, 2 mmol) were each condensed into a fluorine-passivated FEP

reactor. Anhydrous hydrogen fluoride (0.5 l) was added as reactant and solvent at -196 °C. The mixture was homogenized at room temperature. Methyl prop-2-ynoate (83.6 µl, 84.1 mg, 1.0 mmol) was added at -196 °C under nitrogen. The mixture was allowed to warm to room temperature. The solvent was removed overnight at -78 °C. The protonated species II and III were obtained as white solids (Scheme 1).

#### 2.1.2. $[C_6H_9O_4][MF_y]$ (*M* = Sb or B; y = 6 or 4)

The Lewis acids (SbF<sub>5</sub>: 216 mg, 1 mmol; BF<sub>3</sub>: 67 mg, 1 mmol) were each condensed into a fluorine-passivated FEP reactor. Anhydrous hydrogen fluoride (0.5 l) was added as reactant and solvent at -196 °C. The mixture was homogenized at room temperature. Dimethyl fumarate (144 mg, 0.5 mmol) was added at -196 °C under nitrogen. The mixture was allowed to warm to room temperature. The solvent was removed overnight at -78 °C. The protonated species V and VI were obtained as white solids (Scheme 1). A clean Raman spectrum of monoprotonated species VI could not be obtained because it contains impurities of either IV or VII.



#### 2.1.3. $[C_6H_{10}O_4][Sb_2F_{11}]_2$ and $[C_6H_{10}O_4][BF_4]_2$

The Lewis acids (SbF<sub>5</sub>: 216 mg, 1 mmol; BF<sub>3</sub>: 67 mg, 1 mmol) were each condensed into a fluorine-passivated FEP reactor. Anhydrous hydrogen fluoride (0.5 l) was added as reactant and solvent at -196 °C. The mixture was homogenized at room temperature. Dimethyl fumarate (48 mg, 0.33 mmol) was added at -196 °C under nitrogen. The mixture was allowed to warm to room temperature. The solvent was removed overnight at -78 °C. The protonated species **VII** and **VIII** were obtained as white solids.

## 2.2. Single-crystal X-ray diffraction and Raman spectroscopic analysis

Compounds I, III, VI and VII were characterized by singlecrystal X-ray diffraction. Complete data and devices for the X-ray measurements are listed in the CIF in the supporting information. Low-temperature Raman spectroscopic analysis



#### Figure 1

A difference Fourier map of **III** without the H atom between O1 and F1. The green solid lines and red dotted lines show positive and negative density distribution, respectively.

was performed for **I–VIII** using a Bruker MultiRAM FT– Raman spectrometer with Nd:YAG laser excitation ( $\lambda = 1064 \text{ cm}^{-1}$ ) under vacuum. For the measurements, the synthesized compound was transferred to a cooled glass cell.

#### 2.3. Refinement

Crystal data, data collection and structure refinement details are summarized in Table 1. The successful protonation of the target molecules was confirmed by the charge of the asymmetric unit, as well as the interatomic distances. C-O distances have become nearly equal after protonation, as the charge can formally be localized on the C atom resulting in the loss of double-bond character. The positions of the H atoms were identified by Q-peaks on the difference Fourier map and



#### Figure 2

A difference Fourier map of **VI** without the H atom between O1 and F1. The green solid lines and red dotted lines show positive and negative density distribution, respectively.

#### Table 1

Experimental details.

Experiments were carried out with Mo  $K\alpha$  radiation using a Rigaku Xcalibur Sapphire3 diffractometer. Absorption was corrected for by multi-scan methods (*CrysAlis PRO*; Rigaku OD, 2020).

	I	III	IV
Crystal data			
Chemical formula	$C_4H_4O_2$	$C_4H_5O_2^+ \cdot AsF_6^-$	$C_6H_8O_4$
$M_{ m r}$	84.07	274.00	144.12
Crystal system, space group	Monoclinic, $P2_1/n$	Monoclinic, $P2_1/n$	Triclinic, $P\overline{1}$
temperature (K)	111	111	112
<i>a</i> , <i>b</i> , <i>c</i> (Å)	3.8409 (5), 15.593 (2), 7.6149 (10)	6.9609 (5), 8.9319 (7), 13.7189 (9)	3.8726 (11), 5.6546 (10), 8.3778 (18)
$lpha,eta,\gamma~(^\circ)$	90, 99.910 (12), 90	90, 91.664 (7), 90	100.642 (16), 100.42 (2), 105.73 (2)
$V(Å^3)$	449.27 (10)	852.60 (11)	168.30 (7)
Z	4	4	1
$\mu  ({\rm mm}^{-1})$	0.10	4.06	0.12
Crystal size (mm)	$1.00 \times 0.56 \times 0.31$	$0.90\times0.21\times0.15$	$0.52 \times 0.46 \times 0.35$
Data collection			
$T_{\min}, T_{\max}$	0.457, 1.000	0.582, 1.000	0.579, 1.000
No. of measured, independent and observed $[I > 2\sigma(I)]$ reflections	3885, 1060, 891	4205, 2303, 1900	1403, 825, 703
R <sub>int</sub>	0.023	0.029	0.015
$\theta_{max}(\circ)$	27.9	29.1	28.3
$(\sin \theta / \lambda)_{\rm max} ({\rm \AA}^{-1})$	0.658	0.685	0.667
Refinement			
$R[F^2 > 2\sigma(F^2)], wR(F^2), S$	0.036, 0.099, 1.05	0.039, 0.099, 1.05	0.048, 0.144, 1.04
No. of reflections	1060	2303	825
No. of parameters	71	123	51
No. of restraints	0	0	0
H-atom treatment	All H-atom parameters refined	H atoms treated by a mixture of independent and constrained refinement	H atoms treated by a mixture of independent and constrained refinement
$\Delta \rho_{\rm max},  \Delta \rho_{\rm min} \ ({\rm e} \ {\rm \AA}^{-3})$	0.18, -0.15	0.87, -0.92	0.52, -0.24

#### VI

Crystal data		
Chemical formula	$C_6H_9O_4^+ \cdot BF_4^- \cdot 2HF$	$0.5C_6H_{10}O_4^{2+}Sb_2F_{11}^{-}$
M <sub>r</sub>	271.96	525.57
Crystal system, space group	Orthorhombic, Pbca	Orthorhombic, Pbca
temperature (K)	101	101
a, b, c (Å)	12.8759 (5), 11.8899 (4), 14.6252 (7)	7.8461 (6), 15.1531 (11), 19.5536 (17)
$lpha,eta,\gamma(^\circ)$	90, 90, 90	90, 90, 90
$V(Å^3)$	2239.02 (16)	2324.8 (3)
Ζ	8	8
$\mu \text{ (mm}^{-1})$	0.19	4.79
Crystal size (mm)	$0.28\times0.18\times0.11$	$0.25 \times 0.11 \times 0.05$
Data collection		
$T_{\min}, T_{\max}$	0.885, 1.000	0.807, 1.000
No. of measured, independent and observed $[I > 2\sigma(I)]$ reflections	7514, 1717, 1478	13316, 3784, 2878
R <sub>int</sub>	0.026	0.053
$\theta_{\max}$ (°)	23.8	32.2
$(\sin \theta / \lambda)_{\max} ( \text{\AA}^{-1} )$	0.568	0.750
Refinement		
$R[F^2 > 2\sigma(F^2)], wR(F^2), S$	0.040, 0.100, 1.05	0.036, 0.071, 1.04
No. of reflections	1717	3784
No. of parameters	176	169
No. of restraints	1	0
H-atom treatment	H atoms treated by a mixture of independent and constrained refinement	H atoms treated by a mixture of independent and constrained refinement
$\Delta \rho_{\text{max}} \Delta \rho_{\text{min}} (e  \text{\AA}^{-3})$	0.77, -0.35	1.091.01

computer programs: CrysAlis PRO (Rigaku OD, 2020), SHELXT (Sheldrick, 2015a), SHELXL2018 (Sheldrick, 2015b), ORTEP-3 for Windows (Farrugia, 2012) and PLATON (Spek, 2020).

VII

#### research papers



#### Figure 3

A difference Fourier map of **VII** without the H atom between O1 and F1. The green solid lines and red dotted lines show positive and negative density distribution, respectively.

by evaluation of the contacts (Figs. 1–3). Methyl, methylene and acetylenic H atoms were refined under restrictions and the proton positions were modulated.

#### 3. Results and discussion

#### 3.1. Single-crystal X-ray diffraction

#### 3.1.1. Crystal structure of methyl prop-2-ynoate (I)

Compound I crystallizes in the monoclinic space group  $P2_1/n$  with one formula unit per unit cell. Fig. 4 displays the asymmetric unit. The C1–C2 bond length [1.4466 (15) Å] is significantly elongated compared to an average  $Csp^1-Csp^2$  hybridized bond (1.427 Å) determined by X-ray diffraction (Allen *et al.*, 1987). The C2=C3 triple bond [1.1780 (16) Å] is in the same range as average terminal C=C bonds (1.181 Å; Allen *et al.*, 1987). The C1=O1 bond [1.1972 (14) Å] is in the same range as other C=O bonds (1.196 Å) in esters (Allen *et al.*, 1987). The C1–O2 bond [1.3195 (12) Å] is significantly shortened compared to other esters (1.337 Å; Allen *et al.*, 1987). The C4–O2 bond [1.4463 (13) Å] is significantly elongated compared to average CH<sub>3</sub>–O bonds in esters (1.418 Å; Allen *et al.*, 1987).



Figure 4

The asymmetric unit of **I**, with displacement ellipsoids drawn at the 50% probability level.



Hydrogen bonds in the crystal structure of **I**, with displacement ellipsoids drawn at the 50% probability level. [Symmetry codes: (i)  $x - \frac{1}{2}, -y + \frac{3}{2}, z + \frac{1}{2};$  (ii) -x + 1, -y + 1, -z + 2; (iii)  $x + \frac{1}{2}, -y + \frac{3}{2}, z - \frac{1}{2}.$ ]

The crystal structure of **I** displays a layered structure built of weak C3-H1 $\cdots$ O1 hydrogen bonds, according to the classification of Jeffrey (1997). The layered structure is connected *via* weak C4-H2A $\cdots$ O1 hydrogen bonds into a three-dimensional network (Fig. 5), according to the classification of Jeffrey (1997).

### 3.1.2. Crystal structure of (1-methoxyprop-2-yn-1-ylidene) oxidanium hexafluoroarsenate (III)

Salt III crystallizes in the monoclinic space group  $P_{2_1}/n$  with four formula units per unit cell. Fig. 6 displays the asymmetric unit of III. The C1–O1 bond [1.261 (4) Å] is significantly elongated by 0.064 Å due to the protonation compared to I. The C1–O2 bond [1.270 (3) Å] is significantly shortened by 0.049 Å compared to the starting material I. Due to the protonation, the C4–O2 bond [1.484 (4) °] is elongated by 0.038 Å compared to the neutral compound I. The C2==C3 triple bond is not significantly influenced by the protonation.

The three-dimensional network of **III** (Fig. 7) is built by a strong  $O1-H3\cdots F1$  and three weak  $C3-H1\cdots F5$ ,  $C3-H1\cdots F6$  and  $C4-H2B\cdots F3$  hydrogen bonds, according to the classification of Jeffrey (1997). Additionally, the crystal structure forms two interatomic contacts ( $C1\cdots F2$  and  $C1\cdots F5$ ) which are 8% shorter than the sum of the van der Waals radii.



Figure 6

The asymmetric unit of **III**, with displacement ellipsoids drawn at the 50% probability level.



Figure 7

The intramolecular interactions in the crystal structure of **III**, with displacement ellipsoids drawn at the 50% probability level. [Symmetry codes: (i)  $-x + \frac{1}{2}$ ,  $y - \frac{1}{2}$ ,  $-z + \frac{1}{2}$ ; (ii) -x, -y + 1, -z + 1; (iii) -x + 1, -y + 1, -z + 1; (iv) x, y - 1, z; (v)  $-x + \frac{3}{2}$ ,  $y - \frac{1}{2}$ ,  $-z + \frac{1}{2}$ .]

#### 3.1.3. Crystal structure of dimethyl (E)-but-2-enedioate (IV)

The determined crystal structure is in the same range as that reported by Kooijman *et al.* (2004). The formula unit is shown in Fig. 8 and the crystal structure exhibits the same three-dimensional network (Fig. 9).

#### 3.1.4. Crystal structure of 1,4-dimethoxy-4-oxobut-2-en-1ylidene]oxidanium tetrafluoroborate-hydrogen fluoride (1/2) (VI)

The crystal structure of the monoprotonated species **VI** of dimethyl fumarate crystallizes in the orthorhombic space group *Pbca* with eight formula units per unit cell. Fig. 10 displays the asymmetric unit. Similar to fumaric acid and acetylenedicarboxylic acid, the monoprotonated type forms extended chains of cations that are connected *via* strong O3–H9···O1 hydrogen bonds (Jessen & Kornath, 2022; Bayer *et al.*, 2020; Jeffrey, 1997). Due to the protonation, the C1–O1 [1.248 (3) Å] and C4–O3 [1.247 (3) Å] bonds are significantly elongated compared to the neutral compound [1.205 (2) Å]. The C1–O2 and C4–O4 bonds [both 1.289 (3) Å] are shortened by 0.052 Å compared to the starting material **IV**. The CH<sub>3</sub>–O and C2=C3 bonds are not significantly influenced by the monoprotonation.

Besides the strong hydrogen bonding between the cations, the crystal structure forms a medium-strong  $C6-H6\cdots F6$ 



Figure 8

The formula unit of **IV**, with displacement ellipsoids drawn at the 50% probability level. [Symmetry code: (i) -x + 1, -y + 2, -z + 1.]



Figure 9

The intramolecular interactions in the crystal structure of **IV**, with displacement ellipsoids drawn at the 50% probability level. [Symmetry codes: (i) -x + 1, -y + 2, -z + 1; (ii) x - 1, y - 1, z; (iii) -x, -y + 1, -z + 1; (iv) x - 1, y, z; (v) -x, -y + 2, -z + 1; (vi) x + 1, y, z; (vii) x + 1, y + 1, z; (viii) -x + 2, -y + 2, -z + 1; (ix) -x + 2, -y + 3, -z + 1.]

hydrogen bond [3.198 (3) Å] and two weak C5–H3···F4 [3.242 (3) Å] and C6–H7···F2 [3.261 (3) Å] hydrogen bonds. Furthermore, the crystal structure forms six interatomic C···F contacts, *i.e.* C1···F1 [2.900 (3) Å], C1···F6 [2.943 (3) Å], C2···F6 [2.987 (3) Å], C3···F6 [3.048 (3) Å], C4···F2 [3.020 (3) Å] and C4···F5 [2.974 (3) Å], which are shorter than the sum of the van der Waals radii (3.17 Å). The interactions in the crystal structure of **VI** are shown in Fig. 11.

#### 3.1.5. Crystal structure of hemi{[1,4-dimethoxy-4-oxidaniumylidenebut-2-en-1-ylidene]oxidanium} undecafluorodiarsenate (VII)

The crystal structure of the diprotonated species **VII** of dimethyl fumarate crystallizes in the orthorhombic space group Pbca with eight formula units per unit cell. Fig. 12 displays the formula unit. The crystal structure of the dipro-



**Figure 10** The asymmetric unit of **VI**, with displacement ellipsoids drawn at the 50% probability level.



Figure 11

The intramolecular interactions in the crystal structure of **VI**, with displacement ellipsoids drawn at the 50% probability level. [Symmetry codes: (i)  $-x + \frac{1}{2}$ ,  $y - \frac{1}{2}$ , z; (ii)  $-x + \frac{1}{2}$ ,  $y + \frac{1}{2}$ , z; (iii)  $x - \frac{1}{2}$ , y + 1,  $-z + \frac{1}{2}$ ; (iv) x, y + 1, z; (v) -x + 1, -y + 1, -z + 1; (vi) x,  $-y + \frac{1}{2}$ ,  $z + \frac{1}{2}$ ; (vii)  $x - \frac{1}{2}$ ,  $-y + \frac{1}{2}$ , -z + 1.]

tonated species has a C1–O1 bond [1.277 (5) Å] significantly elongated by 0.072 Å compared to the starting material [1.205 (2) Å]. The C1–O2 bond [1.281 (5) Å] is shortened by 0.060 Å compared to the neutral compound [1.341 (1) Å]. The C3–O2 bond [1.489 (6) Å] is elongated by 0.034 Å compared to **IV** [1.455 (2) Å].

The three-dimensional network is formed by a strong O1-H1...F1 hydrogen bond and seven interatomic interactions (Jeffrey, 1997). The interatomic interactions are C1...F4 [3.061 (4) Å], C1...F7 [3.001 (4) Å], C2...F3 [2.958 (5) Å], C2...F7 [3.008 (5) Å], C3...F2 [3.050 (6) Å], O1...F7 [2.985 (4) Å] and O1...F11 [2.864 (4) Å]. Fig. 13 displays the interatomic distances in the crystal structure of **VII**.



Figure 13

The intramolecular interactions in the crystal structure of **VIII**, with displacement ellipsoids drawn at the 50% probability level. [Symmetry codes: (i) -x + 1, -y + 1, -z + 1; (ii)  $-x + \frac{3}{2}$ ,  $y + \frac{1}{2}$ , z; (iii)  $x + \frac{1}{2}$ ,  $-y + \frac{1}{2}$ , -z + 1; (iv) -x + 2, -y + 1, -z + 1; (v)  $x - \frac{1}{2}$ ,  $-y + \frac{1}{2}$ , -z + 1; (vi)  $-x + \frac{1}{2}$ ,  $-y + \frac{1}{2}$ , z; (vii)  $x - \frac{1}{2}$ ,  $-y + \frac{1}{2}$ , -z + 1; (vii) x - 1, y, z.]

#### 3.2. Raman spectroscopy

#### 3.2.1. Raman spectra of I, II and III

Fig. 14 displays the low-temperature Raman spectra of **I**, **II** and **III**. The first evidence for succesful protonation is the significantly red-shifted C=O oscillation from  $1700 \text{ cm}^{-1}$  in the starting material **I** to 1617 (in **II**) and 1615 cm<sup>-1</sup> (in **III**). Due to the protonation, the C–O oscillation is blue-shifted in the Raman spectra to 1414 (in **II**) and 1413 cm<sup>-1</sup> (in **III**) compared to the starting material ( $1278 \text{ cm}^{-1}$ ). The H<sub>3</sub>C–O oscillation is red-shifted by  $42 \text{ cm}^{-1}$  to 948 (in **II**) and 951 cm<sup>-1</sup> (in **III**) compared to **I** (990 cm<sup>-1</sup>). The oscillation of the triple bond is only slightly affected from 2107 cm<sup>-1</sup> to 2141 (in **II**) and 2140 cm<sup>-1</sup> (in **III**).



Figure 12

The formula unit of **VIII**, with displacement ellipsoids drawn at the 50% probability level. [Symmetry code: (i) -x + 1, -y + 1, -z + 1.]

C=O C=C C=O  $H_{3}C=O$   $H_{3}C=O$   $H_{3}C=O$   $C_{4}H_{6}O_{2}][SbF_{6}]$   $C_{4}H_{6}O_{2}][SbF_{6}]$   $C_{4}H_{6}O_{2}][SbF_{6}]$   $C_{4}H_{6}O_{2}][SbF_{6}]$   $C_{4}H_{6}O_{2}][SbF_{6}]$   $C_{4}H_{6}O_{2}][SbF_{6}]$   $C_{4}H_{6}O_{2}][SbF_{6}]$   $C_{4}H_{6}O_{2}][SbF_{6}]$   $C_{4}H_{6}O_{2}][SbF_{6}]$   $C_{6}H_{6}O_{2}][SbF_{6}]$   $C_{6}H_{6}O_{2}]$   $C_{6}H_{6}O_{2}]$   $C_{6}H_{6}O_{2}]$   $C_{6}H_{6}O_{2}]$   $C_{6}H_{6}O_{2}]$   $C_{7}H_{6}O_{2}]$   $C_{7}H_{7}O_{2}$   $C_{7}$ 

Figure 14

The low-temperature Raman spectra of I (black),  $I\!I$  (red) and  $I\!I\!I$  (blue).

#### 3.2.2. Raman spectra of IV, V, VII and VIII

Fig. 15 shows the low-temperature Raman spectra of **IV**, **V**, **VII** and **VIII**. The C==O oscillation is red-shifted by 48 cm<sup>-1</sup> to 1611 cm<sup>-1</sup> compared to **IV** (1659 cm<sup>-1</sup>). Due to the protonation, the C–O oscillation is blue-shifted in the Raman spectra to 1401 cm<sup>-1</sup> (**V**) compared to the neutral compound (1217 cm<sup>-1</sup>). The C==C oscillation is only slightly red shifted to 1705 cm<sup>-1</sup> compared to the starting material (1725 cm<sup>-1</sup>).

The diprotonation is characterized by a red-shifted C=O oscillation by 46 cm<sup>-1</sup> to 1613 cm<sup>-1</sup> in the spectra of **VII** and **VIII** compared to the Raman spectrum of **IV** (1659 cm<sup>-1</sup>). The C=C oscillation is significantly red-shifted by 38 cm<sup>-1</sup> to 1686 cm<sup>-1</sup> in the spectra of **VII** and **VIII** compared to the Raman spectrum of **IV** (1724 cm<sup>-1</sup>). Due to the protonation, the C–O oscillation is blue-shifted in the Raman spectra to 1419 (in **VII**) and 1425 cm<sup>-1</sup> (in **VIII**) compared to the neutral compound (1217 cm<sup>-1</sup>).

#### 4. Conclusion

We present herein the first single-crystal X-ray diffraction and Raman spectroscopy study of the monoprotonated species of methyl prop-2-ynoate and mono- and diprotonated species of dimethyl fumarate. All three protonated species crystallize in the more stable *syn-anti* conformation. Furthermore, the first single-crystal structure of methyl prop-2-ynoate is reported. The protonated species are important intermediates of acidcatalyzed reactions.

#### Acknowledgements

We are grateful to the Department of Chemistry at the Ludwig Maximilian University of Munich, the Deutsche Forschungsgemeinschaft (DFG), the F-Select GmbH and Professor Dr Konstantin Karaghiosoff and Dr Constantin Hoch for their support. Open access funding enabled and organized by Projekt DEAL.

#### **Funding information**

Funding for this research was provided by: Ludwig-Maximilians-University; F-Select GmbH; Deutsche Forschungsgemeinschaft.



Figure 15

The low-temperature Raman spectra of IV (black), V (red), VII (green) and VIII (blue).

#### References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). J. Chem. Soc. Perkin Trans. 2, pp. S1–S19.
- Bayer, M. C., Jessen, C. & Kornath, A. J. (2020). Z. Anorg. Allg. *Chem.* **646**, 333–339.
- Farrugia, L. J. (2012). J. Appl. Cryst. 45, 849-854.
- Hogeveen, H. (1967). Recl Trav. Chim. Pays Bas, 86, 816-820.
- Hogeveen, H. (1968). Recl Trav. Chim. Pays Bas, 87, 1313-1317.
- Hollenwäger, D., Morgenstern, Y., Daumer, L., Bockmair, V. & Kornath, A. J. (2024a). ACS Earth Space Chem. 8, 2101–2109.
- Hollenwäger, D., Thamm, S., Bockmair, V., Nitzer, A. & Kornath, A. J. (2024b). J. Org. Chem. 89, 11421–11428.
- Jeffrey, G. A. (1997). In *An Introduction to hydrogen Bonding*. New York, Oxford: Oxford University Press.
- Jessen, C. & Kornath, A. J. (2022). Eur. J. Inorg. Chem. 2022, e202100965,
- Kooijman, H., Sprengers, J. W., Agerbeek, M. J., Elsevier, C. J. & Spek, A. L. (2004). Acta Cryst. E60, 0917–0918.
- Olah, G. A., O'Brien, D. H. & White, A. M. (1967). J. Am. Chem. Soc. 89, 5694–5700.
- Olah, G. A., Prakash, G. K. S., Molnar, A. & Sommer, J. (2009). In *Superacid Chemistry*, 2nd ed. Hoboken, NJ: Wiley.
- Rigaku OD (2020). CrysAlis PRO. Rigaku Oxford Diffraction Ltd, Yarnton, Oxfordshire, England.
- Sheldrick, G. M. (2015a). Acta Cryst. A71, 3-8.
- Sheldrick, G. M. (2015b). Acta Cryst. C71, 3-8.
- Spek, A. L. (2020). Acta Cryst. E76, 1-11.

Acta Cryst. (2025). C81, 31-37 [https://doi.org/10.1107/S2053229624011653]

The crystal structures of methyl prop-2-ynoate, dimethyl fumarate and their protonated species

### Dirk Hollenwäger, Valentin Bockmair and Andreas J. Kornath

**Computing details** 

Methyl prop-2-ynoate (I)

Crystal data

C<sub>4</sub>H<sub>4</sub>O<sub>2</sub>  $M_r = 84.07$ Monoclinic,  $P2_1/n$  a = 3.8409 (5) Å b = 15.593 (2) Å c = 7.6149 (10) Å  $\beta = 99.910$  (12)° V = 449.27 (10) Å<sup>3</sup> Z = 4

#### Data collection

Rigaku Xcalibur Sapphire3 diffractometer Radiation source: Enhance (Mo) X-ray Source Graphite monochromator Detector resolution: 15.9809 pixels mm<sup>-1</sup>  $\omega$  scans Absorption correction: multi-scan (CrysAlis PRO; Rigaku OD, 2020)  $T_{\min} = 0.457, T_{\max} = 1.000$ 

#### Refinement

Refinement on  $F^2$ Least-squares matrix: full  $R[F^2 > 2\sigma(F^2)] = 0.036$  $wR(F^2) = 0.099$ S = 1.051060 reflections 71 parameters 0 restraints Primary atom site location: structure-invariant direct methods F(000) = 176  $D_x = 1.243 \text{ Mg m}^{-3}$ Mo K\alpha radiation,  $\lambda = 0.71073 \text{ Å}$ Cell parameters from 1269 reflections  $\theta = 3.0-31.5^{\circ}$   $\mu = 0.10 \text{ mm}^{-1}$  T = 111 KNeedle, colorless  $1.00 \times 0.56 \times 0.31 \text{ mm}$ 

3885 measured reflections 1060 independent reflections 891 reflections with  $I > 2\sigma(I)$  $R_{int} = 0.023$  $\theta_{max} = 27.9^{\circ}, \ \theta_{min} = 3.0^{\circ}$  $h = -5 \rightarrow 5$  $k = -12 \rightarrow 20$  $l = -10 \rightarrow 10$ 

Secondary atom site location: difference Fourier map Hydrogen site location: difference Fourier map All H-atom parameters refined  $w = 1/[\sigma^2(F_o^2) + (0.0475P)^2 + 0.0483P]$ where  $P = (F_o^2 + 2F_c^2)/3$  $(\Delta/\sigma)_{max} = 0.001$  $\Delta\rho_{max} = 0.18 \text{ e } \text{Å}^{-3}$  $\Delta\rho_{min} = -0.15 \text{ e } \text{Å}^{-3}$ 

#### Special details

**Geometry**. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement**. Refinement of  $F^2$  against ALL reflections. The weighted *R*-factor *wR* and goodness of fit *S* are based on  $F^2$ , conventional *R*-factors *R* are based on *F*, with *F* set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating *R*-factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. *R*-factors based on  $F^2$  are statistically about twice as large as those based on *F*, and *R*- factors based on ALL data will be even larger.

)
)
)
)
)
)
*
*
*
*
) * * * *

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters  $(\hat{A}^2)$ 

Atomic displacement parameters  $(Å^2)$ 

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
02	0.0382 (4)	0.0368 (4)	0.0295 (4)	-0.0001 (3)	0.0097 (3)	-0.0030 (3)
01	0.0635 (6)	0.0427 (5)	0.0340 (5)	0.0123 (4)	0.0229 (4)	0.0042 (3)
C1	0.0265 (5)	0.0361 (6)	0.0249 (5)	0.0037 (4)	0.0031 (4)	0.0007 (4)
C2	0.0328 (5)	0.0416 (6)	0.0308 (6)	0.0017 (4)	0.0064 (4)	-0.0010 (4)
C4	0.0390 (6)	0.0356 (6)	0.0398 (7)	0.0007 (4)	0.0056 (5)	0.0014 (5)
C3	0.0465 (7)	0.0409 (6)	0.0462 (7)	-0.0022 (5)	0.0124 (5)	0.0041 (5)

*Geometric parameters (Å, °)* 

02—C1	1.3195 (12)	C1—C2	1.4466 (15)
O2—C4	1.4463 (13)	C2—C3	1.1780 (16)
01—C1	1.1972 (14)		
C1—O2—C4	115.83 (9)	O2—C1—C2	111.02 (9)
01—C1—O2	124.88 (10)	C3—C2—C1	179.47 (11)
O1—C1—C2	124.10 (10)		
C4—O2—C1—O1	0.50 (16)	C4—O2—C1—C2	-179.25 (9)

(1-Methoxyprop-2-yn-1-ylidene)oxidanium hexafluoroarsenate (III)

#### Crystal data

 $C_{4}H_{5}O_{2}^{+} \cdot AsF_{6}^{-}$   $M_{r} = 274.00$ Monoclinic,  $P2_{1}/n$  a = 6.9609 (5) Å b = 8.9319 (7) Å c = 13.7189 (9) Å  $\beta = 91.664$  (7)° V = 852.60 (11) Å<sup>3</sup> Z = 4

#### Data collection

Rigaku Xcalibur Sapphire3	4205 measured reflections
diffractometer	2303 independent reflections
Radiation source: Enhance (Mo) X-ray Source	1900 reflections with $I > 2\sigma(I)$
Graphite monochromator	$R_{\rm int} = 0.029$
Detector resolution: 15.9809 pixels mm <sup>-1</sup>	$\theta_{\rm max} = 29.1^\circ, \ \theta_{\rm min} = 2.7^\circ$
$\omega$ scans	$h = -6 \rightarrow 9$
Absorption correction: multi-scan	$k = -9 \rightarrow 12$
(CrysAlis PRO; Rigaku OD, 2020)	$l = -18 \rightarrow 18$
$T_{\min} = 0.582, \ T_{\max} = 1.000$	

#### Refinement

Secondary atom site location: difference Fourier
map
Hydrogen site location: mixed
H atoms treated by a mixture of independent
and constrained refinement
$w = 1/[\sigma^2(F_o^2) + (0.0444P)^2]$
where $P = (F_0^2 + 2F_c^2)/3$
$(\Delta/\sigma)_{\rm max} = 0.001$
$\Delta \rho_{\rm max} = 0.87 \ {\rm e} \ {\rm \AA}^{-3}$
$\Delta \rho_{\min} = -0.92 \text{ e} \text{ Å}^{-3}$

#### Special details

**Geometry**. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

F(000) = 528

 $\theta = 2.3 - 31.5^{\circ}$ 

 $\mu = 4.06 \text{ mm}^{-1}$ 

Needle, colorless

 $0.90 \times 0.21 \times 0.15 \text{ mm}$ 

T = 111 K

 $D_{\rm x} = 2.135 {\rm Mg} {\rm m}^{-3}$ 

Mo *K* $\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 1298 reflections

**Refinement**. Refinement of  $F^2$  against ALL reflections. The weighted *R*-factor *wR* and goodness of fit *S* are based on  $F^2$ , conventional *R*-factors *R* are based on *F*, with *F* set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating *R*-factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. *R*-factors based on  $F^2$  are statistically about twice as large as those based on *F*, and *R*-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters  $(Å^2)$ 

	x	У	Ζ	$U_{ m iso}$ */ $U_{ m eq}$	
As1	0.55531 (4)	0.61946 (4)	0.31141 (2)	0.02340 (12)	
F3	0.3663 (2)	0.7351 (3)	0.28285 (13)	0.0380 (5)	
F5	0.5944 (3)	0.5983 (2)	0.18954 (13)	0.0380 (5)	
F2	0.5053 (3)	0.6364 (3)	0.43228 (14)	0.0462 (6)	

O2	0.0897 (3)	0.1533 (3)	0.50117 (16)	0.0301 (5)
01	0.1633 (3)	0.3569 (3)	0.42071 (18)	0.0310 (5)
F4	0.7411 (3)	0.5020(3)	0.33987 (16)	0.0503 (6)
F1	0.3982 (3)	0.4674 (3)	0.30251 (16)	0.0508 (6)
F6	0.7072 (3)	0.7683 (3)	0.31930 (17)	0.0496 (6)
C1	0.1973 (4)	0.2223 (4)	0.44299 (19)	0.0244 (6)
C2	0.3535 (4)	0.1383 (4)	0.4053 (2)	0.0256 (6)
C3	0.4795 (4)	0.0709 (4)	0.3697 (2)	0.0323 (7)
H1	0.580815	0.016719	0.341095	0.039*
C4	-0.0767 (4)	0.2351 (5)	0.5405 (3)	0.0417 (9)
H2A	-0.030985	0.324746	0.575262	0.063*
H2B	-0.144696	0.170089	0.585530	0.063*
H2C	-0.164273	0.264647	0.486657	0.063*
Н3	0.235 (6)	0.394 (5)	0.387 (3)	0.047 (13)*

Atomic displacement parameters  $(\mathring{A}^2)$ 

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
As1	0.02680 (18)	0.0244 (2)	0.01897 (17)	0.00042 (12)	-0.00046 (12)	0.00004 (12)
F3	0.0363 (9)	0.0460 (13)	0.0313 (9)	0.0157 (10)	-0.0033 (8)	-0.0030 (10)
F5	0.0382 (10)	0.0531 (14)	0.0230 (9)	0.0054 (9)	0.0071 (8)	-0.0007 (9)
F2	0.0596 (13)	0.0605 (16)	0.0187 (8)	0.0094 (11)	0.0016 (9)	-0.0008 (10)
O2	0.0335 (11)	0.0311 (13)	0.0260 (10)	-0.0035 (10)	0.0083 (9)	-0.0020 (10)
01	0.0305 (11)	0.0281 (13)	0.0344 (12)	0.0014 (10)	0.0023 (10)	-0.0015 (11)
F4	0.0579 (12)	0.0450 (14)	0.0469 (12)	0.0253 (11)	-0.0158 (10)	-0.0018 (11)
F1	0.0657 (14)	0.0441 (14)	0.0430 (12)	-0.0300 (12)	0.0066 (10)	-0.0002 (11)
F6	0.0425 (11)	0.0429 (14)	0.0631 (14)	-0.0170 (10)	-0.0034 (10)	-0.0051 (13)
C1	0.0251 (13)	0.0279 (17)	0.0200 (13)	-0.0020 (12)	-0.0033 (11)	-0.0057 (13)
C2	0.0283 (14)	0.0260 (17)	0.0226 (13)	0.0005 (12)	0.0000 (11)	0.0028 (13)
C3	0.0292 (14)	0.0346 (19)	0.0333 (16)	0.0049 (14)	0.0032 (12)	-0.0012 (15)
C4	0.0339 (16)	0.049 (2)	0.0436 (19)	-0.0021 (17)	0.0149 (15)	-0.0097 (19)

Geometric parameters (Å, °)

As1—F6	1.700 (2)	O2—C1	1.270 (3)
As1—F4	1.702 (2)	O2—C4	1.484 (4)
As1—F3	1.7091 (19)	O1—C1	1.261 (4)
As1—F2	1.7109 (19)	C1—C2	1.430 (4)
As1—F5	1.7122 (18)	C2—C3	1.180 (4)
As1—F1	1.746 (2)		
F6—As1—F4	90.03 (12)	F6—As1—F1	179.51 (12)
F6—As1—F3	90.88 (11)	F4—As1—F1	90.37 (13)
F4—As1—F3	179.09 (11)	F3—As1—F1	88.72 (12)
F6—As1—F2	90.72 (11)	F2—As1—F1	89.55 (11)
F4—As1—F2	90.39 (11)	F5—As1—F1	87.85 (10)
F3—As1—F2	89.53 (10)	C1—O2—C4	118.0 (3)
F6—As1—F5	91.87 (11)	O1—C1—O2	120.3 (3)

F4—As1—F5	90.84 (10)	01—C1—C2	123.4 (3)
F3—As1—F5	89.20 (9)	02—C1—C2	116.2 (3)
F2—As1—F5	177.13 (10)	C3—C2—C1	176.8 (3)
C4—O2—C1—O1	0.8 (4)	C4—O2—C1—C2	-178.7 (3)

Dimethyl (E)-but-2-enedioate (IV)

#### Crystal data

$C_6H_8O_4$	Z = 1
$M_r = 144.12$	F(000) = 76
Triclinic, $P\overline{1}$	$D_{\rm x} = 1.422 {\rm Mg} {\rm m}^{-3}$
a = 3.8726 (11)  Å	Mo <i>K</i> $\alpha$ radiation, $\lambda = 0.71073$ Å
b = 5.6546 (10)  Å	Cell parameters from 712 reflections
c = 8.3778 (18)  Å	$\theta = 3.9 - 32.0^{\circ}$
$\alpha = 100.642 \ (16)^{\circ}$	$\mu = 0.12 \text{ mm}^{-1}$
$\beta = 100.42 \ (2)^{\circ}$	T = 112  K
$\gamma = 105.73 \ (2)^{\circ}$	Plate, colorless
$V = 168.30 (7) \text{ Å}^3$	$0.52 \times 0.46 \times 0.35 \text{ mm}$
Data collection	
Rigaku Xcalibur Sapphire3	1403 measured reflections
diffractometer	825 independent reflections
Radiation source: Enhance (Mo) X-ray Source	703 reflections with $I > 2\sigma(I)$
Graphite monochromator	$R_{\rm int} = 0.015$
Detector resolution: 15.9809 pixels mm <sup>-1</sup>	$\theta_{\rm max} = 28.3^{\circ}, \ \theta_{\rm min} = 4.1^{\circ}$
ω scans	$h = -5 \rightarrow 5$
Absorption correction: multi-scan	$k = -5 \rightarrow 7$
(CrysAlis PRO; Rigaku OD, 2020)	$l = -10 \longrightarrow 11$
$T_{\min} = 0.579, \ T_{\max} = 1.000$	
Refinement	

Refinement on $F^2$	Secondary atom site location: difference Fourier
Least-squares matrix: full	map
$R[F^2 > 2\sigma(F^2)] = 0.048$	Hydrogen site location: mixed
$wR(F^2) = 0.144$	H atoms treated by a mixture of independent
S = 1.04	and constrained refinement
825 reflections	$w = 1/[\sigma^2(F_o^2) + (0.1042P)^2 + 0.0039P]$
51 parameters	where $P = (F_o^2 + 2F_c^2)/3$
0 restraints	$(\Delta/\sigma)_{\rm max} < 0.001$
Primary atom site location: structure-invariant	$\Delta  ho_{ m max} = 0.52 \ { m e} \ { m \AA}^{-3}$
direct methods	$\Delta \rho_{\rm min} = -0.24 \text{ e } \text{\AA}^{-3}$

#### Special details

**Geometry**. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement**. Refinement of  $F^2$  against ALL reflections. The weighted *R*-factor *wR* and goodness of fit *S* are based on  $F^2$ , conventional *R*-factors *R* are based on *F*, with *F* set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating *R*-factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. *R*-factors based on  $F^2$  are statistically about twice as large as those based on *F*, and *R*- factors based on ALL data will be even larger.

	x	v	Ζ	$U_{iso}^*/U_{eq}$	
$\overline{02}$	0 2058 (2)	0 46773 (15)	0 30549 (11)	0 0270 (3)	
01	0.5682 (3)	0.77635 (18)	0.21852 (12)	0.0381 (4)	
C2	0.3998 (3)	0.8781 (2)	0.47844 (15)	0.0244 (4)	
C1	0.4062 (3)	0.7084 (2)	0.32036 (14)	0.0237 (4)	
C3	0.1927 (4)	0.2863 (2)	0.15494 (16)	0.0295 (4)	
H1A	0.078071	0.332324	0.056057	0.044*	
H1B	0.046722	0.116090	0.156867	0.044*	
H1C	0.444625	0.288310	0.150486	0.044*	
H1	0.246 (5)	0.796 (3)	0.544 (2)	0.034 (4)*	

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters  $(\hat{A}^2)$ 

Atomic displacement parameters  $(Å^2)$ 

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
O2	0.0328 (5)	0.0188 (5)	0.0261 (5)	0.0011 (4)	0.0112 (4)	0.0039 (4)
01	0.0484 (6)	0.0260 (6)	0.0335 (6)	-0.0030 (4)	0.0210 (5)	0.0035 (4)
C2	0.0249 (6)	0.0231 (7)	0.0238 (6)	0.0045 (5)	0.0075 (4)	0.0056 (5)
C1	0.0235 (6)	0.0206 (6)	0.0242 (6)	0.0027 (5)	0.0059 (4)	0.0048 (5)
С3	0.0354 (7)	0.0197 (6)	0.0278 (7)	0.0028 (5)	0.0086 (5)	0.0006 (5)

Geometric parameters (Å, °)

$\begin{array}{cccccccccccccccccccccccccccccccccccc$					
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	02—C1	1.3414 (14)	C2C2 <sup>i</sup>	1.329 (2)	
$O1-C1$ $1.2055 (15)$ $C1-O2-C3$ $115.47 (9)$ $O1-C1-C2$ $125.05 (11)$ $C2^{i}-C2-C1$ $120.37 (14)$ $O2-C1-C2$ $111.03 (11)$ $O1-C1-O2$ $123.92 (11)$ $C2^{i}-C2-C1-O1$ $-4.2 (2)$ $C3-O2-C1-C2$ $179.80 (9)$ $C2^{i}-C2-C1-O2$ $175.91 (13)$	O2—C3	1.4549 (14)	C2—C1	1.4937 (17)	
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	01—C1	1.2055 (15)			
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	C1—O2—C3	115.47 (9)	01—C1—C2	125.05 (11)	
$O1-C1-O2$ $123.92 (11)$ $C3-O2-C1-O1$ $-0.07 (19)$ $C2^{i}-C2-C1-O1$ $-4.2 (2)$ $C3-O2-C1-C2$ $179.80 (9)$ $C2^{i}-C2-C1-O2$ $175.91 (13)$	$C2^{i}$ — $C2$ — $C1$	120.37 (14)	O2—C1—C2	111.03 (11)	
C3-O2-C1-O1 $-0.07 (19)$ C2 <sup>i</sup> -C2-C1-O1 $-4.2 (2)$ C3-O2-C1-C2179.80 (9)C2 <sup>i</sup> -C2-C1-O2175.91 (13)	O1—C1—O2	123.92 (11)			
C3-O2-C1-C2 179.80 (9) $C2^{i}$ -C2-C1-O2 175.91 (13)	C3—O2—C1—O1	-0.07 (19)	C2 <sup>i</sup> —C2—C1—O1	-4.2 (2)	
	C3—O2—C1—C2	179.80 (9)	C2 <sup>i</sup> —C2—C1—O2	175.91 (13)	

Symmetry code: (i) -x+1, -y+2, -z+1.

1,4-Dimethoxy-4-oxobut-2-en-1-ylidene]oxidanium tetrafluoroborate-hydrogen fluoride (1/2) (VI)

Crystal data  $C_6H_9O_4^+ \cdot BF_4^- \cdot 2HF$   $M_r = 271.96$ Orthorhombic, *Pbca*  a = 12.8759 (5) Å b = 11.8899 (4) Å c = 14.6252 (7) Å V = 2239.02 (16) Å<sup>3</sup> Z = 8F(000) = 1104

 $D_x = 1.614 \text{ Mg m}^{-3}$ Mo K $\alpha$  radiation,  $\lambda = 0.71073 \text{ Å}$ Cell parameters from 2510 reflections  $\theta = 2.7-31.9^{\circ}$  $\mu = 0.19 \text{ mm}^{-1}$ T = 101 KPlate, colorless  $0.28 \times 0.18 \times 0.11 \text{ mm}$  Data collection

Rigaku Xcalibur Sapphire3 diffractometer Radiation source: Enhance (Mo) X-ray Source Graphite monochromator Detector resolution: 15.9809 pixels mm <sup>-1</sup> $\omega$ scans Absorption correction: multi-scan (CrysAlis PRO; Rigaku OD, 2020) $T_{\min} = 0.885, T_{\max} = 1.000$	7514 measured reflections 1717 independent reflections 1478 reflections with $I > 2\sigma(I)$ $R_{int} = 0.026$ $\theta_{max} = 23.8^{\circ}, \theta_{min} = 2.7^{\circ}$ $h = -14 \rightarrow 11$ $k = -13 \rightarrow 6$ $l = -16 \rightarrow 16$
Refinement	
Refinement on $F^2$ Least-squares matrix: full $R[F^2 > 2\sigma(F^2)] = 0.040$ $wR(F^2) = 0.100$ S = 1.05 1717 reflections 176 parameters 1 restraint Primary atom site location: structure-invariant direct methods	Secondary atom site location: difference Fourier map Hydrogen site location: mixed H atoms treated by a mixture of independent and constrained refinement $w = 1/[\sigma^2(F_o^2) + (0.0431P)^2 + 2.1459P]$ where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{max} < 0.001$ $\Delta\rho_{max} = 0.77$ e Å <sup>-3</sup> $\Delta\rho_{min} = -0.35$ e Å <sup>-3</sup>

#### Special details

**Geometry**. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement**. Refinement of  $F^2$  against ALL reflections. The weighted *R*-factor *wR* and goodness of fit *S* are based on  $F^2$ , conventional *R*-factors *R* are based on *F*, with *F* set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating *R*-factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. *R*-factors based on  $F^2$  are statistically about twice as large as those based on *F*, and *R*- factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters  $(\hat{A}^2)$ 

	x	У	Ζ	$U_{ m iso}$ */ $U_{ m eq}$
F1	0.52277 (12)	0.16806 (12)	0.38516 (10)	0.0424 (4)
F5	0.50136 (12)	0.27683 (13)	0.53107 (11)	0.0415 (4)
O4	0.32442 (12)	0.62198 (12)	0.58784 (10)	0.0255 (4)
01	0.13464 (12)	0.86421 (12)	0.36669 (10)	0.0249 (4)
O3	0.36950 (12)	0.47179 (12)	0.50843 (11)	0.0256 (4)
O2	0.18127 (12)	0.71387 (13)	0.28735 (10)	0.0275 (4)
F6	0.36144 (13)	-0.08191 (13)	0.38730 (12)	0.0467 (4)
F2	0.68532 (12)	0.09644 (14)	0.38787 (12)	0.0525 (5)
F4	0.54899 (14)	-0.01889 (13)	0.39813 (14)	0.0598 (5)
F3	0.58307 (16)	0.06993 (16)	0.26486 (12)	0.0681 (6)
C4	0.32631 (17)	0.56549 (18)	0.51262 (15)	0.0223 (5)
C2	0.23256 (17)	0.71960 (18)	0.44011 (16)	0.0229 (5)
C1	0.17955 (17)	0.77141 (18)	0.36205 (15)	0.0222 (5)
C3	0.27536 (18)	0.61916 (19)	0.43439 (15)	0.0232 (5)
C6	0.3732 (2)	0.5736 (2)	0.66902 (16)	0.0340 (6)
H7	0.332494	0.508833	0.689926	0.051*

H6	0.375883	0.630283	0.717592	0.051*	
H8	0.443830	0.549154	0.653995	0.051*	
C5	0.1229 (2)	0.7558 (2)	0.20841 (16)	0.0328 (6)	
H4	0.048343	0.747717	0.219943	0.049*	
Н5	0.141976	0.712458	0.153940	0.049*	
Н3	0.139489	0.835347	0.198520	0.049*	
B1	0.5867 (2)	0.0793 (2)	0.35695 (19)	0.0294 (6)	
H2	0.2735 (18)	0.5775 (19)	0.3809 (16)	0.025 (6)*	
H1	0.2311 (19)	0.758 (2)	0.4951 (17)	0.030 (6)*	
H10	0.509 (3)	0.231 (3)	0.485 (3)	0.088 (14)*	
H11	0.425 (3)	-0.056 (3)	0.396 (3)	0.088 (13)*	
H9	0.363 (3)	0.421 (2)	0.4363 (11)	0.101 (13)*	

Atomic displacement parameters  $(Å^2)$ 

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	<i>U</i> <sup>23</sup>
F1	0.0422 (9)	0.0355 (8)	0.0493 (9)	0.0150 (7)	-0.0058 (7)	-0.0100 (7)
F5	0.0399 (9)	0.0439 (9)	0.0408 (9)	0.0092 (7)	0.0031 (7)	-0.0027 (7)
O4	0.0301 (9)	0.0235 (8)	0.0231 (9)	0.0014 (7)	-0.0017 (7)	-0.0002 (7)
01	0.0253 (9)	0.0184 (8)	0.0311 (9)	0.0009 (7)	-0.0025 (7)	0.0018 (7)
03	0.0245 (9)	0.0187 (8)	0.0336 (9)	0.0026 (7)	0.0009 (7)	0.0017 (7)
02	0.0305 (9)	0.0290 (9)	0.0231 (8)	0.0060 (7)	-0.0015 (7)	0.0002 (7)
F6	0.0404 (10)	0.0405 (9)	0.0593 (11)	-0.0046 (8)	-0.0029 (8)	0.0040 (8)
F2	0.0313 (9)	0.0594 (11)	0.0667 (11)	0.0051 (8)	-0.0063 (8)	-0.0041 (9)
F4	0.0510 (11)	0.0370 (9)	0.0915 (14)	-0.0035 (8)	0.0098 (10)	0.0054 (9)
F3	0.0852 (14)	0.0799 (13)	0.0393 (10)	0.0153 (11)	-0.0028 (9)	-0.0159 (9)
C4	0.0189 (11)	0.0191 (12)	0.0289 (13)	-0.0028 (10)	0.0013 (9)	0.0006 (10)
C2	0.0241 (12)	0.0216 (12)	0.0231 (12)	-0.0011 (10)	0.0019 (10)	-0.0014 (10)
C1	0.0182 (11)	0.0207 (12)	0.0277 (12)	-0.0020 (10)	0.0012 (9)	0.0012 (10)
C3	0.0253 (12)	0.0210 (11)	0.0234 (12)	-0.0016 (10)	-0.0002 (10)	-0.0027 (10)
C6	0.0383 (15)	0.0376 (14)	0.0261 (13)	-0.0023 (12)	-0.0086 (11)	0.0054 (11)
C5	0.0348 (14)	0.0400 (14)	0.0238 (12)	0.0042 (12)	-0.0042 (11)	0.0049 (11)
B1	0.0270 (15)	0.0262 (14)	0.0349 (16)	0.0014 (12)	0.0013 (12)	-0.0065 (12)

Geometric parameters (Å, °)

F1—B1	1.401 (3)	F2—B1	1.363 (3)
O4—C4	1.289 (3)	F4—B1	1.401 (3)
O4—C6	1.461 (3)	F3—B1	1.352 (3)
01—C1	1.247 (3)	C4—C3	1.465 (3)
O3—C4	1.247 (3)	C2—C3	1.318 (3)
O2—C1	1.289 (3)	C2—C1	1.466 (3)
O2—C5	1.465 (3)		
C4—O4—C6	118.69 (18)	O2—C1—C2	115.41 (19)
C1—O2—C5	118.59 (17)	C2—C3—C4	122.2 (2)
O3—C4—O4	121.1 (2)	F3—B1—F2	112.1 (2)
O3—C4—C3	123.4 (2)	F3—B1—F1	109.6 (2)

O4—C4—C3	115.54 (19)	F2—B1—F1	109.7 (2)
C3—C2—C1	121.7 (2)	F3—B1—F4	110.3 (2)
O1—C1—O2	121.6 (2)	F2—B1—F4	107.8 (2)
O1—C1—C2	123.0 (2)	F1—B1—F4	107.3 (2)
C6—O4—C4—O3	-0.6 (3)	C3—C2—C1—O2	-2.4 (3)
C6—O4—C4—C3	179.66 (19)	C1—C2—C3—C4	-178.5 (2)
C5—O2—C1—O1	-4.1 (3)	O3—C4—C3—C2	-178.2 (2)
C5—O2—C1—C2	175.19 (19)	O4—C4—C3—C2	1.5 (3)
C3—C2—C1—O1	176.9 (2)		

Hemi{[1,4-dimethoxy-4-oxidaniumylidenebut-2-en-1-ylidene]oxidanium} undecafluorodiarsenate (VII)

 $D_{\rm x} = 3.003 {\rm Mg} {\rm m}^{-3}$ 

 $\theta = 3.1 - 31.4^{\circ}$  $\mu = 4.79 \text{ mm}^{-1}$ 

Needle, colorless  $0.25 \times 0.11 \times 0.05 \text{ mm}$ 

T = 101 K

 $R_{\rm int} = 0.053$ 

 $h = -11 \rightarrow 11$ 

 $k = -22 \rightarrow 18$ 

 $l = -19 \rightarrow 29$ 

Mo *K* $\alpha$  radiation,  $\lambda = 0.71073$  Å

13316 measured reflections 3784 independent reflections

 $\theta_{\text{max}} = 32.2^{\circ}, \ \theta_{\text{min}} = 2.9^{\circ}$ 

2878 reflections with  $I > 2\sigma(I)$ 

Cell parameters from 3099 reflections

Crystal data

 $0.5C_{6}H_{10}O_{4}^{2+}Sb_{2}F_{11}^{-}M_{r} = 525.57$ Orthorhombic, *Pbca* a = 7.8461 (6) Å b = 15.1531 (11) Å c = 19.5536 (17) Å V = 2324.8 (3) Å<sup>3</sup> Z = 8F(000) = 1920

#### Data collection

Rigaku Xcalibur Sapphire3 diffractometer Radiation source: Enhance (Mo) X-ray Source Graphite monochromator Detector resolution: 15.9809 pixels mm<sup>-1</sup>  $\omega$  scans Absorption correction: multi-scan (CrysAlis PRO; Rigaku OD, 2020)  $T_{\min} = 0.807, T_{\max} = 1.000$ 

#### Refinement

Refinement on  $F^2$ Secondary atom site location: difference Fourier Least-squares matrix: full map  $R[F^2 > 2\sigma(F^2)] = 0.036$ Hydrogen site location: mixed  $wR(F^2) = 0.071$ H atoms treated by a mixture of independent S = 1.04and constrained refinement 3784 reflections  $w = 1/[\sigma^2(F_0^2) + (0.021P)^2]$ 169 parameters where  $P = (F_0^2 + 2F_c^2)/3$ 0 restraints  $(\Delta/\sigma)_{\rm max} = 0.002$  $\Delta \rho_{\rm max} = 1.09 \text{ e } \text{\AA}^{-3}$ Primary atom site location: structure-invariant direct methods  $\Delta \rho_{\rm min} = -1.01 \ {\rm e} \ {\rm \AA}^{-3}$ 

#### Special details

**Geometry**. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement**. Refinement of  $F^2$  against ALL reflections. The weighted *R*-factor *wR* and goodness of fit *S* are based on  $F^2$ , conventional *R*-factors *R* are based on *F*, with *F* set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating *R*-factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. *R*-factors based on  $F^2$  are statistically about twice as large as those based on *F*, and *R*- factors based on ALL data will be even larger.

	<i>x</i>	У	Z	$U_{\rm iso}$ */ $U_{\rm eq}$
Sb1	0.53289 (3)	0.11951 (2)	0.58884 (2)	0.01838 (7)
Sb2	0.87980 (3)	0.29435 (2)	0.64518 (2)	0.01906 (8)
F6	0.6878 (3)	0.20173 (15)	0.64187 (14)	0.0290 (6)
F4	0.3807 (3)	0.04618 (14)	0.54353 (14)	0.0254 (6)
F11	1.0501 (3)	0.37949 (16)	0.64744 (16)	0.0325 (7)
F7	0.7631 (3)	0.34255 (15)	0.57101 (14)	0.0267 (6)
F1	0.4121 (3)	0.22282 (15)	0.56555 (15)	0.0288 (6)
F8	0.7357 (3)	0.35628 (15)	0.70374 (15)	0.0313 (6)
F2	0.4096 (4)	0.11295 (17)	0.66985 (15)	0.0331 (6)
F3	0.6723 (3)	0.13957 (15)	0.51323 (14)	0.0278 (6)
O2	0.4521 (4)	0.52155 (17)	0.62437 (16)	0.0208 (6)
F5	0.6789 (3)	0.03114 (16)	0.61891 (17)	0.0360 (7)
F10	0.9958 (3)	0.21939 (16)	0.58552 (17)	0.0348 (7)
F9	0.9623 (4)	0.23004 (17)	0.71862 (17)	0.0414 (8)
O1	0.4065 (4)	0.37941 (18)	0.61581 (18)	0.0226 (6)
H1	0.400397	0.339397	0.586174	0.034*
C1	0.4470 (5)	0.4521 (2)	0.5871 (2)	0.0175 (8)
C3	0.4204 (6)	0.5136 (3)	0.6992 (3)	0.0283 (10)
H4	0.312992	0.482094	0.706860	0.043*
Н5	0.413086	0.572665	0.719463	0.043*
Н3	0.514077	0.480919	0.720535	0.043*
C2	0.4888 (5)	0.4604 (3)	0.5143 (2)	0.0189 (9)
H2	0.498 (5)	0.410 (3)	0.488 (2)	0.006 (10)*

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters*  $(Å^2)$ 

Atomic displacement parameters (A	nt parameters (Å <sup>2</sup> )
-----------------------------------	---------------------------------

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Sb1	0.02181 (13)	0.01431 (12)	0.01902 (15)	-0.00328 (10)	-0.00136 (11)	0.00006 (12)
Sb2	0.02106 (13)	0.01545 (12)	0.02066 (16)	-0.00268 (10)	-0.00279 (11)	0.00125 (12)
F6	0.0361 (14)	0.0251 (12)	0.0258 (16)	-0.0164 (10)	-0.0012 (12)	-0.0014 (12)
F4	0.0274 (13)	0.0227 (11)	0.0260 (15)	-0.0048 (9)	-0.0033 (11)	-0.0063 (12)
F11	0.0221 (13)	0.0275 (13)	0.048 (2)	-0.0080 (10)	-0.0049 (13)	0.0029 (13)
F7	0.0308 (14)	0.0278 (12)	0.0213 (15)	0.0027 (10)	-0.0024 (11)	0.0062 (12)
F1	0.0360 (15)	0.0194 (12)	0.0311 (17)	0.0062 (10)	-0.0033 (12)	-0.0022 (12)
F8	0.0339 (15)	0.0317 (12)	0.0283 (16)	-0.0049 (11)	0.0051 (12)	-0.0102 (13)
F2	0.0389 (16)	0.0416 (15)	0.0189 (15)	-0.0156 (12)	0.0049 (12)	0.0013 (13)
F3	0.0304 (14)	0.0282 (12)	0.0247 (15)	-0.0035 (10)	0.0094 (11)	-0.0026 (12)
O2	0.0278 (15)	0.0178 (13)	0.0169 (16)	-0.0014 (11)	0.0022 (13)	-0.0029 (13)
F5	0.0330 (14)	0.0241 (12)	0.051 (2)	-0.0011 (10)	-0.0142 (15)	0.0103 (14)
F10	0.0333 (15)	0.0254 (13)	0.046 (2)	0.0047 (10)	0.0046 (13)	-0.0071 (13)

F9	0.0490 (18)	0.0352 (15)	0.040 (2)	-0.0077 (13)	-0.0171 (15)	0.0145 (15)
01	0.0253 (15)	0.0161 (13)	0.0263 (18)	0.0005 (11)	0.0040 (13)	-0.0002 (13)
C1	0.0161 (18)	0.0158 (17)	0.021 (2)	0.0045 (14)	-0.0015 (16)	-0.0022 (18)
C3	0.041 (3)	0.027 (2)	0.017 (2)	0.0022 (18)	0.001 (2)	-0.004 (2)
C2	0.0175 (19)	0.0186 (18)	0.021 (2)	0.0007 (14)	0.0006 (16)	-0.0020 (18)

Geometric parameters (Å, °)

1.856 (2)	Sb2—F7	1.864 (3)
1.858 (2)	Sb2—F10	1.865 (3)
1.859 (3)	Sb2—F6	2.060 (2)
1.864 (3)	O2—C1	1.281 (5)
1.886 (2)	O2—C3	1.489 (6)
2.026 (2)	O1—C1	1.277 (5)
1.852 (3)	C1—C2	1.466 (6)
1.858 (2)	$C2$ — $C2^{i}$	1.334 (8)
1.863 (3)		
96.68 (11)	F9—Sb2—F7	168.79 (11)
92.29 (12)	F11—Sb2—F7	95.72 (12)
90.73 (14)	F8—Sb2—F7	89.00 (12)
95.54 (11)	F9—Sb2—F10	89.65 (14)
90.38 (13)	F11—Sb2—F10	94.98 (12)
171.91 (11)	F8—Sb2—F10	170.88 (11)
93.34 (11)	F7—Sb2—F10	89.53 (13)
169.97 (12)	F9—Sb2—F6	85.50 (11)
89.34 (12)	F11—Sb2—F6	178.88 (11)
88.18 (12)	F8—Sb2—F6	85.34 (11)
176.70 (11)	F7—Sb2—F6	83.29 (11)
84.91 (11)	F10—Sb2—F6	85.55 (11)
84.78 (11)	Sb1—F6—Sb2	150.90 (15)
87.33 (11)	C1—O2—C3	119.2 (3)
85.11 (11)	O1—C1—O2	117.7 (4)
95.48 (12)	O1—C1—C2	123.8 (4)
90.04 (14)	O2—C1—C2	118.4 (4)
94.12 (12)	$C2^{i}$ — $C2$ — $C1$	120.8 (5)
3.3 (5)	01—C1—C2—C2 <sup>i</sup>	172.3 (5)
-176.4 (4)	$O2-C1-C2-C2^{i}$	-8.1 (7)
	$\begin{array}{c} 1.856\ (2)\\ 1.858\ (2)\\ 1.859\ (3)\\ 1.864\ (3)\\ 1.864\ (3)\\ 1.886\ (2)\\ 2.026\ (2)\\ 1.852\ (3)\\ 1.858\ (2)\\ 1.863\ (3)\\ \end{array}$	$\begin{array}{cccccccccccccccccccccccccccccccccccc$

Symmetry code: (i) -x+1, -y+1, -z+1.

### Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D····A	<i>D</i> —H··· <i>A</i>
01—H1…F1	0.84	1.81	2.569 (4)	149
C3—H5…F9 <sup>ii</sup>	0.98	2.58	3.427 (5)	145

			supporting information		
C3—H3…F8	0.98	2.59	3.437 (5)	145	
C2—H2…F10 <sup>iii</sup>	0.92 (4)	2.43 (4)	3.352 (5)	175 (3)	

Symmetry codes: (ii) -*x*+3/2, *y*+1/2, *z*; (iii) *x*-1/2, -*y*+1/2, -*z*+1.