

# A new name and a new synonym in *Miconia* (Melastomataceae)

Susanne S. Renner<sup>1</sup>, Renato Goldenberg<sup>2</sup>

**1** Herbaria and University of Munich, Menzingerstr. 67, Munich, Germany **2** Departamento de Botânica, Universidade Federal do Paraná, 81531-970, Curitiba, Paraná, Brazil

Corresponding author: *Susanne Renner* (renner@lrz.uni-muenchen.de)

---

Academic editor: *Hanno Schaefer* | Received 24 February 2011 | Accepted 7 April 2011 | Published 30 May 2011

**Citation:** Renner SS, Goldenberg R (2011) A new name and a new synonym in *Miconia* (Melastomataceae). *PhytoKeys* 3: 35–37. doi: 10.3897/phytokeys.3.1174

---

## Abstract

The name *Miconia densiflora* Cogn. (1886) is a later homonym of *M. densiflora* (Gardner) Naudin (1851), but since we propose it as a taxonomic synonym of *M. caudata* (Bonpl.) DC. (1828), we do not provide a new name. The name *Miconia longicuspis* Herzog (1909) is a later homonym of *M. longicuspis* Cogn. (1891) and we here propose its replacement by *M. longicuspidata* S.S. Renner & R. Goldenb.

## Keywords

Bolivia, Colombia, Melastomataceae, *Miconia*, homonyms, synonyms, taxonomy

## Introduction

The site Melastomataceae.net provides a portal to open-access databases dealing with Melastomataceae, a pan-tropical family of about 3500 species. Among the databases that can be accessed at this site is “MEL names,” which deals with the ca. 13,278 names of Melastomataceae and Memecylaceae (= Melastomataceae subfamily Olisbeoideae) and provides information on recognized species, synonyms, and relevant literature. In the course of dealing with the 1497 names available for *Miconia* Ruiz & Pav., a genus of at least 1061 accepted species, we discovered two homonymy problems, resolved here. Since one of the homonyms is actually a heterotypic synonym of another species, a replacement name is only needed for one of these species. A comprehensive taxonomic treatment of *Miconia* is currently in preparation, as part of the PBI Miconieae project ([sweetgum.nybg.org/melastomataceae/](http://sweetgum.nybg.org/melastomataceae/)).

## Systematics

### *Miconia longicuspidata* S.S.Renner & R.Goldenb., nom. nov.

urn:lsid:ipni.org:names:77111574-1

[http://species-id.net/wiki/Miconia\\_longicuspidata](http://species-id.net/wiki/Miconia_longicuspidata)

Replaced name: *Miconia longicuspis* Herzog, Feddes Repert. Nov. Sp. 7: 64. 1909.

TYPE: BOLIVIA. Cordillera de Santa Cruz: Cerro Amboró, Cuñucú, 600–1400 m, Oct 1907, T.Herzog 326 (holotype: Z!). Not *Miconia longicuspis* Cogn. in A.DC. & C.DC, Monog. Phan. 7: 850. 1891.

**Comments.** *Miconia longicuspis* Herzog is a later homonym of *Miconia longicuspis* Cogn., a treelet from eastern Brazil (Goldenberg and Reginato 2006). *Miconia longicuspidata* is known only from the holotype at Z, a collection by Theodor Herzog (1880–1961) in the mountains of Cuñucú, Bolivia, in 1907. We have found no isotypes or type photos in G, JE, L, U, W, or WAG. This species most closely matches *M. abbreviata* Markgr., a widespread shrub that ranges from Bolivia throughout the Amazon basin to the Guianas, but differs in the much denser secondary venation. In *M. abbreviata* the secondaries are spaced at about 0.5–0.7 mm apart, in *M. longicuspidata*, only 2–3 mm apart. Theodor Herzog was an expert mountaineer and collector, who explored widely in Bolivia. During his two expeditions to that country in 1907/08 and 1910/11, he collected about 600 species of vascular plants. In 1910, Herzog not only explored the Cordillera Santa Cruz, where *M. longicuspidata* appears to be endemic, but also surveyed the Cordillera de Cocabata, a range of peaks near Cochabamba extending northwest for about 100 kilometers. A description of the flora of the Cordillera de Santa Cruz is given in Herzog (1910). A color photograph of the type is available at <http://www.zuerich-herbarien.uzh.ch>.

### *Miconia caudata* (Bonpl.) DC., Prodr. 3: 187. 1828.

[http://species-id.net/wiki/Miconia\\_caudata](http://species-id.net/wiki/Miconia_caudata)

Basionym: *Melastoma caudata* Bonpl., Monogr. Melast. 1: 13. t. 7. 1807.

New synonym: *Miconia densiflora* Cogn., Bot. Jahrb. 8(1): 22, 1887 [1886]; et in A. DC. & C. DC. Monog. Phan. 7: 744, 1891, *syn. nov.* TYPE: COLOMBIA. Cauca: Popayan, Apr 1883, Lehmann 2798 (Isotype: US!). Non *Miconia densiflora* (Gardner) Naudin, Ann. Sc. Nat. Ser. 3, 16: 245, 1851, basionym *Chaenopleura densiflora* Gardner, Hook. Lond. Journ. Bot. 2: 349, 1843.

**Comments.** *Miconia densiflora* Cogn. is a later homonym of *Miconia densiflora* (Gardner) Naudin, which in turn is a taxonomic synonym of *Miconia pusilliflora* (DC.) Triana, a shrub from eastern Brazil (Cogniaux 1891; Goldenberg 2000). By contrast, Cogniaux' *M. densiflora* is based on a collection made by Lehmann near Popayán, in

the State of Cauca, Colombia, in 1883. The name is undoubtedly a taxonomic synonym of the common and frequently collected *Miconia caudata* (Bonpl.) DC. Cogniaux (1891: 736, 739) erroneously states that *M. caudata* has glandular-pubescent filaments, while *M. densiflora* has glabrous ones. However, all 15 specimens of *M. caudata* that we checked had completely glabrous filaments. A color photograph of *M. densiflora*'s type is available at <http://www.botany.si.edu/types>.

## Acknowledgements

The “Mel Names” project was funded by seed money from the “Electronic Catalogue of Names of Known Organisms” project organized by GBIF International 2005. We thank Werner Greuter for his advice and R. D. Stone for his careful review of our manuscript.

## References

- Cogniaux CA (1891) Melastomataceae. In: De Candolle A, De Candolle C (Eds) *Monographiae Phanerogamarum*, vol. 7. G. Masson, Paris, 1–1256.
- Goldenberg R (2000) O gênero *Miconia* Ruiz & Pav. (Melastomataceae). I. listagens analíticas, II. Revisão taxonômica da seção *Hypoxanthus* (Rich. ex DC.) Hook. f. PhD Thesis, Universidade Estadual de Campinas, Brazil.
- Goldenberg R, Reginato M (2006) Sinopse da família Melastomataceae na Estação Biológica de Santa Lúcia (Santa Teresa, Espírito Santo). *Boletim do Museu de Biologia Mello Leitão – Nova Série* 20: 33–58.
- Herzog T (1910) Pflanzenformationen Ostbolivias. *Botanische Jahrbücher für Systematik, Pflanzengeschichte und Pflanzengeographie* 44: 346–405.
- Melastomataceae.net: a site with information on the biodiversity of Melastomataceae. <http://www.melastomataceae.net>.