

# Lack of the Long Pentraxin PTX3 Promotes Autoimmune Lung Disease but not Glomerulonephritis in Murine Systemic Lupus Erythematosus

Maciej Lech<sup>1,9</sup>, Christoph Römmele<sup>1,9</sup>, Onkar P. Kulkarni<sup>1</sup>, Heni Eka Susanti<sup>1</sup>, Adriana Migliorini<sup>1</sup>, Cecilia Garlanda<sup>2</sup>, Alberto Mantovani<sup>2,3</sup>, Hans-Joachim Anders<sup>1\*</sup>

**1** Medizinische Poliklinik, University of Munich, Munich, Germany, **2** Istituto Clinico Humanitas, IRCCS, Rozzano, Italy, **3** Department of Translational Medicine, University of Milan, Rozzano, Italy

## Abstract

The long pentraxin PTX3 has multiple roles in innate immunity. For example, PTX3 regulates C1q binding to pathogens and dead cells and regulates their uptake by phagocytes. It also inhibits P-selectin-mediated recruitment of leukocytes. Both of these mechanisms are known to be involved in autoimmunity and autoimmune tissue injury, e.g. in systemic lupus erythematosus, but a contribution of PTX3 is hypothetical. To evaluate a potential immunoregulatory role of PTX3 in autoimmunity we crossed *Ptx3*-deficient mice with *Fas*-deficient (*lpr*) C57BL/6 (B6) mice with mild lupus-like autoimmunity. PTX3 was found to be increasingly expressed in kidneys and lungs of B6/*lpr* along disease progression. Lack of PTX3 impaired the phagocytic uptake of apoptotic T cells into peritoneal macrophages and selectively expanded CD4/CD8 double negative T cells while other immune cell subsets and lupus autoantibody production remained unaffected. Lack of PTX3 also aggravated autoimmune lung disease, i.e. peribronchial and perivascular CD3+ T cell and macrophage infiltrates of B6/*lpr* mice. In contrast, histomorphological and functional parameters of lupus nephritis remained unaffected by the *Ptx3* genotype. Together, PTX3 specifically suppresses autoimmune lung disease that is associated with systemic lupus erythematosus. Vice versa, loss-of-function mutations in the *Ptx3* gene might represent a genetic risk factor for pulmonary (but not renal) manifestations of systemic lupus or other autoimmune diseases.

**Citation:** Lech M, Römmele C, Kulkarni OP, Susanti HE, Migliorini A, et al. (2011) Lack of the Long Pentraxin PTX3 Promotes Autoimmune Lung Disease but not Glomerulonephritis in Murine Systemic Lupus Erythematosus. PLoS ONE 6(5): e20118. doi:10.1371/journal.pone.0020118

**Editor:** Mehrdad Matlobian, University of California, San Francisco, United States of America

**Received:** December 28, 2010; **Accepted:** April 25, 2011; **Published:** May 27, 2011

**Copyright:** © 2011 Lech et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Funding:** The work was supported by a grant from the Deutsche Forschungsgemeinschaft (AN372/11-1 and GRK 1202) to H.J.A. and European Commission (ERC project HHS to A. Mantovani). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing Interests:** The authors have declared that no competing interests exist.

\* E-mail: hjanders@med.uni-muenchen.de

<sup>9</sup> These authors contributed equally to this work.

## Introduction

Systemic lupus erythematosus (SLE) involves polyclonal autoimmunity against multiple nuclear autoantigens and presents clinically in a broad spectrum of manifestations ranging from mild fever, skin rashes, and arthralgia to severe inflammation of kidney, lungs, or brain [1]. It has become evident that SLE is not a single disease with a uniform trigger but rather a syndrome that can develop from many different causes [2]. The pathogenesis of SLE is largely based on variable combinations of genetic variants that promote loss-of-tolerance or tissue inflammation [2,3]. For example, some gene affect apoptosis, opsonization of dying cells, phagocytosis or the digestion of self-DNA which increase the exposure of nuclear particles to the immune system [4]. Another set of risk genes enhance the immune recognition of self nucleic acids by Toll-like receptors (TLR) in dendritic cells which increases the production of type I interferon [5,6] and eventually the expansion of autoreactive lymphocytes [7]. A third class of genetic lupus risk factors affects tissue inflammation [4].

Pentraxins belong to the first (and the third) group of molecules. The short pentraxins, C-reactive protein (CRP) and serum amyloid P (SAP), are acute phase proteins that are strongly

induced in hepatocytes in response to IL-6 [8]. CRP and SAP bind to all types of microorganisms, dead cells, and other particles and facilitate complement-mediated killing as well as uptake of the particle into phagocytes [8]. As such the short pentraxins foster the rapid clearance of pathogens and dead cells from the extracellular space [9]. The latter is particularly important in order to prevent an exposure of nuclear particles to the immune system [10]. In analogy to complement deficiency genetic lack of CRP or SAP is associated with impaired clearance of apoptotic cells and the onset of lupus [11,12,13]. Pentraxin gene polymorphisms are unlikely to account broadly for human SLE but, interestingly, serum CRP and SAP levels are usually low in the majority of lupus patients in the absence of infection despite significant SLE activity [14,15]. This has been attributed to anti-pentraxin antibodies as well as to the suppressive effect of IFN- $\alpha$  on the promoter activity of the short-pentaxins [16,17]. A recent study also examined serum levels of the long pentraxin PTX3 which were high in patients with all kinds of rheumatic diseases but remain low in patients affected by SLE [18,19]. Anti-PTX3 antibodies behaved the opposite way [18,19].

The long pentraxin PTX3, in contrast to the short pentraxins, is produced outside the liver by neutrophils, macrophages, myeloid

dendritic cells, as well as a number of non-immune cells in response to IL-1, TNF- $\alpha$ , and TLR agonists [20]. PTX3 shares some immunomodulatory functions with the short pentraxins such as binding to C1q and activation of the classical complement pathway [21], and inhibiting the amplification loop of the alternative complement pathway [22], and accelerating host defense to pathogens [8]. However, PTX3 seems to also have unique immunoregulatory functions such as modulating the phagocytic uptake of apoptotic cells by macrophages and dendritic cells [23,24,25,26], and interacting with P-selectin which inhibits leukocyte recruitment [27]. Although all of the aforementioned mechanisms might be involved in the pathogenesis of autoimmune diseases, a contribution of PTX3 to SLE is speculative to date. PTX3 might promote SLE via modulating the clearance of apoptotic cells or by driving complement-mediated tissue pathology. PTX3 might also protect from SLE manifestations by suppressing P-selectin-mediated leukocyte recruitment to affected organs.

In order to test whether *Ptx3* serves as a modifier gene on established SLE we generated *Ptx3*-deficient C57BL/6<sup>*lpr/lpr*</sup> (B6<sup>*lpr*</sup>) mice and compared the phenotype with wild type B6<sup>*lpr*</sup> mice, an autoimmune mouse strain that develop lupus autoantibodies but only mild SLE manifestations in kidneys and lung [28].

## Results

### Lack of PTX3 impairs the clearance of apoptotic cells

PTX3 was reported to regulate the C1q-mediated phagocytosis of apoptotic cells *in vitro* [21]. As impaired clearance of apoptotic cells is a well established pathomechanism of SLE [29] we first tested the ability of *Ptx3*-deficient mice to clear apoptotic cells by phagocytosis *in vivo*. PHrodo-labeled apoptotic cells were injected into the peritoneal cavities of *Ptx3*-deficient and wild type B6<sup>*lpr*</sup> mice. After 45 minutes peritoneal lavage fluids were analyzed by flow cytometry for pHrodo+ F4/80 macrophages. F4/80+ peritoneal macrophages of *Ptx3*-deficient mice displayed a significantly reduced capacity to take up apoptotic cells as compared to wild type B6<sup>*lpr*</sup> mice (Figure 1). Thus, lack of PTX3 is associated with a reduced clearance of apoptotic cells.

### PTX3 expression in autoimmune B6<sup>*lpr*</sup> mice

Next we characterized the expression of PTX3 mRNA in solid organs of 6 week old B6<sup>*lpr*</sup> mice. PTX3 mRNA was highly expressed in bone marrow (Figure 2A). Among the solid organs PTX3 mRNA levels were much higher heart and lungs as compared to the kidneys and the urinary bladder. PTX3 protein expression was confirmed by Western blot in spleens, kidneys and lungs of B6<sup>*lpr*</sup> mice. Spleen and kidney PTX3 expression decreased over time when autoimmunity progresses in B6<sup>*lpr*</sup> mice (Figure 2B). By contrast, PTX3 protein expression increased in lungs at 6 months of age in B6<sup>*lpr*</sup> mice indicating local production of PTX3 in lungs which are usually affected by autoimmune tissue injury in B6<sup>*lpr*</sup> mice. Thus, PTX3 is increasingly expressed in lungs during the progression of autoimmunity of B6<sup>*lpr*</sup> mice.

### PTX3 suppresses lymphoproliferation in B6<sup>*lpr*</sup> mice

Next we generated *Ptx3*-deficient B6<sup>*lpr*</sup> mice. The autoimmune phenotype of homozygous B6<sup>*lpr*</sup> mice is introduced only by mutation of a single lupus susceptibility gene (*lpr*) which impairs Fas-induced apoptosis of autoreactive B and T cells [28]. Because B6<sup>*lpr*</sup> mice develop only mild autoimmune syndrome, litters of B6<sup>*lpr/Ptx3*-/-</sup> mice could be bred along Mendelian ratios from B6<sup>*lpr/Ptx3*+/-</sup> mice and revealed no differences in body weight gain between the two genotypes (Figure 3A). For SLE phenotype

analysis we first evaluated the size of spleens and lymph nodes in 6 months old B6<sup>*lpr*</sup> and B6<sup>*lpr/Ptx3*-/-</sup> mice. Spleens and lymph nodes were significantly enlarged in B6<sup>*lpr/Ptx3*-/-</sup> mice as compared to B6<sup>*lpr*</sup> mice. This was evident from spleen and cervical lymph node weights (Figure 3B).

### PTX3 specifically suppresses CD4/CD8 double negative T cells in B6<sup>*lpr*</sup> mice

Because we had found *Ptx3*-deficiency impaired the rapid clearance of apoptotic cells we hypothesized that sustained exposure to dead cells would modulate the activation of dendritic cells, cells that handle lupus autoantigens and drive the expansion of autoreactive lymphocytes in SLE, like in E8-Mag or DNase 1-deficient mice [30,31]. We performed flow cytometry to quantify and characterize the activation state of CD11c+ dendritic cells without additional stimuli directly after the spleen harvest at 6 months of age. However, the numbers of CD11c+/CD40+ or the CD11b/MHCII+ cells were identical as well as the total numbers of monocytes or neutrophils in both genotypes (Figure 4A). Consistent with this finding serum levels of IL-12 were unaffected by the PTX3 genotype (Figure 4B). Does lack of PTX3 affect T cell populations in B6<sup>*lpr*</sup> mice? The numbers of CD4/CD8 double negative 'autoreactive' T cells were increased in *Ptx3*-deficient B6<sup>*lpr*</sup> mice (Figure 4C). In contrast, the numbers of all other T cell subsets, i.e. CD4+, CD8+, and CD4+/CD25+ T cells were comparable between the two genotypes (Figure 4C). Thus, PTX3-deficiency is associated with a selective expansion of CD4/CD8 double negative T cells in B6<sup>*lpr*</sup> mice.


### PTX3 does not regulate B cell expansion and autoantibody production in B6<sup>*lpr*</sup> mice

Flow cytometry did not reveal any difference in numbers of mature B cells, follicular B cells, marginal zone B cells, and plasma cells in spleens (Figure 5A) which was consistent with the comparable size of the IgM+ plasma cell areas in spleens of B6<sup>*lpr*</sup> and B6<sup>*lpr/Ptx3*-/-</sup> mice (Figure 5B). Consistent with the numbers of B cells, *Ptx3*-deficient B6<sup>*lpr*</sup> mice displayed similar serum IgG levels as compared to 6 months old B6<sup>*lpr*</sup> wild-type mice (Figure 6A). At that time lack of PTX3 did also had no effect on serum dsDNA autoantibody levels from total IgG, IgG1, IgG2a/c, and IgG3. The specificity of dsDNA autoantibodies was confirmed by using the *Critidia luciliae* assay. Diluted serum from B6<sup>*lpr/Ptx3*-/-</sup> mice showed comparable binding to the dsDNA of the flagellate's kinetoplast as serum of B6<sup>*lpr*</sup> mice (not shown). In addition, lack of PTX3 did not affect the levels of anti-Sm IgG, anti-U1snRNP IgG, and rheumatoid factor as compared to B6<sup>*lpr*</sup> mice (Figure 6B). In addition, *Ptx3*-deficient B6 mice did not reveal any sign of spontaneous autoimmunity, e.g. autoantibodies against ANA, dsDNA or rheumatoid factor up to 12 months of age. Thus, PTX3 is redundant for the expansion of B cells and plasma cells as well as for the production of numerous autoantibodies against nuclear autoantigens in B6<sup>*lpr*</sup> mice lack of PTX3 alone does not induce autoimmunity against DNA.


### PTX3 suppresses autoimmune lung but not kidney disease in B6<sup>*lpr*</sup> mice

SLE may be associated with little or severe autoimmune tissue injury [32]. B6<sup>*lpr*</sup> mice do not develop major autoimmune tissue injuries although mild glomerulonephritis develops from 6 months of age [28]. *Ptx3*-deficient B6<sup>*lpr*</sup> mice revealed significant peribronchial and perivascular neutrophils, CD3+ T cell infiltrates accompanied by Mac2 + macrophages while significant pulmonary pathology was absent in age-matched B6<sup>*lpr*</sup> mice (Figure 7,




A



B




C




**Figure 1. PTX3 fosters the phagocytic uptake of apoptotic cells.** A:  $2 \times 10^5$  pHrodo-labeled apoptotic cells were injected intraperitoneally into *B6/prPtx3*-deficient and *B6/pr* wildtype mice and 45 min later peritoneal lavage fluids were prepared for flow cytometry. Figure A displays the percentage of F4/80+ peritoneal macrophages of all cells in lavage fluids which were not different in *B6/pr* (black bar) from *B6/pr Ptx3*-deficient mice (white bar). Figure B displays the number of F4/80+ cells also positive for pHrodo, a dye which gets only activated in the acidic environment of phagolysosomes. Data represent means  $\pm$  SEM from three independent experiments; \*  $p < 0.05$  versus wild type. Figure C displays representative dot plots from this experiment. C: Representative density plots of F4/80+Phrodo+ cells in *B6/prPtx3*-deficient and *B6/pr* wildtype mice. doi:10.1371/journal.pone.0020118.g001

8A and 8B). Given the known role of PTX3 for P-selectin-mediated lung leukocyte recruitment we next determined the mRNA expression levels of P-, E-, and L-selectin, PECAM, IP10 and Cxcr3 in lungs of both mouse strains. Lack of PTX3 was associated with increased mRNA expression of P-selectin and E-selectin (but not L-selectin, PECAM, IP10 or Cxcr3) in lungs of 6 months old *B6<sup>lpr</sup>* mice (Figure 8C). Consistent with mRNA data,

lack of PTX3 was associated with increased P-selectin expression on the protein level (Figures 8D and 8E). In contrast, the *Ptx3* Genotype had no effect on the activity of lupus nephritis, proteinuria, renal P-selectin and E-selectin mRNA expression, complement immunostaining, and renal leukocyte numbers (Figure 9). Together, PTX3 protects *B6<sup>lpr</sup>* mice from autoimmune lung disease but not from lupus nephritis



**Figure 2. PTX3 expression in mice.** A: RNA was isolated from organs of 6 week old *B6<sup>lpr</sup>* mice for real-time RT-PCR. Data are expressed as means of the ratio of the specific mRNA versus that of 18S rRNA  $\pm$  SEM. B: Protein samples were prepared from spleens, kidneys, and lungs of *B6<sup>lpr</sup>* mice at 1, 3, and 6 months of age. PTX3 Western blot indicates the quantitative (20  $\mu$ g protein load per lane) PTX3 protein expression in each organ over time. The histogram represents the ratio of PTX3 expression to the expression of the  $\beta$ -actin loading control. Data represent means  $\pm$  SEM from three independent experiments. doi:10.1371/journal.pone.0020118.g002




**Figure 3. PTX3 and lymphoproliferation in B6<sup>lpr</sup> mice.** A: Body weight increased similarly over time in *Ptx3*-deficient or wild type B6<sup>lpr</sup> mice. Data are means  $\pm$  SEM from at least 18 mice in each group. B: At 6 months of age *Ptx3*-deficient B6<sup>lpr</sup> mice revealed splenomegaly and hyperplasia of cervical lymph nodes as compared to age-matched B6<sup>lpr</sup> control mice. Quantitative data on spleen and lymphnode weights are means  $\pm$  SEM from at least 18 mice in each group, \*  $p < 0.05$  versus B6<sup>lpr</sup> mice. doi:10.1371/journal.pone.0020118.g003

## Discussion

The long pentraxin PTX3, like the short pentraxins, has multiple regulatory roles on innate immunity. It modulates opsonization (including dead cell clearance), complement activation, and leukocyte recruitment, all processes that affect autoimmunity and autoimmune tissue injury [8]. The experimental strategy of generating *Ptx3*-deficient autoimmune B6<sup>lpr</sup> mice intended to test the role of *Ptx3* as a potential modifier gene for established autoimmunity. Our data now demonstrate that the evolution of autoimmunity and lupus autoantibodies in B6<sup>lpr</sup> mice seems to be independent of PTX3 but that PTX3 has a non-redundant role in suppressing autoimmune lung injury.

Interstitial or alveolar lung disease is rare in humans and usually absent in 6 months old B6<sup>lpr</sup> mice [33] but is well described in MRL<sup>lpr</sup> mice which suffer from more advanced lupus-like autoimmune tissue injuries at this age [34]. The peribronchial and perivascular lymphocyte infiltrates that we observed in B6<sup>lpr</sup>/*Ptx3*<sup>-/-</sup> mice were similar to those reported from MRL<sup>lpr</sup> mice and to those that we had previously observed in B6<sup>lpr</sup> mice with accelerated SLE [35]. Hence, lack of PTX3 specifically acceler-

ated the evolution of autoimmune lung disease, albeit not kidney disease, in B6<sup>lpr</sup> mice. The knowledge about organ-specific pathomechanisms in SLE is limited. Kidney disease in autoimmune mice (and lupus patients) mainly develops from immune complex disease and depends on glomerular complement activation and macrophage recruitment [36]. By contrast, lung disease in MRL<sup>lpr</sup> mice involves the recruitment of CXCR3 positive T cells via local secretion of CXCL10 [37], and the endothelial expression of selectins [38] and intercellular adhesion molecule (ICAM)-1 [39]. In addition, TNF- $\alpha$  is a crucial mediator of lung injury in experimental lupus [40,41]. Unspecific immunosuppressants like cyclophosphamide or dihydroorotate dehydrogenase inhibitors suppress pulmonary and renal manifestations in experimental lupus [42,43]. However, the aforementioned molecular and cellular pathomechanisms of autoimmune lung injury differ, at least in part, from those of lupus nephritis because *Icam-1*-deficiency as well as TNF- $\alpha$  antagonism protects MRL<sup>lpr</sup> mice from lung but not from kidney disease [39,41]. Furthermore, P-selectin is only induced in lungs but not in kidneys upon immune stimulation with LPS [38] or, as we found here, in experimental SLE. The latter finding is of particular interest because PTX3 was




**Figure 4. Lack of PTX3 specifically affects T cells subsets in B6<sup>lpr</sup> mice.** A: Spleen monocytes, neutrophils, CD11b+ and CD11c+ cells were quantified in B6<sup>lpr/Ptx3<sup>-/-</sup></sup> and B6<sup>lpr</sup> wild-type mice by flow cytometry also using surface MHCII and CD40 expression as a marker of cell activation as described in methods. Data represent means  $\pm$  SEM from 10 mice in each group. B: Serum IL-12p40 levels were determined in 6 months old B6<sup>lpr</sup> of both genotypes by ELISA and are shown as dot blot. C: Flow cytometry was used to determine the total number of distinct T cell subsets in spleens of 6 months old B6<sup>lpr/Ptx3<sup>-/-</sup></sup> and B6<sup>lpr</sup> wild type mice. The graphs in B–C present means  $\pm$  SEM of 8–14 mice in each group \*  $p < 0.05$ . doi:10.1371/journal.pone.0020118.g004

recently identified as an endogenous P-selectin inhibitor that limits leukocyte recruitment to the lung [27]. We therefore assume that the phenotype of *Ptx3*-deficient B6<sup>lpr</sup> mice underscores the organ-specific immunopathology in SLE and might relate to the potential of PTX3 to specifically impair leukocyte recruitment to the lung, e.g. by interacting with P-selectin.

Surprisingly, *Ptx3*-deficiency only marginally affected systemic autoimmunity in B6<sup>lpr</sup> mice. The *lpr* mutation induces autoimmunity by not allowing immature and potentially autoreactive T lymphocytes to undergo Fas-mediated apoptosis [28]. As a consequence CD4/CD8 double negative T cells accumulate and eventually undergo secondary necrosis which increases the exposure of potential lupus autoantigens to phagocytes and antigen-presenting cells [28,44]. Our data clearly demonstrate that lack of PTX3 is associated with an additional expansion of CD4/CD8 double negative cells indicating that PTX3 has a non-redundant role in suppressing the expansion of this cell population in B6<sup>lpr</sup> mice. How PTX3 suppresses the expansion of the CD4/CD8 double negative T cells is less clear. Three studies consistently reported that soluble PTX3 impairs the capacity of cultured

macrophages and dendritic cells to take up apoptotic cells which was concluded to counterbalance the opsonizing effect on dead cells of the short pentraxins [23,25,26]. A more recent study reported the opposite effect for membrane-bound PTX3 on apoptotic neutrophils which acts as an ‘eat me’ signal and fosters phagocytic clearance by macrophages [24]. The results from our own in-vivo phagocytosis assay are consistent with the latter report and document that PTX3 fosters the rapid clearance of apoptotic T cells by peritoneal macrophages, a process that may keep lupus autoantigens away from dendritic cells and avoid the activation of autoreactive T cells. Vice versa, impaired apoptotic cell clearance by antigen-presenting cells may serve as a stimulus for the selective expansion of CD4/CD8 double negative ‘autoreactive’ T cells in our study. It is of note that the numbers of CD4/CD8 double negative cells or their ratio to CD4/CD25/Foxp3 regulatory T cells are usually taken as a surrogate marker of the autoimmune process in experimental lupus [45,46]. However, in our study the isolated expansion of the CD4/CD8 double negative T cells population was not associated with more lupus autoantibody production which raises doubts about their causal role for systemic autoimmunity.



**Figure 5. PTX3 and B cell subsets in B6<sup>lpr</sup> mice.** A: Flow cytometry was used to determine the total number of distinct B cell subsets in spleens of 6 months old B6<sup>lpr</sup>/Pttx3<sup>-/-</sup> and B6<sup>lpr</sup> wild type mice. The histogram presents means  $\pm$  SEM of 8–14 mice in each group. B: Spleens from the same mice were stained for IgM to localize plasma cell areas. The images are representative for 6 mice in each group. Original magnification  $\times$ 100. doi:10.1371/journal.pone.0020118.g005

Together, the long pentraxin PTX3 is required to suppress lung disease in systemic autoimmunity. PTX3 does not regulate lupus nephritis of B6<sup>lpr</sup> mice. Furthermore, PTX3 is redundant for the production of lupus autoantibodies in these mice, albeit it fosters the clearance of apoptotic cells and (thereby) inhibits the expansion of CD4/CD8 double negative T cells. These results add on to previous data that have documented organ-specific pathomechanisms for lupus manifestations. In addition, it is now intriguing to speculate that loss-of-function mutations in the *Ptx3* gene might represent a genetic risk factor for pulmonary manifestations of human SLE or that recombinant PTX3 or other PTX3 agonists might have the potential to specifically suppress autoimmune lung disease.

## Materials and Methods


### Animal studies

*Ptx3*-deficient mice were generated as previously described [47] and backcrossed to the C57BL/6 strain (B6, CharlesRiver Laboratories, Calco, Italy) to the N11 generation. B6<sup>Ptx3<sup>-/-</sup></sup> and B6<sup>lpr</sup> mice (Charles River) were mated to generate B6<sup>lpr</sup>/Pttx3<sup>-/+</sup> mice which were then mated among each other to generate B6<sup>lpr</sup>/Pttx3<sup>+/+</sup> and B6<sup>lpr</sup>/Pttx3<sup>-/-</sup> mice as described [35]. Littermate female were used for all experimental procedures. In each

individual mouse the genotype was assured by PCR. Mice were housed in groups of 5 mice in sterile filter top cages with a 12 hour dark/light cycle and unlimited access to autoclaved food and water. One cohort of mice was sacrificed by cervical dislocation at 24 weeks of age. All experimental procedures were performed according to the German animal care and ethics legislation and had been approved by the local government authorities (Regierung von Oberbayern, Az 55.2-1-54-2531-11-10).

### Phagocytosis assay and generation of apoptotic cells

EL4 cells ( $1 \times 10^6$ /ml) were incubated with 90  $\mu$ M of the DNA topoisomerase I inhibitor Camptothecin (Calbiochem, San Diego, USA) for 4 h at 37°C to induce apoptosis. Apoptosis was confirmed by flow cytometry after staining with FITC Annexin V Apoptosis Detection Kit 1 (BD Biosciences Pharmingen, San Diego, CA). After 4 h the apoptotic cells ( $1 \times 10^6$ /ml) were stained with 125 ng/ $\mu$ l pHrodo (Invitrogen, Eugene, OR). Mice were injected intraperitoneally with 500  $\mu$ l of 4% thioglycollate medium (BD, Franklin Lakes, USA). After 73 h of incubation the mice were injected intraperitoneally with 200  $\mu$ l of the apoptotic pHrodo-stained apoptotic EL4 cells. 40 min later peritoneal fluid were centrifuges and resuspended in HBSS (PAN-Biotech GmbH, Aldenbach, Germany) +4 mM EDTA (Biochrom KG, Berlin, Germany). The cells were stained with anti-



**Figure 6. PTX3 and humoral immunity in lupus of B6<sup>lpr</sup> mice.** B6<sup>lpr/Ptx3<sup>-/-</sup></sup> (□) and B6<sup>lpr</sup> wild-type mice (■) were bled at monthly intervals to determine serum levels of IgG (A). Data represent means  $\pm$  SEM from at least 12 mice at each time point and genotype. B: DsDNA autoantibodies and various lupus autoantibodies were analysed by ELISA at 6 months only and are shown as dot blots. 6 mice at each time point and genotype. No significant differences were detected between the two genotypes for any of the parameters. doi:10.1371/journal.pone.0020118.g006

F4/80 IgG (BD Pharmingen, Heidelberg, Germany) for flow cytometry analysis.

### Flow cytometry

Anti-mouse CD3, CD4, CD8 and CD25 antibodies (BD Pharmingen, Heidelberg, Germany) were used to detect CD3+CD4<sup>-</sup>CD8<sup>-</sup> double negative T cells and CD4+CD25+ regulatory T cells populations in spleens. Anti-CD11c was used to identify dendritic cells and their activation was assed by co-staining for CD40 (BD Pharmingen, Heidelberg, Germany). Anti-mouse B220, CD21, CD23, IgD, IgM antibodies (BD Pharmingen, Heidelberg, Germany) were used to detect mature B cells (B220+IgD+IgM<sup>+</sup>), marginal zone B cells (B220+CD21<sup>high</sup>CD23<sup>low</sup>) and follicular B cells (B220+CD21<sup>low</sup>CD23<sup>high</sup>). Plasma cells were identified by using anti-mouse antibodies for Ig  $\kappa$  light chain and CD138 (BD Pharmingen, Heidelberg, Germany). Respective isotype antibodies were used to demonstrate specific staining of cell subpopulations [35]. Quantification of cell number was done using counting beads for FACS (Invitrogen).

### Evaluation of autoimmune tissue injury


Lungs, spleens and kidneys from all mice were fixed in 10% buffered formalin, processed, and embedded in paraffin. Two  $\mu\text{m}$  sections for periodic acid-Schiff (PAS) stains were prepared following routine protocols [48]. The severity of the renal lesions was graded using the activity and chronicity indices for human

lupus nephritis as described [49]. Autoimmune lung injury was scored semiquantitatively (0–4) by assessing the extent of peribronchial, perivascular or interstitial lymphocyte infiltrates as described [34]. For lung immunostaining: anti-mCD3e (clone 500A2, 1:50), Mac2 (macrophages, Cederlane, Ontario, Canada, 1:5000), rat anti-mouse neutrophils (Serotec, Oxford, UK, 1:50) were used. For IgM staining of the spleen sections, anti-mouse IgM-mu-chain specific antibodies (Vector, Burlingame, CA) were used. The slides were scanned using an Olympus BX 61 microscope and recorded via CellP software.

Serum IL-12 levels were determined by ELISA following the manufacturer's protocols (OptEiA, BD, Biosciences, Heidelberg, Germany).

### Ptx3 protein expression analysis

Western blot analysis were performed from kidney, lung and spleen protein extracts, which were incubated in two times loading buffer for 5 min at 95°C, resolved by 10% SDS-PAGE, and transferred to an Immobilon-P membrane (Millipore, Eschborn, Germany). After blocking with 1% Western Blocking Reagent (Roche, Germany), the filter was incubated with Rat anti mouse Ptx3 antibody (1:1000, Alpha Diagnostic International, San Antonio USA) overnight in TBS;  $\beta$ -actin (1:1000, Cell Signaling Technology, Beverly, MA) for 1 hour), anti-P-selectin monoclonal antibody (mAb) RB40 (anti-P-selectin were prepared from hybridoma cells). Immune complexes were



**Figure 7. PTX3 and lung injury in B6<sup>pr</sup> mice.** Lung sections were stained either with PAS or with specific antibodies for neutrophils, macrophages or T lymphocytes. Images are representative for at least 12 mice in each group. Original magnification  $\times 100$  (PAS) and  $\times 200$  (immunostaining).


doi:10.1371/journal.pone.0020118.g007

visualized using a peroxidase-conjugated anti-rat IgG Ab (1:10000, Cell Signaling Technology, Beverly, MA) for 1 h and processed for detection by ECL (Amersham Pharmacia Biotech Europe, Freiburg, Germany).

#### Autoantibody analysis

Serum antibody levels were determined by ELISA as described [50,51]. Anti-dsDNA antibodies: NUNC maxisorp ELISA plates were coated with poly-L-lysine (Trevigen, Gaithersburg, MD, USA) and mouse embryonic stem cell dsDNA. After incubation with mouse serum, dsDNA-specific IgG, IgG<sub>1</sub>, IgG<sub>2a/c</sub>, IgG<sub>2b</sub>,

IgG<sub>3</sub> and serum IgG levels were detected by ELISA (Bethyl Labs, Montgomery, TX, USA). *Critidia luciliae* assay: 1:50 diluted serum was applied to fixed *C. luciliae* slides (BioRad Laboratories, Redmond USA). Binding to *C. luciliae* kinetoplast was detected with FITC-conjugated goat anti-mIgG (1:1000, Invitrogen, Oregon USA). DAPI staining (Vector Laboratories, Burlingame CA) allowed colocalization with kinetoplast dsDNA. For quantitation of kinetoplast staining intensity a semiquantitative score from 0–3 was used. Anti-Sm: NUNC maxisorp ELISA plates were coated with Smith (Sm) antigen (Immunovision, Springdale, AR). A horseradish peroxidase-conjugated goat anti-mouse IgG (Rock-




**Figure 8. Lung damage and gene expression in B6<sup>lpr</sup> mice.** A: Peribronchial and interstitial inflammation was scored from 0–4 as described in methods. Data represent means  $\pm$  SEM of 8–12 mice in each group. B: Neutrophils, macrophages or T lymphocytes numbers were quantify, data represent means  $\pm$  SEM from at least 12 mice in each group, \*  $p < 0.05$ , \*\*  $p < 0.005$ , \*\*\*  $p < 0.0005$  versus B6<sup>lpr</sup> mice. C: RNA was isolated from lungs of 6 months old B6<sup>lpr/Ptx3<sup>-/-</sup></sup> and B6<sup>lpr</sup> wild-type mice for real-time RT-PCR. Data are expressed as means of the ratio of the specific mRNA from at least 5 mice versus that of 18S rRNA  $\pm$  SEM. \*  $p < 0.05$  versus B6<sup>lpr</sup> mice. D: Lung sections were stained with anti-P-selectin P. Images are representative for at least 12 mice in each group. Original magnification  $\times 100$ . E: Protein samples were prepared from lungs of B6<sup>lpr</sup> or B6<sup>Ptx3<sup>-/-</sup>lpr</sup> mice. PTX3 Western blot indicates the quantitative (20  $\mu$ g protein load per lane) P-selectin protein expression. The histogram represents the ratio of P-selectin to the respective  $\beta$ -actin expression. Data represent means  $\pm$  SEM from three independent experiments. doi:10.1371/journal.pone.0020118.g008

land, Gilbertsville, PA) was used for detection. The same procedure was followed for anti-SmRNP and anti-nucleosome antibodies as for anti-Sm except the ELISA plates were captured with Sm-RNP complex (Immunovision) or dsDNA together with histones (USB Corporation, Ohio USA) respectively instead of Sm antigen. Rheumatoid factor: ELISA plates were coated with 10  $\mu$ g/ml IgG (Jackson ImmunoResearch, West Grove, PA, USA) overnight at 4°C. C57BL/6 10 week mouse serum was used as negative control. HRP conjugated anti-mouse IgG was used as secondary antibody.

### Real-time quantitative RT-PCR

Real time RT-PCR was performed on mRNA from mouse organs as previously described [52]. SYBR Green Dye detection system was used for quantitative real-time PCR on Light Cycler 480 (Roche, Mannheim, Germany). All the technical steps were performed according to The Minimum Information for Publication of Quantitative Real-Time PCR Experiments (MIQE) guidelines [53]. For relative quantification the  $2^{-\Delta\Delta Ct}$  analysis method was used. Controls consisting of ddH<sub>2</sub>O were negative for target and housekeeper genes. Gene-specific primers



**Figure 9. PTX3 and lupus nephritis in *B6<sup>lpr</sup>* mice.** A: Kidney sections were stained with PAS. Images are representative for at least 12 mice in each group, original magnification  $\times 400$ . B: The composite activity index of lupus nephritis (0–24) was assessed as described in methods. Data represent means  $\pm$  SEM of 8–12 mice in each group. C: Urinary albumin/creatinine ratios were determined at 6 months as a functional marker of glomerular damage. D: Renal P-selectin and E-selectin mRNA levels were determined by real-time RT-PCR and are expressed as means of the ratio of the specific mRNA versus that of 18S rRNA  $\pm$  SEM (n = 12 in each group). E: Neutrophils, macrophages or T lymphocytes numbers were quantify, data represent means  $\pm$  SEM from at least 12 mice in each group. F: C9 complement staining was performed on kidney sections. No significant differences were detected between the two genotypes. doi:10.1371/journal.pone.0020118.g009

(300 nM, Metabion, Martinsried, Germany) were designed to be cDNA specific and to target possibly all known transcripts of gene of interest. In silico specificity screen (BLAST) was performed. PTX3: ID NM\_008987 right: 5'-CCTGCTTTGTGCTCTC TGGT-3', left: 5'-TCTCCAGCATGATGAACAGC-3'; E-Selectin: ID NM\_011345 right: 5'-TCTATTTCCCACGATG-CATTT-3', left: 5'-CTGCCAAAGCCTTCAATCAT-3'; L-Selectin: ID NM\_011346 right: 5'-TTCATGGCTTTCCTTT-CACA-3'; left: 5'-CTGGCATTCTCATT TGGCT-3'; P-

Selectin: ID NM\_011347 right: 5'-GGACACTTGATGGCTT-CACA-3', left: 5'-CAGTTCATGTGCGATGAAGG-3'; PE-CAM-1: ID NM\_001032378 right: 5'-TCCTTCCTGCTTC-TTGCTAGCT-3', left: 5'-GAGCCCAATCACGTTTCAGT-TT-3'; IP-10: ID NM\_021274 right: 5'-GGCTGGTC-ACTTTTCAGAAG-3', left: 5'-ATGGATGGACAGCAGA-GAGC-3'; Cxcr3: ID NM\_009910, right: 5'- TCTCGT-TTCCCATAATCG-3', left: 5'-AGCCAAGCCATGTACC-TTGA-3'.

## Statistical analysis

One-way ANOVA followed by post-hoc Bonferroni's test was used for multiple comparisons using GraphPad Prism 4.03 version. Single groups were compared by unpaired two-tailed Students t-test or non-parametric Mann-Whitney test. Data were expressed as mean  $\pm$  SEM. Statistical significance was assumed at a p value of  $<0.05$ .

## Acknowledgments

The expert technical assistance of Dan Draganovic and Janina Mandelbaum is gratefully acknowledged. Parts of this work were performed as a

medical thesis project by C.R. at the Medical Faculty of the University of Munich.

## Author Contributions

Conceived and designed the experiments: ML CR H-JA. Performed the experiments: ML CR OPK HES A. Migliorini. Analyzed the data: ML CR OPK H-JA. Contributed reagents/materials/analysis tools: CG A. Montovani. Wrote the paper: ML CR H-JA.

## References

- Kotzin BL (1996) Systemic lupus erythematosus. *Cell* 85: 303–306.
- Goodnow CC (2007) Multistep pathogenesis of autoimmune disease. *Cell* 130: 25–35.
- Gregersen PK, Olsson LM (2009) Recent advances in the genetics of autoimmune disease. *Annu Rev Immunol* 27: 363–391.
- Kanta H, Mohan C (2009) Three checkpoints in lupus development: central tolerance in adaptive immunity, peripheral amplification by innate immunity and end-organ inflammation. *Genes Immun* 10: 390–396.
- Theofilopoulos AN, Baccala R, Beutler B, Kono DH (2005) Type I interferons (alpha/beta) in immunity and autoimmunity. *Annu Rev Immunol* 23: 307–336.
- Hom G, Graham RR, Modrek B, Taylor KE, Ortmann W, et al. (2008) Association of systemic lupus erythematosus with C8orf13-BLK and ITGAM-ITGAX. *N Engl J Med* 358: 900–909.
- Marshak-Rothstein A, Rifkin IR (2007) Immunologically active autoantigens: the role of toll-like receptors in the development of chronic inflammatory disease. *Annu Rev Immunol* 25: 419–441.
- Bottazzi B, Doni A, Garlanda C, Mantovani A (2010) An integrated view of humoral innate immunity: pentraxins as a paradigm. *Annu Rev Immunol* 28: 157–183.
- Bijl M, Horst G, Bijzet J, Bootsma H, Limburg PC, et al. (2003) Serum amyloid P component binds to late apoptotic cells and mediates their uptake by monocyte-derived macrophages. *Arthritis Rheum* 48: 248–254.
- Anders HJ (2009) Pseudoviral immunity - a novel concept for lupus. *Trends Mol Med* 15: 553–561.
- Bickerstaff MC, Botto M, Hutchinson WL, Herbert J, Tennent GA, et al. (1999) Serum amyloid P component controls chromatin degradation and prevents antinuclear autoimmunity. *Nat Med* 5: 694–697.
- Russell AI, Cunningham-Graham DS, Shepherd C, Robertson CA, Whittaker J, et al. (2004) Polymorphism at the C-reactive protein locus influences gene expression and predisposes to systemic lupus erythematosus. *Hum Mol Genet* 13: 137–147.
- Truedsson L, Bengtsson AA, Sturfelt G (2007) Complement deficiencies and systemic lupus erythematosus. *Autoimmunity* 40: 560–566.
- Voss A, Nielsen EH, Svehag SE, Junker P (2008) Serum amyloid P component-DNA complexes are decreased in systemic lupus erythematosus: inverse association with anti-dsDNA antibodies. *J Rheumatol* 35: 625–630.
- Becker GJ, Waldburger M, Hughes GR, Pepys MB (1980) Value of serum C-reactive protein measurement in the investigation of fever in systemic lupus erythematosus. *Ann Rheum Dis* 39: 50–52.
- Shoenfeld Y, Szyper-Kravitz M, Witte T, Doria A, Tsutsumi A, et al. (2007) Autoantibodies against protective molecules—C1q, C-reactive protein, serum amyloid P, mannose-binding lectin, and apolipoprotein A1: prevalence in systemic lupus erythematosus. *Ann N Y Acad Sci* 1108: 227–239.
- Enocsson H, Sjowall C, Skogh T, Eloranta ML, Ronnblom L, et al. (2009) Interferon-alpha mediates suppression of C-reactive protein: explanation for muted C-reactive protein response in lupus flares? *Arthritis Rheum* 60: 3755–3760.
- Hollan I, Bottazzi B, Cuccovillo I, Forre OT, Mikkelsen K, et al. (2010) Increased levels of serum pentraxin 3, a novel cardiovascular biomarker, in patients with inflammatory rheumatic disease. *Arthritis Care Res (Hoboken)* 62: 378–385.
- Bassi N, Ghirardello A, Blank M, Zampieri S, Sarzi-Puttini P, et al. (2010) IgG anti-pentraxin 3 antibodies in systemic lupus erythematosus. *Ann Rheum Dis* 69: 1704–1710.
- Garlanda C, Bottazzi B, Bastone A, Mantovani A (2005) Pentraxins at the crossroads between innate immunity, inflammation, matrix deposition, and female fertility. *Annu Rev Immunol* 23: 337–366.
- Nauta AJ, Bottazzi B, Mantovani A, Salvatori G, Kishore U, et al. (2003) Biochemical and functional characterization of the interaction between pentraxin 3 and C1q. *Eur J Immunol* 33: 465–473.
- Deban L, Jarva H, Lehtinen MJ, Bottazzi B, Bastone A, et al. (2008) Binding of the long pentraxin PTX3 to factor H: interacting domains and function in the regulation of complement activation. *J Immunol* 181: 8433–8440.
- Baruah P, Dumitriu IE, Peri G, Russo V, Mantovani A, et al. (2006) The tissue pentraxin PTX3 limits C1q-mediated complement activation and phagocytosis of apoptotic cells by dendritic cells. *J Leukoc Biol* 80: 87–95.
- Jailion S, Jeannin P, Hamon Y, Fremaux I, Doni A, et al. (2009) Endogenous PTX3 translocates at the membrane of late apoptotic human neutrophils and is involved in their engulfment by macrophages. *Cell Death Differ* 16: 465–474.
- Rovere P, Peri G, Fazzini F, Bottazzi B, Doni A, et al. (2000) The long pentraxin PTX3 binds to apoptotic cells and regulates their clearance by antigen-presenting dendritic cells. *Blood* 96: 4300–4306.
- van Rossum AP, Fazzini F, Limburg PC, Manfredi AA, Rovere-Querini P, et al. (2004) The prototypic tissue pentraxin PTX3, in contrast to the short pentraxin serum amyloid P, inhibits phagocytosis of late apoptotic neutrophils by macrophages. *Arthritis Rheum* 50: 2667–2674.
- Deban L, Russo RC, Sironi M, Moalli F, Scanziani M, et al. (2010) Regulation of leukocyte recruitment by the long pentraxin PTX3. *Nat Immunol* 11: 328–334.
- Cohen PL, Eisenberg RA (1991) Lpr and gld: single gene models of systemic autoimmunity and lymphoproliferative disease. *Annu Rev Immunol* 9: 243–269.
- Munoz LE, Lauber K, Schiller M, Manfredi AA, Herrmann M (2010) The role of defective clearance of apoptotic cells in systemic autoimmunity. *Nat Rev Rheumatol* 6: 280–289.
- Hanayama R, Tanaka M, Miyasaka K, Aozasa K, Koike M, et al. (2004) Autoimmune disease and impaired uptake of apoptotic cells in MFG-E8-deficient mice. *Science* 304: 1147–1150.
- Napirei M, Karsunky H, Zevnik B, Stephan H, Mannherz HG, et al. (2000) Features of systemic lupus erythematosus in Dnase1-deficient mice. *Nat Genet* 25: 177–181.
- Rahman A, Isenberg DA (2008) Systemic lupus erythematosus. *N Engl J Med* 358: 929–939.
- Pego-Reigosa JM, Medeiros DA, Isenberg DA (2009) Respiratory manifestations of systemic lupus erythematosus: old and new concepts. *Best Pract Res Clin Rheumatol* 23: 469–480.
- Sunderrajan EV, McKenzie WN, Lieske TR, Kavanaugh JL, Braun SR, et al. (1986) Pulmonary inflammation in autoimmune MRL/Mp-lpr/lpr mice. Histopathology and bronchoalveolar lavage evaluation. *Am J Pathol* 124: 353–362.
- Lech M, Kulkarni OP, Pfeiffer S, Savarese E, Krug A, et al. (2008) Tlr8/Sigirr prevents murine lupus by suppressing the immunostimulatory effects of lupus autoantigens. *J Exp Med* 205: 1879–1888.
- Kulkarni O, Anders HJ (2008) Chemokines in lupus nephritis. *Front Biosci* 13: 3312–3320.
- Shiozawa F, Kasama T, Yajima N, Odoi T, Isozaki T, et al. (2004) Enhanced expression of interferon-inducible protein 10 associated with Th1 profiles of chemokine receptor in autoimmune pulmonary inflammation of MRL/lpr mice. *Arthritis Res Ther* 6: R78–R86.
- Harari OA, Marshall D, McHale JF, Ahmed S, Haskard DO (2001) Limited endothelial E- and P-selectin expression in MRL/lpr lupus-prone mice. *Rheumatology (Oxford)* 40: 889–895.
- Lloyd CM, Gonzalo JA, Salant DJ, Just J, Gutierrez-Ramos JC (1997) Intercellular adhesion molecule-1 deficiency prolongs survival and protects against the development of pulmonary inflammation during murine lupus. *J Clin Invest* 100: 963–971.
- Deguchi Y, Kishimoto S (1991) Tumour necrosis factor/cachectin plays a key role in autoimmune pulmonary inflammation in lupus-prone mice. *Clin Exp Immunol* 85: 392–395.
- Kim N, Ussin L, Cheng X, Murali R, Sullivan KE (2002) TNFalpha inhibition in MRL/lpr mice ameliorates pulmonary but not renal disease. *J Autoimmun* 19: 215–222.
- Kulkarni OP, Sayyed SG, Kantner C, Ryu M, Schnurr M, et al. (2010) 4SC-101, a novel small molecule dihydroorotate dehydrogenase inhibitor, suppresses systemic lupus erythematosus in MRL-(Fas)lpr mice. *Am J Pathol* 176: 2840–2847.
- Kulkarni O, Eulberg D, Selve N, Zollner S, Allam R, et al. (2009) Anti-Ccl2 Spiegelmer permits 75% dose reduction of cyclophosphamide to control diffuse

- proliferative lupus nephritis and pneumonitis in MRL-Fas(lpr) mice. *J Pharmacol Exp Ther* 328: 371–377.
44. Kakkanaiah VN, Pyle RH, Nagarkatti M, Nagarkatti PS (1990) Evidence for major alterations in the thymocyte subpopulations in murine models of autoimmune diseases. *J Autoimmun* 3: 271–288.
  45. Gutierrez-Ramos JC, Andreu JL, Revilla Y, Vinuela E, Martinez C (1990) Recovery from autoimmunity of MRL/lpr mice after infection with an interleukin-2/vaccinia recombinant virus. *Nature* 346: 271–274.
  46. Singer PA, Theofilopoulos AN (1990) Novel origin of lpr and gld cells and possible implications in autoimmunity. *J Autoimmun* 3: 123–135.
  47. Garlanda C, Hirsch E, Bozza S, Salustri A, De Acetis M, et al. (2002) Non-redundant role of the long pentraxin PTX3 in anti-fungal innate immune response. *Nature* 420: 182–186.
  48. Kulkarni O, Pawar RD, Purschke W, Eulberg D, Selve N, et al. (2007) Spiegelmer inhibition of CCL2/MCP-1 ameliorates lupus nephritis in MRL-Fas/lpr mice. *J Am Soc Nephrol* 18: 2350–2358.
  49. Allam R, Pawar RD, Kulkarni OP, Hornung V, Hartmann G, et al. (2008) Viral 5'-triphosphate RNA and non-CpG DNA aggravate autoimmunity and lupus nephritis via distinct TLR-independent immune responses. *Eur J Immunol* 38: 3487–3498.
  50. Patole PS, Pawar RD, Lichtnekert J, Lech M, Kulkarni OP, et al. (2007) Coactivation of Toll-like receptor-3 and -7 in immune complex glomerulonephritis. *J Autoimmun* 29: 52–59.
  51. Lech M, Skuginna V, Kulkarni OP, Gong J, Wei T, et al. (2010) Lack of SIGIRR/TIR8 aggravates hydrocarbon oil-induced lupus nephritis. *J Pathol* 220: 596–607.
  52. Lech M, Avila-Ferrufino A, Skuginna V, Susanti HE, Anders HJ (2010) Quantitative expression of RIG-like helicase, NOD-like receptor and inflammation-related mRNAs in humans and mice. *Int Immunol* 22: 717–728.
  53. Bustin SA, Benes V, Garson JA, Hellemans J, Huggett J, et al. (2009) The MIQE guidelines: minimum information for publication of quantitative real-time PCR experiments. *Clin Chem* 55: 611–622.