

Zitteliana

An International Journal
of Palaeontology and Geobiology

Series A / Reihe A
Mitteilungen der Bayerischen Staatssammlung
für Paläontologie und Geologie

53



München 2013



Pliensbachian nannofossils from Kachchh: Implications on the earliest Jurassic transgressive event on the western Indian margin

Jyotsana Rai¹ & Sreepat Jain^{2*}

- Zitteliana A 53, 105 – 120
- München, 31.12.2013
- Manuscript received 05.04.2013; revision accepted 16.11.2013
- ISSN 1612 - 412X

¹Birbal Sahni Institute of Palaeobotany, 53 University Road, 226007, Lucknow, India

²DG-2, Flat no. 52D, SFS Flats, Vikas Puri, New Delhi, 110018, India

*Author for correspondence and reprint requests: E-mail: sreepatjain@gmail.com

Abstract

The oldest rocks within the Kachchh Basin belong to the sediments of Kaladongar Formation exposed in Kuar Bet, Pachchham Island (western India). The Formation's lowest unit, the Dingi Hill Member has yielded a moderately diversified calcareous nannofossil assemblage that includes the marker species of *Lotharingius contractus* and *Triscutum sullivanii* of late Early Aalenian age associated with reworked species of *Biscutum finchii*, *Bussonius prinsii*, *Crucirhabdus primulus*, *Crepidolithus pliensbachensis*, *Discorhabdus criotus* and *D. striatus* suggesting an age spanning NJ4a to NJ7 Zones (Early Pliensbachian, Tethyan ammonite Jamesoni Zone to Middle Toarcian, Variabilis Zone). Additionally, samples from four other Kachchh domal localities (Kachchh Mainland: Jara, Jumara and Habo and the Island belt, Waagad) have also yielded reworked Pliensbachian-Toarcian age (~183 Ma) nannofauna viz. *Crepidolithus granulatus*, *Diductius constans*, *Mazaganella protensa*, *Mitrolithus elegans*, *Parhabdolithus liasicus*, *Similiscutum orbiculus*, and *Triscutum tiziense*. This nannofauna age is much earlier than the ammonite-based Earliest Bajocian date (~171.6 Ma) based on the presence of ammonite *Calliphylloceras heterophyloides* (Oppel). Additional reworked assemblages have been studied from the Callovian sediments at Jara, the Bathonian-Callovian sediments at Jumara, the Callovian sediments at Habo and the Oxfordian-Kimmeridgian sediments at Wagad. The present data, thus, indicates the presence of a marine connection at least since the Pliensbachian in Kachchh. It is proposed that both global eustatic rise and local tectonics were responsible for this Pliensbachian inundation of the Kachchh Basin. Similar record of Late Pliensbachian age reworked nannofossils have also come from the Masirah Island of the Sultanate of Oman and of ?Aalenian-Bajocian age nannofossils from Kuwait, lending credence to our findings and of the presence of at least Pliensbachian-Toarcian age sediments in Kachchh.

Key words: Calcareous nannofossils, Pliensbachian, Toarcian, Aalenian, Kuar Bet, Pachchham Island, Kachchh, western India


Zusammenfassung

Die ältesten Sedimente des Kachchh Beckens gehören zur Kaladongar Formation und sind in Kuar Bet, Pachchham (West-Indien), aufgeschlossen. Aus der untersten Einheit dieser Formation (Dingi Hill Member) konnte eine relativ diverse Vergesellschaftung kalkiger Nannofossilien gewonnen werden, die u.A. die Marker-Taxa *Lotharingius contractus* und *Triscutum sullivanii* (oberes unteres Aalenium) enthält sowie umgearbeitete Exemplare von *Biscutum finchii*, *Bussonius prinsii*, *Crucirhabdus primulus*, *Crepidolithus pliensbachensis*, *Discorhabdus criotus* und *D. striatus*, welche eine Zuordnung der Sedimente zwischen NJ4a und NJ7 indizieren (unteres Pliensbachium, Tethys Ammoniten Jamesoni Zone bis mittleres Toarcium, Variabilis Zone). Darüber hinaus wurden in vier anderen Lokalitäten des Kachchh Beckens (Kachchh Festland: Jara, Jumara und Habo sowie Inselgürtel, Waagad) ebenfalls umgearbeitete Nannofossilien aus dem Pliensbachium-Toarcium (~183 Ma) gefunden, nämlich *Crepidolithus granulatus*, *Diductius constans*, *Mazaganella protensa*, *Mitrolithus elegans*, *Parhabdolithus liasicus*, *Similiscutum orbiculus*, und *Triscutum tiziense*. Die Nannofossilien indizieren ein höheres Alter der Sedimente, als die Datierung mit Ammoniten, welche auf der Basis des Vorkommens von *Calliphylloceras heterophyloides* (Oppel) die Sedimente in das unterste Bajocium (~171.6 Ma) stellt. Um das ungewöhnliche Vorkommen von Nannofauna aus dem Pliensbachium-Toarcium in Kachchh zu überprüfen, wurden fünf weitere Lokalitäten erfolgreich auf Nannofossilien hin untersucht. Diese Proben stammen aus Sedimenten des Calloviums von Jara und des Bathoniums-Calloviums von Jumara, aus dem Callovium von Habo sowie aus dem Oxfordium-Kimmeridgium von Wagad. Unsere Ergebnisse lassen den Schluss zu, dass in Kachchh mindestens seit dem Pliensbachium eine marine Verbindung existierte. Es ist wahrscheinlich, dass sowohl der weltweite eustatische Anstieg des Meeresspiegels als auch lokale tektonische Aktivitäten für die Überflutung des Kachchh Beckens im Pliensbachium verantwortlich waren. Ähnliche Vorkommen von umgearbeiteten Nannofossilien aus jungen Pliensbachium sind auch von der Insel Masirah, Sultanat Oman bekannt. Nannofossilien aus dem ?Aalenium-Bajocium kennt man aus Kuwait. Beide Vorkommen stützen unsere Ergebnisse und bestätigen das Vorkommen von


1. Introduction

Calcareous nannofossils from the oldest marine sediments exposed in Kuar Bet area of Pachchham Island, Kachchh Basin, are studied to decipher the earliest epeiric transgressive event in western India in response to the opening of the Ethiopian gulf. The Kachchh Basin (Textfig. 1) is a small sedimentary basin situated on the eastern fringe of the southern extension of the Neotethys at a palaeo-latitude of around 33° S during Early-Middle Jurassic times (Dercourt et al. 2000). Current reports indicate that after a phase of terrestrial sedimentation in Late Triassic (Koshal 1984), marine sedimentation started during the Early Jurassic (Earliest Bajocian) in response to the opening of the Arabian Sea. This date is based on the earliest ammonite record of *Calliphylloceras heterophylloides* from the ?Earliest Bajocian (~171.6 Ma; Pandey et al. 2013), recorded from the Dingy Hill Member (Pachchham Island; Textfig. 2).

We report here a moderately diversified late Early Aalenian age calcareous nannofossil assemblage with reworked Early Jurassic Pliensbachian-Toarcian interval nannotauna from the middle part of the Dingy Hill Member of Kaladongar Formation exposed at Point 16 hillock at Kuar Bet in Pachchham Island (Textfig. 1). This assemblage includes the marker species *Lotharingius contractus* and *Triscutum sullivanii* of late Early Aalenian age which co-occurs with reworked nannotauna of *Biscutum finchii*, *Bussonius prinsii*, *Crepidolithus granulatus*, *C. pliensbachensis*, *Discorhabdus criotus*, *D. striatus* and *Mitrolithus elegans* (Pl. 3) of Pliensbachian-Toarcian interval. Rai (2007) and Rai & Jain (2012) have also previously reported Pliensbachian-Toarcian (Early Jurassic) interval reworked nannofossils from the Pachchham Island of Kachchh.



Textfigure 1: Kachchh Locality map showing the sampling localities at Pachchham, Jara, Jumara, Habo and Wagad. Inset is the detailed map of Kuar Bet showing Point 16, the sampling locality of the oldest beds in Kachchh.



Textfigure 2: Stratigraphy of the Jurassic sediments of Kachchh and the presence of reworked nannotaxa recorded from Jurassic localities of Pachchham (Kuar Bet), Jara, Jumara, Habo and Wagad.

2. Geological setting

The Mesozoic rocks exposed in Kachchh Basin range in age from Aalenian with no datable elements (Middle Jurassic; 178 Ma) to Albian (Early Cretaceous; 98.9 Ma) (Biswas 1991; Rai 2006). These Mesozoic rocks are exposed in six localities viz. Kachchh Mainland, Wagad, Pachchham, Khadir, Bela “Islands” and Chorad Hill (Textfig. 1) and are separated by vast spans of arid Banni Plains or grassland. The western-most island belt of Pachchham comprises two main island parts: the Pachchham and the Kuar Bet (Textfig. 1). The geological sequence exposed in

the Pachchham Island comprises the shallow, lower Kaladongar Formation and the overlying relatively deeper Goradongar Formation (Textfig. 2).

The Kaladongar Formation is a ~472 m thick sequence of conglomerates, sandstones and shales exposed at Kaladongar (Textfig. 1) constituting the oldest stratigraphic unit in Kachchh (Textfig. 2) assigned to ?Aalenian-Early Bajocian (Biswas 1991). On the basis of its lithological attributes, the formation is divisible into three informal Members: upper Babia Cliff Sandstone, middle Kaladongar Sandstone and lowest Dingi Hill (from where the nannofossil record described in this study comes). On the other hand,

the ~152 m thick Goradongar Formation is exposed on the southern flank of Modar Hill at the eastern-most point of the Goradongar Range. Its lower part is predominantly shales with thin flaggy fossiliferous calcareous bands and the upper part is dominated by sandstones with minor shales.

The nannofossil-yielding middle part of the Dingi Hill Member of the Kaladongar Formation (Fürsich et al. 2001) exposed at Point 16 hillock at Kuar Bet in the Pachchham Island (Textfig. 1), ~10 m high, contains broadly alternating sandy fossiliferous limestones and calcareous sandstones with planar and concurrent stratification and flaser bedding and well preserved ichnotaxa viz. *Thalassinoides*, *Rhizocorallium*, and *Diplocriterion* (Textfig. 3). Near the top of this unit is a shell-hash deposit with small sized bivalves and rare gastropods. The base is marked by a conglomerate. The calcareous sandstone level shows prominent bedding fissility and one such level represented by sample number PAT-2 (GPS location: 23° 59' 40" N: 69° 42' 28" E; Textfig. 3) has yielded datable calcareous nannofossils of moderate diversity (Pls 1, 2).

3. Age controversy: Kuar Bet in Pachchham Island

The Kuar Bet beds of Kaladongar Formation contain alternating sandy limestones, conglomerates, red-green sandstones, and occasional shales; the upper part has yellow massive sandstones with calcareous bands (Textfig. 2). For these Kuar Bet beds palaeontological evidences favour a Middle Jurassic age (Wynne 1872; Waagen 1875; Rajnath 1932, 1942; Pandey & Dave 1993; Satyanarayana et al. 1999), whereas palaeobotanical and palynological evidence suggests an Early Cretaceous age (Mathur 1972). Biswas (1977) suggested a Bathonian age but later revised the same down to Bajocian and/or Lias (Biswas 2002). Both Wynne (1872) and Waagen (1875) considered these units as being Bathonian in age, whereas Rajnath (1932, 1942) gave an age range from Bathonian to Early Callovian. Singh et al. (1982) used the marker ammonite *Leptosphinctes* from the overlying Goradongar Formation to date the stata as latest Bajocian. Mathur (1972), who studied plant megafossils, reported the occurrence of the Early Cretaceous marker taxon *Onychiopsis cf. psilotoides*. However, a re-analysis (Jana & Hilton 2007) of the same sediments yielded a broad Middle Jurassic age. Satyanarayana et al. (1999) recorded fossil vertebrae and limb elements of a dinosaur along with petrified tree trunks and suggested an Aalenian-Bathonian age.

The nannofossils assemblage described in this paper, containing *Lotharingius contractus* and *Triscutum sullivanii* from Kuar Bet, now permits a more precise dating of the sediments as late Early Aalenian (Pls 1, 2). The assemblage also contains reworked Pliensba-

chian-Toarcian interval nannotaxa, including *Axopodorhabdus cylindratus*, *Biscutum finchii*, *Bussonius prinsii*, *Crucirhabdus primulus*, *Crepidolithus pliensbachensis*, *Discorhabdus criotus*, *D. striatus*, *Lotharingius contractus*, and *Triscutum sullivanii* (Pl. 3).

Four additional localities were studied to ascertain the anomalous occurrence of Pliensbachian-Toarcian nannotaxa in Kachchh, and all have produced essentially the same results (Pl. 4), strongly suggesting that the timing of the oldest epeiric transgressive event in the Kachchh Basin (western India) occurred at least during the Pliensbachian. These other assemblages of reworked nannofossils come from Jara (in Callovian sediments; Rai 2006; Textfig. 1), Jumara (in Bathonian-Callovian sediments), Habo (in Callovian sediments) and Wagad (in Oxfordian-Kimmeridgian sediments) (Pl. 4, Fig. 2).


4. Material and methods

Permanent duplicate slides, one containing comparatively coarser and the other having finer fraction of the samples for nannofossil productivity and study were prepared using the usual preparation technique described in Bown & Young (1998). These slides were examined under Leitz polarizing microscope (LM) with 10x (magnification) or 12.5x oculars and microphotographs were taken under 100x oil immersion objective. The recovered nannofossils were compared with global marker charts (Bown 1998; Mattioli & Erba 1999; Sandoval et al. 2012).

5. New nannofossil record

The nannotaxa recovered from the sediments studied include the following species (Pls 1, 2): *Axopodorhabdus cylindratus* (Noël, 1965) Wind & Wise in Wise & Wind, 1977, *Biscutum finchii* Crux, 1984, *B. novum* (Goy in Goy et al., 1979) Bown, 1987, *Biscutum* sp., *Bussonius prinsii* (Noël, 1973) Goy, 1979, *Crepidolithus crassus* (Deflandre in Deflandre & Fert, 1954) Noël, 1965, *C. granulatus* Bown, 1987, *C. pliensbachensis* Crux, 1985, *Crucirhabdus primulus* Rood et al., 1973, *Diazmatolithus lehmanii* Noël, 1965, *Discorhabdus* cf. *D. criotus* Bown, 1987, *Discorhabdus striatus* Moshkovitz & Ehrlich, 1976, *Ethmorhabdus gallicus* Noël, 1965, *Lotharingius contractus* Bown & Cooper, 1989, *Micula staurophora* (Gardet, 1955) Stradner, 1963, *Mitrolithus elegans* Deflandre in Deflandre & Fert, 1954, *Octopodorhabdus* sp., *Parhabdolithus liasicus* Deflandre in Grassé, 1952, *Schizosphaerella* sp., *Triscutum sullivanii* de Kaenel & Bergen, 1993, *Tubirhabdus patulus* Rood et al., 1973, *Watznaueria barnesae* (Black in Black & Barnes, 1959) Perch-Nielsen, 1968 and *W. fossacincta* (Black, 1971) Bown in Bown & Cooper, 1998.

The Pliensbachian-Aalenian age reworked nannofossils (Pl. 4) have been also found in the Bathonian-



Textfigure 3: Sampling location from the middle part of the Dingi Hill Member of the Kaladongar Formation exposed at the Point 16 hillock ($23^{\circ} 59' 40''$ N; $69^{\circ} 42' 28''$ E) at Kuar Bet in the Pachchham Island.

Callovian sediments of the three Mainland domes (i.e. Jara, Jumara and Habo), as well as in the Oxfordian-Kimmeridgian sediments of Wagad. All strongly suggest that the timing of the oldest epeiric transgressive event in the Kachchh Basin (western India) occurred at least during the Pliensbachian.

6. Ammonite data

Ammonites in general provide precise dates. However, the Early Jurassic basal sediments of the Kachchh Basin lack ammonite fossils. Hence, to

better appreciate this early duration, it is imperative to look beyond Kachchh, and into the Indian subcontinental region and neighboring ammonite yielding localities such as the Himalayas (Nepal, Tibet and Spiti), Pakistan and Madagascar to assign a comparative age for the basal sediments of Kachchh (Tab. 1). The Pliensbachian and Aalenian are devoid of ammonites from the Indian subcontinent, whereas early Toarcian sediments are quite rich in ammonites. Higher sea level (Textfig. 4) might explain this rich record that not only provided newer niches but also made the previously unavailable land areas available to aquatic animals. However, the available




Plate 1: Calcareous nannofossil assemblage identified from sample number PAT-2, middle part of the Dingi Hill Member of Kaladongar Formation exposed at Point 16 hillock ($23^{\circ} 59' 40''$ N: $69^{\circ} 42' 28''$ E) at Kuar Bet in the Pachchham Island; all forms magnified at 1200x.




Plate 2: Calcareous nannofossil assemblage identified from sample number PAT-2, middle part of the Dingi Hill Member of Kaladongar Formation exposed at Point 16 hillock ($23^{\circ} 59' 40''$ N: $69^{\circ} 42' 28''$ E) at Kuar Bet in the Pachchham Island; all forms magnified at 1200x.




Plate 3: Marker calcareous nannofossil species identified from sample number PAT-2, middle part of the Dingi Hill Member of Kaladongar Formation exposed at Point 16 hillock ($23^{\circ} 59' 40''$ N: $69^{\circ} 42' 28''$ E) at Kuar Bet in the Pachchham Island; all forms magnified at 1200x.

ammonite data do not help in dating the basal sediments of Kachchh, but it is intriguing to ponder as to why Pliensbachian ammonites are not recorded from Kachchh, whereas other body fossils (pelecypods, gastropods and corals) and microfossils (nannofossils) are profuse. Closer and careful sampling might yield Pliensbachian and Aalenian ammonites from Kachchh.

7. Discussion

7.1 Biostratigraphy

There is little consensus amongst Early Jurassic nannofossil workers concerning the ranges of marker species (de Kaenel & Bergen 1993; de Kaenel et al. 1996; Mattioli & Erba 1999; Perilli & Duarte 2006; Veiga de Oliveira et al. 2007). Hence, for now, we

have followed the most commonly used stratigraphic ranges put forward by Mattioli & Erba (1999; see Tab. 2).

The presence of *Lotharingius contractus* Bown & Cooper (FAD at base of NJ8b) and *Triscutum sullivanii* (FAD NJ8b) suggests that the lowest sediments of Kuar Bet Island are late Early Aalenian in age (Tab. 2). The assemblage of *B. prinsii* (FAD NJ4 and LAD NJ8b), *Biscutum finchii* (FAD at NJ4b and LAD at NJ6), *Crepidolithus granulatus* (FAD NJ4a and LAD NJ4b), *C. pliensbachensis* (LAD base of NJ4b), *Crucirhabdus primulus* (LAD NJ5a), *Discorhabdus criotus* (FAD base of NJ7), *D. striatus* (FAD straddling the NJ6 and NJ7), and *Mitrolithus elegans* (FAD NJ2a and LAD Upper Pliensbachian) can be placed between the NJ4a and NJ7 Zones of Early Pliensbachian to Middle Toarcian age, and are here considered to be reworked into the Aalenian (Pls 1–3). This assemblage equates with the Tethyan ammonite in-




Plate 4: Pliensbachian-Toarcian age reworked nannofossils from Wagad (1–4), Jumara (5–11), Habo (12) and Jara (13–14) domes (Kachchh).

Table 1: Ammonite records from the Indian subcontinental region and neighboring areas such as Ladakh Himalaya, Nepal, Pakistan and Madagascar.


Age	Ammonite species	Locality	Reference
Bajocian	<i>Calliphylloceras heterophylloides</i>	Sadhara Dome, Goradongar	Pandey et al. (2013)
	<i>Leptosphinctes</i> sp.	Kaladongar, Kachchh	Singh et al. (1982)
	<i>Dorstensis, Emileia, Sonninia</i> and <i>Witchellia</i>	Central Nepal	Bordet et al. (1971); Gradstein et al. (1989)
	<i>Stephanoceras, Witcheillia</i> and <i>Fontanessia</i>	South Tibet	Hayden (1907); Westermann & Wang (1988)
Aelian is devoid of ammonites from the Indian subcontinent			
Early Toarcian	<i>Alocolytoceras</i>	Spiti	Stoliczka (1866)
	Associated with <i>Alocolytoceras</i> are <i>Paltarpites</i> , <i>Dactylioceras</i> , <i>Fuciniceras</i> , <i>Peronoceras</i> , <i>Protogrammoceras</i> , <i>Sphenarpites</i> , <i>Harpoceras</i> and <i>Hildoceras</i>	Madagascar and Pakistan	Holland (1909)
	<i>Bouleiceras</i> sp.		Arkell (1956); Fatmi (1972, 1986); Fatmi et al. (1986); Shah (1978)
Pliensbachian is devoid of ammonites from the Indian subcontinent			
Late Sinemurian (Oxynotum Zone)	<i>Sulciferites</i> and <i>Gleviceras</i>	South Tibet	Holland (1909); Shah (1978); Wignall et al. (2006)
Early Sinemurian (Bucklandi Zone)	<i>Arietites</i> , <i>Coroniceras</i> and <i>Oxynoticeras</i>	Baluchistan	
Earliest Hettangian (Planorbis Zone)	<i>Psiloceras</i> gr. <i>planorbis</i>	Ladakh Himalaya (Lama Yuru Formation)	Krishna et al. (1997)

terval between the Early Pliensbachian Jamesoni to the Middle Toarcian Variabilis (Gradstein et al. 2004) zones not yet recorded from India.

All data gathered to date suggest that the age of the present nannofossil assemblage from the Dingi Hill Member of the Kaladongar Formation represents a 9 Ma interval span (~189–180 Ma) and that the earliest marine sediments in Kachchh date back to the Early Pliensbachian NJ4a nannofossil Zone and within the Tethyan ammonite Jamesoni Zone (Textfig. 4).

7.2 Palaeogeography

The recorded assemblage has wider palaeogeographical implications. Based on this new record, it appears that after faulting of the Indian plate in its western margin, the transgressive event within the Kachchh basin took place at least during the Pliensbachian. As a result, this early transgression is ~11 Ma older than the previously proposed Early Bajocian ammonite-based date (Pandey et al. 2013). In this context, the record of coeval Late Pliensbachian nannofossils from the Masirah Island (Sultanate of Oman, Arabia; Von Salis & Immenhauser 1997) and



Textfigure 4: Sea level at the Pliensbachian-Toarcian boundary interval (modified after Dera et al. 2009) and the corresponding Tethyan ammonite Zones.

?Aalenian-Bajocian (NJ8b Zone) age nannofossils from Kuwait (Kadar et al. 2012) further strengthens the present record (Textfig. 5). Additionally, reworked Pliensbachian-Aalenian age nannofossils (*Crepidolithus granulatus*, *Diductius constans*, *Mazaganella protensa*, *Mitrolithus elegans*, *Parhabdolithus liasicus*, *Similiscutum orbiculus* and *Triscutum tiziense*) have also been recovered from Jara (in Callovian sediments; Rai 2006; Textfig. 1), Jumara (in Bathonian-Callovian sediments), Habo (in Callovian sediments) and Wagad (in Oxfordian-Kimmeridgian sediments) (Pl. 4, Fig. 2). These findings attest to the presence of marine conditions in the Kachch Basin at least since the Pliensbachian (see also Rai 2007; Rai & Jain 2012). In this context, it is very interesting to note that from the sub-surface sediments in a well in Banni south of Pachchham (Textfig. 1), Rheatian-Liassic (Latest Triassic-Earliest Jurassic) palynoflora have also been recorded (Koshal 1975).

It is proposed that both global eustatic rise (Textfig. 4) coupled with local tectonics during the Pliens-


bachian-Toarcian boundary interval led to the early introduction of this new nannofossil assemblage marking the presence of marine conditions within the Kachch Basin, for the first time.

Table 2 lists the presence of global marker species recorded from Kachch.

8. Conclusions

The oldest rocks exposed in Kuar Bet (Pachchham Island) are of marine origin and are of late Early Aalenian age containing reworked Pliensbachian-Toarcian age nannotaxa. This nannofossil assemblage is of low diversity with moderate preservation.

Based on the presence of *Lotharingius contractus* (FAD at base of NJ8b) and *Triscutum sullivanii* (FAD NJ8b), the lowest sediments of the Dingi Hill Member (Kaladongar Formation) in Kuar Bet (Pachchham Island) are assigned an early Late Aalenian age. The presence of *Biscutum finchii* (FAD NJ5, LAD NJ6),



Textfigure 5: Palaeogeographical map of Pliensbachian-Toarcian time (~190–185 Ma) and Early-Middle Jurassic nannofossil record localities mentioned in the text (modified after Dera et al. 2010; Sandoval et al. 2012).

Bussonius prinsii (NJ5B), *Crepidolithus granulatus* (FAD NJ4a and LAD NJ4b), *Crucirhabdus primulus* (NJ5B), *Discorhabdus criotus* (FAD NJ7) and *Mitolithus elegans* (FAD NJ2a and LAD Upper Pliensbachian) in this assemblage suggests an age between NJ4a–NJ7 nannofossil Zones, straddling the Pliensbachian-Toarcian boundary interval.

Reworked Pliensbachian-Toarcian age nannofossils have also been found from Bathonian-Oxfordian sediments of Jara, Jumara and Habo domes (Kachchh Mainland) and in the Oxfordian-Kimmeridgian sediments of Wagad.

Based on a data set of five sampling localities covering the entire Kachchh basin, the earliest transgressive event in the Kachchh basin occurred ~11 Ma prior to the previously proposed ammonite-inferred [*Calliphylloceras heterophylloides* (Oppel)] Earliest Bajocian age.

It is proposed that the global eustatic rise coupled with local tectonics were responsible for this Pliensbachian-Toarcian age nannofossil introduction in the Kachchh Basin Record of Pliensbachian-Toarcian

age nannofossils from Masirah Island of Sultanate of Oman and? Aalenian-Bajocian (NJ8b Zone) age nannofossils from Kuwait respectively supports this contention.

Acknowledgements

JR expresses her thanks to Dr. Sunil Bajpai, Director Birbal Sahni Institute of Palaeobotany, Lucknow (India) for providing the work facility. Dr. BG Desai of the School of Petroleum Technology (Gandhinagar) and Mr. Niral Patel of the Gujarat State Petroleum Corporation (GSPC, Ahmedabad) are duly thanked for their help during field work by one of us (JR). Dr. Abha Singh of BSIP (Lucknow) is thanked for preparing some of the illustrations. Dr. Sudeep Kanungo (University of Utah, U.S.A.) is thanked for providing hard-to-find literature. SJ thanks Dr. Emanuela Mattioli (France) and Dr. Mikhail Rogov (Russia) for assistance with literature.

Table 2: Global marker species (in bold). Those not recorded from Kachchh are marked by an asterisk.

Time in Ma	Series	Mattioli and Erba (1999)	de Kaenel et. al. 1996			Bown & Cooper 1998		
		STAGES	Italy/S.France	Portugal (Bergen)	Morocco	Boreal		
159.4		CALLOVIAN	U M L	A. helvetica S. hexum L. velatus S. bigotii bigotii ★ S. speciosum S. speciosum octum T. expansus C. torquatus S. hexum V. stradneri A. raha O. decussatus A. hamsonii H. cuvillieri	L. velatus S. bigotii bigotii S. speciosum octum S. spec. speciosum T. expansus S. hexum V. stradneri A. raha O. decussatus A. hamsonii H. cuvillieri	A. helvetica L. velatus S. bigotii bigotii S. speciosum octum S. spec. speciosum T. expansus S. hexum V. stradneri A. raha O. decussatus A. hamsonii H. cuvillieri	S. bigotii maximum S. hexum S. bigotii S. speciosum ★ P. enigma S. hexum A. helvetica ★ T. shawensis	
164.4		BATHONIAN	U M L	C. wiedmannii ★ ★ LCO Discorhabdus spp. W. barnesae H. magharensis	T. shawensis S. speciosum octum A. helvetica Acme B. striatum C. superbus T. sullivanii W. manivitiae W. aff. manivitiae W. aff. contracta W. communis W. britannica T. patulus ★ LCO Biscutum spp. L. sigillatus ★ L. umbriensis C. margerellii H. magharensis W. contracta R. incompta "Small" Calyculus ★ C. cantaluppii D. criotus T. sullivanii B. depravatus	T. shawensis S. speciosum octum A. helvetica Acme B. striatum C. superbus C. magharensis C. spec. speciosum D. constans T. sullivanii acme C. margerellii T. tiziense acme E. britannica P. enigma ★ D. constans T. sullivanii C. magharensis T. tiziense L. umbriensis C. poulinabroniei P. liasicus ★ C. cavus R. incompta "Small" Calyculus ★ C. cantaluppii D. striatus W. colacicchi ★ D. ignotus L. velatus C. superbus C. poulinabroniei C. cantaluppii ★ E. sigillatus Calyculus spp. L. barozzi ★ L. umbriensis, L. frodoi ★ L. hauffii B. finchii L. primigenius ★ ★ P. robustus C. pleinsbachensis B. novum B. aff. B. dubium S. orbiculus S. cruciculus ★ C. crassus M. lenticularis ★	A. helvetica L. velatus S. bigotii bigotii S. speciosum octum S. spec. speciosum T. expansus S. hexum V. stradneri A. raha O. decussatus A. hamsonii H. cuvillieri	S. hexum A. helvetica ★ T. shawensis H. cuvillieri T. shawensis C. margerellii P. enigma ★
169.2		BAJOCIAN	U	★ LCO Discorhabdus spp. W. barnesae H. magharensis	A. helvetica Acme B. striatum C. superbus T. sullivanii W. manivitiae W. aff. manivitiae W. aff. contracta W. communis W. britannica T. patulus ★ LCO Biscutum spp. L. sigillatus ★ L. umbriensis C. margerellii H. magharensis W. contracta R. incompta "Small" Calyculus ★ C. cantaluppii D. criotus T. sullivanii B. depravatus	T. shawensis S. speciosum octum A. helvetica Acme B. striatum C. superbus C. magharensis C. spec. speciosum D. constans T. sullivanii acme C. margerellii T. tiziense acme E. britannica P. enigma ★ D. constans T. sullivanii C. magharensis T. tiziense L. umbriensis C. poulinabroniei P. liasicus ★ C. cavus R. incompta "Small" Calyculus ★ C. cantaluppii D. striatus W. colacicchi ★ D. ignotus L. velatus C. superbus C. poulinabroniei C. cantaluppii ★ E. sigillatus Calyculus spp. L. barozzi ★ L. umbriensis, L. frodoi ★ L. hauffii B. finchii L. primigenius ★ ★ P. robustus C. pleinsbachensis B. novum B. aff. B. dubium S. orbiculus S. cruciculus ★ C. crassus M. lenticularis ★	★ T. shawensis C. superbus T. sullivanii C. superbus S. speciosum W. britannica P. grassei ★ C. impontus/cavus B. prinsii L. contractus T. tiziense R. incompta A. depravatus B. intermedium ★ acme L. hauffii D. criotus ★ O. hamiltoniae B. criotum B. striatum D. constans C. superbus L. sigillatus ★ ★ C. jansae B. striatum P. grandis C. superbus S. finchii C. primulus L. sigillatus ★ S. cruciculus B. profundum A. atavus D. novus, L. hauffii P. dubia Lotharingius spp. C. granulatus C. pleinsbachensis S. geyeri ★ B. prinsii S. cruciculus P. cavus ★ S. precarium ★ C. granulatus P. robustus ★ O. hamiltoniae ★ C. crassus	C. superbus S. speciosum W. britannica P. grassei ★ C. impontus/cavus B. prinsii L. contractus T. tiziense R. incompta A. depravatus B. intermedium ★ acme L. hauffii D. criotus ★ O. hamiltoniae B. finchii C. superbus C. primulus L. sigillatus C. impontus/cavus ★ L. hauffii B. finchii C. pleinsbachensis ★ P. robustus S. cruciculus ★ M. lenticularis ★ C. crassus ★ P. marthae C. pleinsbachensis M. elegans P. liasicus
176.5		AALENIAN	U M					
180.1		TOARCIAN	U					
189.6		PLEINSBACHIAN	U					
195.3		SINEMURIAN	U					
201.9		HETTANGIAN	U M					
205.7			M	"Small" Crepidolithus P. liasicus C. crassus ? T. patulus C. pleinsbachensis M. jansae M. elegans		★ Species not found in Kachchh Basin	S. punctulata ★ P. triassica	

9. References

- Arkell WJ. 1956. Jurassic Geology of the World. London, Oliver and Boyd, 801 pp.
- Biswas SK. 1977. Mesozoic Rock stratigraphy of Kutch, Gujarat. Quarterly Journal of Geological, Mining and Metallurgical Society of India 49, 1–62.
- Biswas SK. 1991. Stratigraphy and Sedimentary evolution of the Mesozoic Basin of Kutch, West India. In: SK Tondon, CC Pant, SM Casshyap (Eds), Sedimentary Basins of India: Tectonic Context, Gyanodaya Prakashan, 74–103.
- Biswas SK. 2002. Mesozoic Rock Stratigraphy, In: Lecture Notes on DST Sponsored contact programme cum Field Workshop on Structure, Tectonics and Mesozoic Stratigraphy of Kachchh 14–20th January 2002, Department of Geology, Maharaja Sayajirao University of Baroda, Vadodara, 29 pp.
- Black M, Barnes B. 1959. The structure of coccoliths from the English Chalk. Geological Magazine 96, 321–328.
- Black M. 1971. Coccoliths of the Speeton Clay and Sutterby Marl. Proceedings of the Yorkshire Geological Society 38, 381–424.
- Bordet P, Colchen M, Krummenacher D, Le Fort P, Mouterde R, Remy M. 1971. Recherches géologiques dans l'Himalaya du Népal, région de la Thakhola. Paris, Centre de la Recherche Scientifique, 279 pp.
- Bown PR, Cooper MKE. 1989. New calcareous nannofossils from the Jurassic. Journal of Micropalaeontology 8, 91–96.
- Bown PR, Cooper MKE. 1998. Jurassic. In: PR Bown (Ed.), Calcareous Nannofossil Biostratigraphy. Dordrecht, Kluwer Academic Publishers, 34–85.
- Bown PR, Young, JR. 1998. Techniques. In: PR Bown (Ed.), Calcareous Nannofossil Biostratigraphy. Dordrecht, Kluwer Academic Publishers, 16–28.
- Bown PR. 1987. Taxonomy, evolution, and biostratigraphy of Late Triassic-Early Jurassic calcareous nannofossils. The Palaeontological Association, Special Papers in Paleontology 38, 1–118.
- Crux JA. 1984. Biostratigraphy of Early Jurassic calcareous nannofossils from southwest Germany. Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen 169, 160–186.
- Crux JA. 1985. *Crepidolithus pliensbachensis* nomen novum pro *Crepidolithus ocellatus* Crux 1984 non (Bramlette and Sullivan) Noël 1965, INA Newsletter 7, 31 pp.
- de Kaenel E, Bergen JA, Perch-Nielsen K. 1996. Jurassic calcareous nannofossil biostratigraphy of western Europe. Compilation of recent studies and calibration of bioevents. Bulletin of the Geological Society, France 167, 15–28.
- de Kaenel E, Bergen JA. 1993. New Early and Middle Jurassic coccolith taxa and biostratigraphy from the eastern Proto-Atlantic (Morocco, Portugal and DSDP Site 547B). Eclogae Geologicae Helvetiae 86, 861–907.
- Deflandre G, Fert C. 1954. Observations sur les Coccolithophorides actuels et fossiles en microscopie ordinaire et electronique, Ann. Paleontol. 40, 115–176.
- Dera G, Pellenard P, Neige P, Deconinck JF, Puéat E, Dommergues J. 2009. Distribution of clay minerals in Early Jurassic Peritethyan seas: Palaeoclimatic significance inferred from multiproxy comparisons. Palaeogeography, Palaeoclimatology, Palaeoecology 271, 39–51.
- Dercourt J, Gaetani M, Vrielynck B, Barrier E, Biju-Duval B, Brunet MF, Cadet JP, Crasquin S, Sandulescu M (Eds). 2000. Atlas Peri-Tethys of Palaeoenvironmental Maps. Paris, Gauthier Villars, 269 pp.
- Fatmi AN, Hydari I, Anwar M, Mengol J. 1986. Stratigraphy of Zidi Formation (Ferozabad Group) and Parh Group; Khuzdar District Balochistan Pakistan. Records Geological Survey of Pakistan 1, 1–32.
- Fatmi AN. 1972. Stratigraphy of Jurassic and Lower Cretaceous rocks and Jurassic Ammonites from northern areas of West Pakistan. Bulletin British Museum (Natural History), Geology 20, 297–380.
- Fatmi AN. 1986. Lower and Middle Jurassic ammonites from "Windar Group" in Phusi Jhal (Kanrach) and Sand (Windar Nai) sections of Lasbela Baluchistan, Pakistan. Memoirs of the Geological Survey of Pakistan. Palaeontologia Pakistanica 16, 1–17.
- Fürsich FT, Pandey DK, Callomon JH, Jaitly AK, Singh, I.B. 2001. Marker beds in the Jurassic of the Kachchh Basin, Western India: their depositional environment and sequence stratigraphic significance. Journal of the Palaeontological Society of India 46, 173–198.
- Gardet M. 1955. Contribution à l'étude des coecolites des terrains néogenes de l'Algérie. Publications du Service de la Carte Géologique de l'Algérie, Serie 2, Bulletin 5, 477–550.
- Goy G, Noel D, Bussonet G. 1979. Les conditions de sédimentation des schistes-carton (Toarcien Inf.) du Bassin de Paris déduites de l'étude des nannofossiles calcaires et des diagraphies. Documents des laboratoires de géologie de la Faculté des sciences de Lyon 75, 22–57.
- Goy G. 1979. Les « Schistes carton » (Toarcien inférieur) du Bassin de Paris en af fl eurements et en sondages. Étude par diagraphie, pétrographie, nanno fl ore calcaire, conditions de sédimentation. PhD thesis, Université Pierre et Marie Curie, Paris, France 187 pp.
- Gradstein FM, Ogg JG, Smith AG (Eds). 2005. A Geologic Time Scale 2004. Cambridge, Cambridge University Press, 589 pp.
- Gradstein FM, Gibling MR, Jansa LF, Kaminski MA, Ogg JG, Sastri M, Thurow TW, Vonrad U. Westermann GEG. 1989. Mesozoic Stratigraphy of Thakkola, Central Nepal (Report of the Lost Ocean Expedition, 1988). Centre for Marine Geology, Dalhousie University, Halifax, Canada, Special Report no. 1, 114 pp.
- Grassé PP. 1952. Traité de Zoologie. Anatomie, Systématique, Biologie. Tome 1, Fasc.1: Phylogénie, Protozoaires: Généralités. Flagellés. Paris, Masson, xii + 1071 pp.
- Hayden HH. 1907. The Geology of the provinces of Tsang and U in Tibet. Memoir Geological Survey of India 36, 122–201.
- Holland TH. 1909. General report of the Geological Survey of India for the year 1908. Records of the Geological Survey of India 38, 1–70.
- Jain S. 2008. Integrated Jurassic biostratigraphy: a closer look at nannofossil and ammonite evidences from the Indian subcontinent. Current Science 95, 326–331.
- Jana BN, Hilton J. 2007. Resolving the age of the Mesozoic Kuar Bet Beds (Kachchh, Gujarat, India): A reinvestigation of paleobotanical and palynological assemblages. Journal of the Asian Earth Sciences 30, 457–463.
- Kadar AP, Crittenden S, Karam KA. 2013. Middle Jurassic to Early Cretaceous calcareous nannofossils from Onshore North Kuwait: A new record. GeoArabia, 18(2), Late Jurassic-Early Cretaceous Evaporite-Siliciclastic Systems of the Arabian Plate, Abstracts of the EAGE's Fourth Arabian Plate Geology Workshop, 9–12 December, 2012, Abu Dhabi, United Arab Emirates, 216–219.
- Koshal VN. 1975. Palynozonation or Mesozoic sub-surface sediments of Banni Kutch, Gujarat. The Quarterly Journal of the Geological, Mining and Metallurgical Society of India 47, 79–81.
- Koshal VN. 1984. Differentiation of Rhaetic sediments in the sub-surface of Kutch based on palynofossils. Petroleum Asia Journal 7, 102–105.
- Krishna J, Sinha AK, Upadhyay R. 1997. First find of the Hettangian *Psiloceras* (Jurassic Ammonitina) from the Indus-Tsangpo Suture, Ladakh Himalaya: diverse implications. Himalayan Geology 18, 145–151.
- Krishna J. 1987. An overview of the Mesozoic stratigraphy of Kachchh and Jaisalmer basins. Journal of Palaeontological Society of India 32, 136–152.
- Mathur YK. 1972. The plant fossils from the Kuar Bet, Patcham Island, Kutch. Current Science 41, 488–489.
- Mattioli E, Erba E. 1999. Synthesis of calcareous nannofossil events in Tethyan Lower and Middle Jurassic successions. Rivista Italiana di Paleontologia e Stratigrafia 105, 343–376.
- Moshkovitz S, Ehrlich A. 1976. Distribution of Middle and Upper Jurassic calcareous nannofossils in the northeastern Negev, Israel and in Gebel Maghara, northern Sinai. Bulletin of the Geological Survey of Israel 69, 1–47.
- Noël D. 1973. Nannofossiles calcaires de sédiments jurassiques finement laminés. Bulletin du Muséum National d'Histoire Naturelle, 3e Serie 75, 95–156.

- Noël D. 1965. Sur les Coccolithes du Jurassique Européen et d'Afrique du Nord: Essai de Classification Fossiles. Paris, Éditions du Centre National de la Recherche Scientifique, Paris, 211 pp.
- Pandey B, Pathak DB, Krishna. J. 2013. *Calliphylloceras heterophyllumoides* (Oppel, 1856) from the basalmost Jurassic succession of Sadpara Dome, Kachchh, India. Journal of the Palaeontological Society of India 58, 61–65.
- Pandey DK, Dave A. 1993. Studies in Mesozoic Foraminifera and chronostratigraphy of Western Kutch Gujarat. Palaeontographica Indica 1, 1–219.
- Perch-Nielsen K. 1968. Der Feinbau und die Klassifikation der Coccolithen aus dem Maastrichtien von Dänemark. Det Kongelige Danske Videnskabernes Selskab, Biologiske Skrifter 16, 1–96.
- Perch-Nielsen K, Immenhauser A. 1997. Mesozoic calcareous nannofossils from Masirah Island (Sultanate of Oman). Journal of Nannoplankton Research 19, 95–101.
- Perilli N, Duarte LV. 2006. Toarcian nannobiohorizons from Lusitanian Basin (Portugal) and their calibration against ammonite zones. Rivista Italiana di Paeontologia e Stratigrafia 112, 417–434.
- Rai J, Jain S. 2012. Early Jurassic Gondwanaland break up – A Nannofossil Story. In: DST Sponsored Field Workshop and Brainstorming Session on 'Geology of Kachchh Basin, Western India: Present Status and Future Perspectives', 26th to 29th January 2012, Kachchh University, Bhuj, Kachchh (Abstract).
- Rai J. 2006. Reworked Pliensbachian-Aalenian nannofossils from Jara Dome, Kutch: Early Jurassic Palaeobiogeography of western India at The XIth International Nannoplankton Association at Nebraska, U.S.A., 84–85.
- Rai J. 2007. Early Jurassic calcareous nannofossils from Patcham Island, Kutch, western India. XXIth Indian Colloquium on Micro-palaeontology and Stratigraphy, Nov. 16–17th, 2007, B.S.I.P., Lucknow, 143.
- Rajnath, 1932. A contribution to the stratigraphy of Cutch. Quarterly Journal of the Geological Mining and Metallurgical Society of India 4, 162–174.
- Rajnath, 1942. Jurassic rocks of Cutch – their bearing on the stratigraphy of some problem of Indian Geology. Proceedings of the Indian Science Congress 29, 93–106.
- Rood AP, Hay WW, Barnard T. 1973. Electron microscope studies of Lower and Middle Jurassic coccoliths. Eclogae Geologicae Helvetiae 66, 365–382.
- Salis, AK von, Immenhauser A. 1997. Mesozoic calcareous nannofossil from Masirah Island (Sultanat of Oman). Journal of Nannoplankton Research 19, 95–101.
- Sandoval J, Bill M, Aguado R, O'Dogherty L, Rivas P, Morad A, Guex J. 2012. The Toarcian in the Subbetic basin (southern Spain): Bio-events (ammonite and calcareous nannofossils) and carbon-isotope stratigraphy. Palaeogeography, Palaeoceanography, Palaeoclimatology, Palaeoecology 342/343, 40–63.
- Satyanarayana K, Dsagupta DK, Dave A, Das KK. 1999. Record of skeletal remains of dinosaur from early Middle Jurassic of Kuar Bet, Kutch, Gujarat. Current Science 77, 639–641.
- Shah SMI. 1978. Stratigraphy of Pakistan. Memoir of the Geological Survey of Pakistan 5, 1–250.
- Singh CSP, Jaitly AK, Pandey DK. 1982. First report of some Bajocian-Bathonian (Middle Jurassic) ammonoids, and age of the oldest sediments from Kachchh (Gujarat), India. Newsletter on Stratigraphy 11, 37–40.
- Stoliczka F. 1866. Geological sections across the Himalayan Mountains, From Wangtubridge on the river Sutlej to Sungdo on the Indus: with an account on the formations in Spiti, accompanied by a revision of all known fossils from that districts. Memoirs of the Geological Survey of India 5, 1–153.
- Stradner H. 1963. New contributions to Mesozoic stratigraphy by means of nannofossils. Proceedings of the Sixth World Petroleum Congress, Frankfurt am Main, Sect. 1, paper 4, 167–184.
- Veiga de Oliveira LC, Duarte L, Lemos VB, Comas-Rengifo MJ, Perilli N. 2007. Calcareous nannofossil biostratigraphy and correlation with ammonites zones of the Pliensbachian-lowermost Toarcian (Lower Jurassic) of Peniche (Lusitanian Basin, Portugal). In: de IS Carvalho, RCT Cassab, C Schwanke, MA Carvalho, ACS Fernandes, MAC Rodrigues, MSS Carvalho, M Araújo, MEQ Oliveira (Eds), Paleontologia: Cenarios de Vida. Rio de Janeiro, Editora Interciência, 411–420.
- Waagen W. 1875. Abstract of results of the examination of the ammonite fauna of Kutch with remarks on their distribution among the beds and probable age. Records of the Geological Survey of India 4, 89–101.
- Westermann GEG, Wang YG. 1988. Middle Jurassic Ammonites of Tibet and the age of the Lower Spiti Shales. Palaeontology 1, 295–339.
- Wignall PB, Hallam A, Newton RJ, Sha JG, Reeves E, Mattioli E, Crowley S. 2006. An eastern Tethyan (Tibetan) record of the Early Jurassic (Toarcian) mass extinction event. Geobiology 4, 179–190.
- Wise SW, Wind FH. 1977. Mesozoic and Cenozoic calcareous nannofossils recovered by DSDP Leg 36 drilling on the Falkland Plateau, southwest Atlantic sector of the southern ocean. In: PF Barker, IWD Dalziel, MG Dinkelmann, DH Elliott, AM Gombos, A Lonardi, G Plafker, J Tarney, RW Thompson, RC Tjalsma, CC von der Borch, SW Wise (Eds), Deep Sea Drilling Project, Washington, Initial Reports, v. 36, 269–492.
- Wynne AB. 1872. Memoir on the geology of Kutch, to accompany a map compiled by A. B. Wynne and F. Fedden during the seasons 1867–68 and 1868–69. Memoirs of the Geological Survey of India 9, 1–289.

**Appendix: Index of calcareous nannofossils
identified in the present study.**

1. *Axopodorhabdus atavus* (Grün et al., 1974) Bown, 1987
2. *Axopodorhabdus cylindratus* (Noël, 1965) Wind & Wise in Wise & Wind, 1977
3. *Biscutum finchii* Crux, 1984
4. *Biscutum novum* (Goy in Goy et al., 1979) Bown, 1987
5. *Biscutum* sp.
6. *Bussonius prinsii* (Noël, 1973) Goy, 1979
7. *Crepidolithus crassus* (Deflandre in Deflandre & Fert, 1954) Noël, 1965
8. *Crepidolithus granulatus* Bown, 1987
9. *Crepidolithus pliensbachensis* Crux, 1985
10. *Crucirhabdus primulus* Rood et al., 1973
11. *Diazmatolithus lehmanii* Noël, 1965
12. *Discorhabdus cf. D. criotus* Bown, 1987
13. *Discorhabdus striatus* Moshkovitz & Ehrlich, 1976
14. *Ethmorhabdus gallicus* Noël, 1965
15. *Lotharingius contractus* Bown & Cooper, 1989
16. *Micula staurophora* (Gardet, 1955) Stradner, 1963
17. *Mitrolithus elegans* Deflandre in Deflandre & Fert, 1954
18. *Octopodorhabdus* sp.
19. *Parhabdolithus liassicus* Deflandre in Grassé, 1952
20. *Schizosphaerella* sp.
21. *Triscutum sullivanii* de Kaenel & Bergen, 1993
22. *Tubirhabdus patulus* Rood et al., 1973
23. *Watznaueria barnesae* (Black in Black & Barnes, 1959) Perch-Nielsen, 1968
24. *Watznaueria fossacincta* (Black, 1971) Bown in Bown & Cooper, 1998