

Early Life Nutritional Programming of Obesity: Mother-Child Cohort Studies

Michael E. Symonds^a Michelle A. Mendez^b Helle Margrete Meltzer^c
Berthold Koletzko^d Keith Godfrey^e Stewart Forsyth^f Eline M. van der Beek^g

^aEarly Life Nutrition Research Unit, Academic Division of Child Health School of Medicine, Queen's Medical Centre, University Hospital, University of Nottingham, Nottingham, UK; ^bCarolina Population Center, University of North Carolina-Chapel Hill, Chapel Hill, N.C., USA; ^cNorwegian Institute of Public Health (NIPH), Division of Environmental Medicine, Oslo, Norway; ^dDr. von Hauner Children's Hospital, University of Munich Medical Centre, Munich, Germany, ^eMRC Lifecourse Epidemiology Unit, University of Southampton, Southampton General Hospital, Southampton, UK, and ^fDSM – Martek Division, Dundee, UK; ^gDanone Baby Nutrition – Research Centre for Specialised Nutrition, Singapore, Singapore

Key Words

Pregnancy · Obesity · Insulin resistance

Abstract

Background: The obesity epidemic has resulted in more overweight/obese women before and during pregnancy. Their offspring tend to have higher birth weights and more body fat, and carry an increased risk of obesity later in life. These effects may partly be related to the heightened risk of gestational diabetes, occurring in at least 16% of all pregnancies irrespective of current body weight. **Methods:** An ILSI Europe workshop reviewed the key contributors leading to adverse outcomes in pregnancy and childhood, including gestational weight gain and nutrition. New research opportunities from prospective mother-child cohort studies were explored. **Results:** Simple measures of gestational weight gain provide insufficient detail of the underlying physiological and metabolic adaptations occurring in pregnancy, and should be complemented by measures of body composition, metabolic and endocrine responses. Recordings of maternal dietary intake and nutrient status are often limited and potential correlations with gestational weight gain have been poorly studied. Many pregnancies in overweight/

obese women are uncomplicated and result in offspring of normal weight, leaving the main determinants of later adverse outcomes to be clarified. **Conclusions:** The workshop provided insights of primary measurements for the characterization of sustainable nutritional intervention strategies in the mother, infant and child for preventing obesity in later life.

Copyright © 2013 S. Karger AG, Basel

Introduction

The current obesity epidemic has resulted in an increasing number of overweight and obese women entering pregnancy [1]. These mothers are more likely to have infants with increased birth weight, including macrosomia (birth weight >95% percentile) and higher neonatal fat mass at birth, as well as an increased risk of having infants born preterm or small for gestational age [2, 3]. Ultimately, these children carry a greater risk of development of abdominal or general obesity and related metabolic disorders later in life [2]. Despite the growing evidence showing both short and long-term consequences of maternal obesity for child health, there are important gaps in the knowledge about the mechanisms involved.




Fig. 1. Summary of the potential lifetime benefits of early intervention in the mother and/or infant for preventing later adverse outcomes of obesity. Derived from Godfrey et al. [73].

These include the potential contribution of factors such as maternal diet, gestational weight gain and metabolic perturbations during pregnancy, and the extent to which the postnatal diet and feeding practices may modify or determine the long-term health risks.

A workshop organized by the International Life Sciences Institute (ILSI) Europe in October 2011 was designed to discuss how current and future cohort studies can increase our knowledge and understanding of the mechanisms, especially those which are nutrition related, that may be associated with an increased risk of adverse pregnancy and childhood outcomes in obese pregnant women. Although maternal obesity during pregnancy has been linked to other adverse child health outcomes, the scope of the workshop was focused upon risk of obesity in childhood and related metabolic disorders. The aim of the workshop was to explore the potential value of combining cohort data in relation to how maternal obesity and diet influence the early nutritional environment of the fetus and interact with the postnatal environment to determine childhood body composition.

This paper summarizes the findings from the workshop and in particular discusses current knowledge in this area to identify research priorities and/or strategies that could help to identify optimal methods for undertaking observational research in humans to address those research priorities, and considers the key challenges in translating this information to potential intervention strategies. Following this approach we address the opportunities and possible limitations of using existing cohorts to increase our understanding of long-term effects of ma-

ternal nutrition in pregnancy on child obesity risk. Ultimately we aim to provide guidelines for future research that will enable us to establish the long-term benefits for both the mother and child. Ideally this could establish whether the benefits for long-term metabolic health are enhanced the earlier an intervention is introduced to prevent obesity, as summarized in figure 1.

In many developed countries, and especially in Europe, a significant number of studies on pregnancy, birth and child cohorts are ongoing, enabling collaboration and complementary research in a way not feasible just a decade ago. An overview of these can be found at <http://www.birthcohorts.net/>. Combining the results from such studies therefore opens the possibility to determine the potential contributions of diet and maternal prepregnancy weight, pregnancy weight gain, birth weight and postnatal growth to adult body composition, as illustrated by the pooled analysis of data from several EU-based cohorts under the CHICOS initiative [4] as well as the recent initiative to combine data from cohorts in low and medium income countries [5, 6].

How Do Maternal Obesity and Diet Influence the Early Nutritional Environment of the Fetus?

The mechanisms through which maternal obesity may 'program' obesity and related metabolic disorders during pregnancy and after birth are uncertain. An increasing number of publications indicate that maternal obesity can result in chronic inflammation and oxidative stress [7]. These conditions may, for example, impair pla-

centration and establishment of adequate blood supply, perhaps altering fetal metabolism [8]. Fetal changes in growth and body composition may also be related to the heightened risk of insulin resistance early in pregnancy and of the development of gestational diabetes [9], though much remains uncertain about the long-term consequences of these metabolic disturbances [10]. Moreover, while obese women are at greater risk, there is increasing evidence that at least 16% of all women can show signs of metabolic disturbances especially in glucose homeostasis irrespective of body weight [11], as raised glucose alone can increase the risk of offspring obesity [12]. Thus, obesity risk in children linked to maternal nutritional status may occur among women at both extremes, both high [13] and low [14] body mass index (BMI), as well as normal weight women.

Differences in dietary intake and diet quality during pregnancy among overweight or obese women may play an important role in the early development of the fetus and increase child obesity risk. There have been a large number of epidemiological studies in both historical and present day mother-child cohorts as well as animal intervention studies that have explored the impact of maternal diet during pregnancy and lactation on the long-term health of the resulting offspring [15]. However, studying these relationships in more detail in animal studies is often hampered by the fact that more extreme over-nutrition than seen in human populations (e.g. a 4-fold increase in total fat intake compared with controls [16]) as well as under-nutrition are the most used approach. Also, data collection on nutrient intake in epidemiological studies are often limited to a single point in time during pregnancy and lack specificity due to an absence of biomarker measurements, e.g. nutrient status. Thus, the evidence-based pool of knowledge enabling us to provide targeted nutritional advice to women, especially those that are obese, to improve nutrition and health status before, during or after pregnancy, remains limited. Although maternal weight gain is clearly associated with birth size, data collected at week 27 of pregnancy in a large cohort study did not show an association between birth weight and the total intake of either energy or specific macronutrients [17]. It was suggested that the immediate effects of maternal obesity on the offspring may be exacerbated only upon further exposure to an obesogenic diet. This has led to an increased focus on the possible contribution of maternal excessive weight gain throughout pregnancy, though again, links with child obesity remain to be established [18, 19].

Clearly, factors other than diet alone can also be important and will include a range of lifestyle components such as exercise and socio-economic status [16]. Indeed, the implications of maternal obesity, gestational weight gain and diet during pregnancy for a child's long-term risk of obesity remain uncertain and this may be partly related to the extent to which such associations are attributable to shared family environments beyond the womb, such as postnatal diet and activity patterns, which have not always been taken into account [20].

An imbalance between food intake and level of physical activity is frequently reported to be a primary mechanism behind the global rise in obesity although again there are likely to be many modifying factors [21]. It is noted that even a small change in energy balance on a daily basis may, over time, lead to obesity [22], which means our ability to prevent excess weight gain, both in women of reproductive age as well as their offspring, remains a challenging target.

A wide range of studies in (non-pregnant) adults are now starting to identify the optimum diets for both promoting and maintaining weight loss [23], as well as for preventing weight gain. Critical components of these diets include their overall energy content together with the quantity and quality of fat and fiber, in addition to glycemic index and protein and micronutrient content. These factors further interact and add more complexity to the topic as a simple focus on energy intake may not solve the problem. Also, a trend for increased intake of nutritional supplements by pregnant women is seen irrespective of their nutritional status and current body weight, and the long-term consequences are currently unknown [24]. In many women, dietary preferences can change, especially in early pregnancy [25], although whether this contributes to the consumption of a nutritionally imbalanced diet or later adverse effects, especially in obese women, remains to be explored. A clear challenge is to establish whether or not these aspects of maternal diet, at defined stages of gestation, may be correlated not only with maternal metabolic status and body composition, but also with gestational weight gain, and importantly with subsequent body composition development in the offspring. Moreover, the extent to which prenatal effects may interact with the long-term impact of postnatal feeding practices, such as duration of breastfeeding and/or timing and type of weaning diets adopted, is of interest [26].

It is now apparent from studies into pregnancy-induced preeclampsia that both the severity of the disease and the birth outcomes are dependent on whether maternal complications commence in early or late pregnancy

[27, 28]. A comparable situation may also occur with regard to potential adverse effects of maternal obesity as well as gestational diabetes (irrespective of maternal body weight before pregnancy). It will therefore be important to identify those women who may show an early adverse response to pregnancy, such as compromised placental function, and consider the relative contribution of total fat as well as body fat distribution and early weight gain together with measurements of both macro- and micro-nutrient intake [29]. Central as opposed to peripheral adiposity is a primary determinant of the metabolic syndrome [30], so a more detailed assessment of relative fat distribution both before and through pregnancy is necessary. These types of measurements can then be related to body composition outcomes both at birth and through infancy, and are now being examined directly by a European Union Framework project entitled Early Nutrition [31].

An adverse body composition, in terms of relative muscle and fat distribution, and how this may change and/or be nutritionally sensitive in the fetus and offspring remains to be defined. These characteristics also change in women with more extreme BMI, which in terms of dietary guidelines and the validation of biomarkers will mean a range of 'at risk' classifications. For example, prevention of preterm/still birth and/or small for gestational age infants in women with a very high BMI could require separate dietary guidelines and/or interventions. Ultimately we need to provide an evidence-based consensus of the impact of maternal diet and its relationship with metabolism at birth and longer-term outcomes relating to obesity risk. Establishing optimal gestational weight gain in relation to health outcomes in mother and offspring would enable weight control from preconception to postpartum to be integrated into clinical practice. In summary, more research is needed to unravel how maternal obesity, diet during pregnancy and (early) gestational weight gain influence the early nutritional environment of the infant [2].

How Do Maternal Obesity and Diet Interact with the Postnatal Nutritional Environment to Influence Body Composition of the Child?

At present, the specific aspects of maternal diet during pregnancy that may influence later child obesity remain uncertain. Few consistent results have been obtained thus far from trials and observational studies in developed country settings [32]. To date, most animal studies that

have induced maternal obesity tend to compare diets that differ greatly in their macronutrient composition and result in a rapid, rather than gradual, increase in maternal body weight [16]. At the same time most interventions do not produce offspring whose birth weight is increased to enable a direct comparison with the human situation. Also, when subjected to meta-analysis they provide a limited evidence base to suggest targeted interventions in obese women [33]. Nonetheless, some studies do provide interesting findings for further investigation. For example, one recent rodent study suggested that shifting from a high-fat diet to standard chow 1 month before mating may help reverse risk of obesity and metabolic programming [34]. Though observational studies have not confirmed this finding, further research is warranted given a potential link between maternal carbohydrate consumption with epigenetic changes and childhood adiposity [35]. For example, an association between 20% increments in methylation of the retinoid X receptor- α chr9:136355885+ in umbilical cord blood and either total or percent fat mass in the offspring at either 6 or 9 years of age was shown in children of normal body weight [35]. Further insights may be gained from well-designed experimental research exploring more moderate changes in micro- as well as macronutrients [36]. These studies may also help us to understand the relative contribution of imbalances in the macronutrient composition, e.g. a relative increase in fat, protein or carbohydrate in an isocaloric setting.

Brown Adipose Tissue Thermogenesis

It should also be noted that the fat mass of newborn infants is much greater than the majority of other animal species used to study adipose tissue development experimentally [37]. Moreover, adipose tissue in the newborn comprises different depots of both brown and white fat. Whereas the latter represents a storage depot, brown fat contains a unique uncoupling protein that is rapidly activated at birth and is essential in enabling the newborn to effectively adapt to cold exposure of the extra-uterine environment [38]. The activity of brown adipose tissue has now been shown to be greater in healthy children than adults [39] and specific depots of brown fat, including those located around the supraclavicular depots, are retained into adulthood and can have an important role in energy balance [40]. Adult obesity is accompanied by a significant reduction in the amount of brown adipose tissue [41], although the extent to which body composition changes with age, BMI and during gestation has yet to be

quantified despite this now being feasible [39]. It is therefore important that the growth, development and function of different fat depots are examined together with how these can be influenced by both the pre- and postnatal nutritional environment. Furthermore, changes in fat growth could be accompanied by changes in the development of appetite control that is centrally regulated and continues into postnatal life [42], both of which are likely to be influenced by the maternal/infant genotype as well as current diet. To date, these aspects have not been included in any mother-child cohorts. Altogether these arguments emphasize the enormous potential of early intervention and the need to identify key diet- and nutrient-related determinants during pregnancy as the most effective way forward to change the developmental path towards future obesity and metabolic-related problems.

Adipose Tissue Growth, Gender and Diet

In terms of examining the long-term outcomes, adipose tissue is one of the fastest growing organs in early life [43] and simply measuring body weight or total fat mass at a fixed time point in infants and/or children provides very limited insights into later body composition or whether that individual is at greater risk of later obesity. Birth weight and body composition, especially fat mass, are gender dependent and these differences persist through infancy, at least in breast-fed infants [44]. A range of factors determine fetal adiposity, including maternal body composition [45], whilst formula feeding can affect adiposity [46], a response that can be transiently enhanced in offspring of obese mothers [47]. The effect of gender on adiposity may be amplified with maternal obesity, as it has recently been shown that post-term girls are slimmer than term girls at birth whereas in boys, those born post-term show accelerated weight gain and subsequently have a much higher incidence of obesity compared to term boys [48]. It remains to be established whether formula feeding per se can have a long-term influence on either weight and/or fat growth in a gender-dependent manner, although accelerated postnatal growth is an established risk of later obesity.

Ultimately the child's own diet and activity patterns, both important influences on obesity risk, will be largely dependent on the type of diet and lifestyle within the family, as will family decisions regarding the duration of breast feeding and both the timing and type of diet introduced at weaning. Currently, the extent to which breast feeding is protective against excess weight gain in early

life [49] remains to be fully established although feeding from the breast as opposed to a bottle may influence appetite control [50]. Multi-generational studies also suggest transmission of obesity risk across generations is linked to high birth weight [51], but the transmission of dietary habits is also likely to contribute. Numerous dietary factors are likely to be relevant for child obesity risk, starting with breast versus formula feeding habits and protein intakes, but evaluating the role of multiple factors poses a challenge [52]. Lower breast feeding rates as reported among obese women are likely to contribute to obesity risk and perhaps differences in appetite regulation among their children [53]. With respect to the quality of postweaning diets, it is important to note that the extent to which familial dietary environments are shared is not well established and correlation of intakes within parent child dyads is modest [54].

It is clear that a focus on maternal diet during pregnancy should be combined with a greater understanding of the impact of postnatal factors. For example, the substantial change in plasma concentration of placental hormones following the establishment of the placenta can be accompanied by the onset of maternal nausea and decline and/or change in her food preferences [55]. Ultimately this is a developmental process in which responses that occur initially within the mother on becoming pregnant can impact on embryogenesis and placentation. This can determine placental function and nutrient supply to the fetus, thereby influencing fetal growth and weight and body composition at birth [56]. For those infants that are breast fed, maternal diet may not only influence mammary gland development but also milk composition as seen in a porcine model [57]. However, for those infants who are fed entirely on formula milk, it is easier to record their food intake and make a more direct association between dietary composition and current growth and body composition.

As is the case for maternal diet, animal studies to date on postnatal feeding and obesity have not addressed several key issues. This includes the extent to which formula feeding as opposed to breast feeding influences the intake of bioactive compounds [58] and thus later body composition, and will include modifications in formula with time [59]. Investigations indicate that changes in macronutrient intake together with the amount of food consumed impact on fetal growth and development [60], but the extent to which these adaptations persist, or are modified through lactation or beyond, are unknown.

Table 1. Summary of optimal requirements and outputs when undertaking mother-child cohort studies in order to assess the role of maternal and infant nutrition on later obesity

Dietary intake/ pattern	Nutrient status assessment	Body composition	Other measurements	Stage of development	Potential biases/ confounders
<i>Mother</i>					
Specific dietary components versus patterns of food eaten	Glucose tolerance; prefeeding metabolite concentration	Fat distribution; placental size and function	Physical activity; socio-economic status; parity; ethnicity	Preconception; changes with gestation	Gender of fetus
<i>Infant</i>					
Breast versus formula	Hormones and metabolites in cord blood	Birth and growth weight; adipose tissue mass, distri- bution and composition; muscle mass	Gender	From birth up to weaning	
<i>Child</i>					
Weaning; childhood	Glucose tolerance; prefeeding metabolite concentration	Adipose tissue mass, distri- bution and composition; muscle mass	Gender; physical activity	From time of weaning up to puberty	

What Are the Implications and Learning Points for Mother-Child Cohort Studies?

Existing cohorts that have recruited women during pregnancy and begun to follow children to monitor health outcomes provide an opportunity to explore the potential role of specific aspects of the maternal diet, her genotype and social environment on the offspring. In addition to substantial variability in environmental exposures such as pollutants [61], these cohorts span countries with a wide range of food cultures and prevalence of maternal and child obesity [62]. Importantly, within the context of the current obesity epidemic, this enables women of different body mass and nutritional status to be studied in much more detail. It is recognized that dietary assessment is inherently difficult and that the adoption of simplified approaches to characterize diet-based concepts such as ‘a prudent dietary pattern’ may offer distinct advantages [63]. Furthermore, in cohorts studying a range of disease outcomes, combining data from multiple 24-hour recalls either with data from food frequency questionnaires (FFQs) that better reflect episodic foods [19], or with biomarkers which provide objective estimates of intakes [64], has been shown to increase validity and reduce bias in estimates of associations between dietary intakes and health. These approaches are increasingly feasible with the advent of technological advances that reduce the cost and logistical complexity of collecting valid 24-hour recalls via web-based tools among both adults and older children [65].

As these new cohorts also follow offspring into childhood, they provide the opportunity not only to assess health outcomes including obesity, but to evaluate the role played by a child’s postnatal diet and activity. These would have the potential to assess the role of genotype [66] as well as socio-economic and related demographic factors [67].

This introduces the challenge of how to unify the findings made, as each study varies considerably in the breadth and depth of dietary data collected. In addition, most of these data are based on food intake and the extent to which this is translated systemically, perhaps even dependent on maternal health and BMI and/or the pregnancy stage, are not studied.

Examples of the types of data integration that are needed include:

1. Adequate assessment to determine the extent to which data allow an estimate and ranking of intake of specific nutritional factors and related metabolic factors. Dietary assessments will vary in terms of capturing sufficient dietary detail to adequately estimate variability in intakes for the relevant nutrients, foods and food groups, and when precisely data were collected. For example, in FFQs, are different types of fish and seafood assessed in sufficient detail to rank subjects with respect to fatty acid intakes, or types of vegetables disaggregated to allow accurate estimation of folate intake?

2. Measurement error and misreporting may also vary across different protocols, with FFQ tending to

have more error and bias, noting that these issues will be relevant to all methods that rely on self-reported intake. This is particularly problematic in studies on obesity, as misreporting and bias have been found to be strongly linked to weight status [68]. Addressing error and bias may be essential for obtaining valid results. For some dietary factors, biomarkers of intake may be a useful objective tool for calibrating and reducing error and bias in reported intake data. Though dietary recalls and records which also involve self-reporting tend to suffer less from error and bias than FFQs, use of these methods in large-scale studies are currently limited by cost and participant burden. For new studies, including follow-up of children, a range of new technologies and platforms are being developed that would allow more cost-effective assessment of both diet and physical activity via recalls or records, potentially improving data quality and precision [69].

3. Dietary intakes at certain times in pregnancy, including around the time of conception and embryogenesis, may be especially relevant for long-term obesity risk [70]. Although diet is thought to change relatively little during pregnancy, data collected at times closer to the most relevant periods may yield stronger associations with later outcomes [71].

The objectives of undertaking pooled analyses of dietary and related data would aim:

- to provide a more comprehensive review of priority areas that should be addressed in future (and current) studies in order to establish the main mechanisms involved;
- to determine the limitations of individual studies;
- to maximize the research potential of the data through analysis of new pathways;
- to examine the causes of any heterogeneity;
- to assess a wider range of dietary exposure;
- to undertake comparative analyses between different stages of gestation – a key challenge will be to establish the relevant contributions of single compared with multiple dietary factors.

In summary, the majority of mother-child birth cohorts have been studied in isolation with maternal dietary information taken at only one particular stage of gestation. The potential to combine the large number of current studies and include both maternal and infant nutritional data is clearly attractive. It could also lead to the identification of relevant biomarkers from either the mother or her infant to enable early intervention in those women most at risk of producing offspring with disproportionate fat mass. The workshop provided a summary

of primary measurements to enable the characterization of sustainable intervention strategies in the mother, infant and child for preventing obesity in later life (table 1).

This type of strategy could match the advances made from a comparable approach adopted within genome-wide association studies [72]. It would therefore require significant computational input together with the utilization of mathematical modeling.

Ultimately, individual dietary factors, or groups of dietary compounds, could be identified as promoting beneficial child health outcomes such as reduced risk of obesity among women defined as being obese or overweight. This could foster the translation to randomized clinical trials in 'at risk' women either before or during pregnancy. In addition, evidence-based public health campaigns could be introduced across the European Union aimed at raising women's awareness of the vital importance of preventing (the development of) excess fat mass to both their own and their infant's future health.

Acknowledgements

This work was commissioned by the Metabolic Imprinting Task Force of the European branch of the International Life Sciences Institute (ILSI Europe). Industry members of this task force are Abbott Nutrition, Danone, DSM, Mead Johnson Nutrition, Nestlé, Royal FrieslandCampina and Unilever. This publication was coordinated by Fanny Rollin, Scientific Project Manager at ILSI Europe. For further information about ILSI Europe, please email info@ilsieurope.be or call +32 2 771 00 14. The opinions expressed herein and the conclusions of this publication are those of the authors and do not necessarily represent the views of ILSI Europe nor those of its member companies.

The work of Michael E. Symonds, Berthold Koletzko, Keith Godfrey and Eline M. van der Beek in this area is carried out with partial financial support from the Commission of the European Communities, specific RTD Programme 'Quality of Life and Management of Living Resources', within the 7th Framework Programme, contract FP7-289346-EARLY NUTRITION. The manuscript does not necessarily reflect the views of the Commission and in no way anticipates the future policy in this area.

References

- 1 Heslehurst N, Bell R, Rankin J: Tackling maternal obesity: the challenge for public health. *Perspect Public Health* 2011;131:161–162.
- 2 Poston L, Harthoorn LF, Van Der Beek EM: Obesity in pregnancy: implications for the mother and lifelong health of the child – a consensus statement. *Pediatr Res* 2011;69: 175–180.

- 3 Djelantik A, Kunst A, van der Wal M, Smit H, Vrijkotte T: Contribution of overweight and obesity to the occurrence of adverse pregnancy outcomes in a multi-ethnic cohort: population attributive fractions for Amsterdam. *BJOG* 2012;119:283–290.
- 4 Guxens M, Ballester F, Espada M, Fernandez MF, Grimalt JO, Ibarluzea J, Olea N, Rebagliato M, Tardon A, Torrent M, Vioque J, Vrijheid M, Sunyer J: Cohort profile: the INMA – Infancia y Medio Ambiente – (environment and childhood) project. *Int J Epidemiol* 2012;41:930–940.
- 5 Kuzawa CW, Hallal PC, Adair L, Bhargava SK, Fall CH, Lee N, Norris SA, Osmond C, Ramirez-Zea M, Sachdev HS, Stein AD, Victora CG, COHORTS Group: Birth weight, postnatal weight gain, and adult body composition in five low and middle income countries. *Am J Hum Biol* 2012;24:5–13.
- 6 Norris SA, Osmond C, Gigante D, Kuzawa CW, Ramakrishnan L, Lee NR, Ramirez-Zea M, Richter LM, Stein AD, Tandon N, Fall CH, COHORTS Group: Size at birth, weight gain in infancy and childhood, and adult diabetes risk in five low- or middle-income country birth cohorts. *Diabetes Care* 2012;35:72–79.
- 7 Leibowitz KL, Moore RH, Ahima RS, Stunkard AJ, Stallings VA, Berkowitz RI, Chittams JL, Faith MS, Stettler N: Maternal obesity associated with inflammation in their children. *World J Pediatr* 2012;8:76–79.
- 8 King JC: Maternal obesity, metabolism, and pregnancy outcomes. *Annu Rev Nutr* 2006;26:271–291.
- 9 Larsson G, Spjuth J, Ranstam J, Vikbladh I, Saxtrup O, Astedt B: Prognostic significance of birth of large infant for subsequent development of maternal non-insulin-dependent diabetes mellitus: a prospective study over 20–27 years. *Diabetes Care* 1986;9:359–364.
- 10 Kim SY, England JL, Sharma JA, Njoroge T: Gestational diabetes mellitus and risk of childhood overweight and obesity in offspring: a systematic review. *Exp Diabetes Res* 2011;2011:541308.
- 11 Catalano PM, McIntyre HD, Cruickshank JK, McCance DR, Dyer AR, Metzger BE, Lowe LP, Trimble ER, Coustan DR, Hadden DR, Persson B, Hod M, Oats JJ, Group HSCR: The hyperglycemia and adverse pregnancy outcome study: associations of GDM and obesity with pregnancy outcomes. *Diabetes Care* 2012;35:780–786.
- 12 Lawrence JM, Contreras R, Chen W, Sacks DA: Trends in the prevalence of preexisting diabetes and gestational diabetes mellitus among a racially/ethnically diverse population of pregnant women, 1999–2005. *Diabetes Care* 2008;31:899–904.
- 13 Heslehurst N, Simpson H, Ells LJ, Rankin J, Wilkinson J, Lang R, Brown TJ, Summerbell CD: The impact of maternal BMI status on pregnancy outcomes with immediate short-term obstetric resource implications: a meta-analysis. *Obes Rev* 2008;9:635–683.
- 14 Belogolovkin V, Alio AP, Mbah AK, Clayton HB, Wathington D, Salihu HM: Patterns and success of fetal programming among women with low and extremely low pre-pregnancy BMI. *Arch Gynecol Obstet* 2009;280:579–584.
- 15 McMillen IC, Adam CL, Muhlhausler BS: Early origins of obesity: programming the appetite regulatory system. *J Physiol* 2005;565:9–17.
- 16 Symonds ME, Sebert S, Budge H: The obesity epidemic: from the environment to epigenetics – not simply a response to dietary manipulation in a thermoneutral environment. *Front Genet* 2011;2:24.
- 17 Lagiou P, Tamimi RM, Mucci LA, Adami HO, Hsieh CC, Trichopoulos D: Diet during pregnancy in relation to maternal weight gain and birth size. *Eur J Clin Nutr* 2004;58:231–237.
- 18 Fernandez ID, Olson CM, De Ver Dye T: Discordance in the assessment of prepregnancy weight status of adolescents: a comparison between the Centers for Disease Control and Prevention sex- and age-specific body mass index classification and the Institute of Medicine-Based classification used for maternal weight gain guidelines. *J Am Diet Assoc* 2008;108:998–1002.
- 19 Schack-Nielsen L, Michaelsen KF, Gamborg M, Mortensen EL, Sorensen TI: Gestational weight gain in relation to offspring body mass index and obesity from infancy through adulthood. *Int J Obes (Lond)* 2010;34:67–74.
- 20 Branum AM, Parker JD, Keim SA, Schempf AH: Prepregnancy body mass index and gestational weight gain in relation to child body mass index among siblings. *Am J Epidemiol* 2011;174:1159–1165.
- 21 McAllister EJ, Dhurandhar NV, Keith SW, Aronne LJ, Barger J, Baskin M, Benca RM, Biggio J, Boggiano MM, Eisenmann JC, Elobeid M, Fontaine KR, Gluckman P, Hanlon EC, Katzmarzyk P, Pietrobelli A, Redden DT, Ruden DM, Wang C, Waterland RA, Wright SM, Allison DB: Ten putative contributors to the obesity epidemic. *Crit Rev Food Sci Nutr* 2009;49:868–913.
- 22 Levine JA, Lanningham-Foster LM, McCrady SK, Krizan AC, Olson LR, Kane PH, Jensen MD, Clark MM: Interindividual variation in posture allocation: possible role in human obesity. *Science* 2005;307:584–586.
- 23 Simpson SA, Shaw C, McNamara R: What is the most effective way to maintain weight loss in adults? *BMJ* 2011;343:d8042.
- 24 Alwan NA, Greenwood DC, Simpson NA, McArdle HJ, Cade JE: The relationship between dietary supplement use in late pregnancy and birth outcomes: a cohort study in British women. *BJOG* 2010;117:821–829.
- 25 Pepper GV, Craig Roberts S: Rates of nausea and vomiting in pregnancy and dietary characteristics across populations. *Proc Biol Sci* 2006;273:2675–2679.
- 26 Grote V, Theurich M, Koletzko B: Do complementary feeding practices predict the later risk of obesity? *Curr Opin Clin Nutr Metab Care* 2012;15:293–297.
- 27 Rolfo A, Many A, Racano A, Tal R, Tagliarfero A, Ietta F, Wang J, Post M, Caniggia I: Abnormalities in oxygen sensing define early and late onset preeclampsia as distinct pathologies. *PLoS One* 2010;5:e13288.
- 28 van der Merwe JL, Hall DR, Wright C, Schubert P, Grove D: Are early and late preeclampsia distinct subclasses of the disease – what does the placenta reveal? *Hypertens Pregnancy* 2010;29:457–467.
- 29 Blumfield ML, Hure AJ, Macdonald-Wicks L, Smith R, Collins CE: Systematic review and meta-analysis of energy and macronutrient intakes during pregnancy in developed countries. *Nutr Rev* 2012;70:322–336.
- 30 Despres JP, Lemieux I: Abdominal obesity and metabolic syndrome. *Nature* 2006;444:881–887.
- 31 Koletzko B, Brands B, Poston L, Godfrey K, Demmelmaier H, Early Nutrition Project: Early nutrition programming of long-term health. *Proc Nutr Soc* 2012;71:371–378.
- 32 Hauner H, Much D, Vollhardt C, Brunner S, Schmid D, Sedlmeier EM, Heimberg E, Schuster T, Zimmermann A, Schneider KT, Bader BL, Amann-Gassner U: Effect of reducing the n-6:n-3 long-chain PUFA ratio during pregnancy and lactation on infant adipose tissue growth within the first year of life: an open-label randomized controlled trial. *Am J Clin Nutr* 2012;95:383–394.
- 33 Muhlhausler BS, Gibson RA, Makrides M: The effect of maternal omega-3 long-chain polyunsaturated fatty acid (n-3 LCPUFA) supplementation during pregnancy and/or lactation on body fat mass in the offspring: a systematic review of animal studies. *Prostaglandins Leukot Essent Fatty Acids* 2011;85:83–88.
- 34 Zambrano E, Martinez-Samayoa PM, Rodriguez-Gonzalez GL, Nathanielsz PW: Dietary intervention prior to pregnancy reverses metabolic programming in male offspring of obese rats. *J Physiol* 2010;588:1791–1799.
- 35 Godfrey KM, Sheppard A, Gluckman PD, Lillycrop KA, Burdge GC, McLean C, Rodford J, Slater-Jefferies JL, Garratt E, Crozier SR, Emerald BS, Gale CR, Inskip HM, Cooper C, Hanson MA: Epigenetic gene promoter methylation at birth is associated with child's later adiposity. *Diabetes* 2011;60:1528–1534.
- 36 Sen S, Simmons RA: Maternal antioxidant supplementation prevents adiposity in the offspring of western diet-fed rats. *Diabetes* 2010;59:3058–3065.
- 37 Widdowson EM: Chemical composition of newly born animals. *Nature* 1950;116:626–628.
- 38 Cannon B, Nedergaard J: Brown adipose tissue: function and physiological significance. *Physiol Rev* 2004;84:277–359.

- 39 Symonds ME, Henderson K, Elvidge L, Bosman C, Sharkey D, Perkins AC, Budge H: Thermal imaging to assess age-related changes of skin temperature within the supraclavicular region co-locating with brown adipose tissue in healthy children. *J Pediatr* 2012;161:892–898.
- 40 Nedergaard J, Bengtsson T, Cannon B: Unexpected evidence for active brown adipose tissue in adult humans. *Am J Physiol Endocrinol Metab* 2007;293:E444–E452.
- 41 Vijgen GH, Bouvy ND, Teule GJ, Brans B, Schrauwen P, van Marken Lichtenbelt WD: Brown adipose tissue in morbidly obese subjects. *PLoS One* 2011;6:e17247.
- 42 Kirk SL, Samuelsson AM, Argenton M, Dhonye H, Kalamatianos T, Poston L, Taylor PD, Coen CW: Maternal obesity induced by diet in rats permanently influences central processes regulating food intake in offspring. *PLoS One* 2009;4:e5870.
- 43 Symonds ME, Pope M, Sharkey D, Budge H: Adipose tissue and fetal programming. *Diabetologia* 2012;55:1597–606.
- 44 Diouf I, Charles MA, Thiebaugeorges O, Forhan A, Kaminski M, Heude B, Group EM-CCS: Maternal weight change before pregnancy in relation to birthweight and risks of adverse pregnancy outcomes. *Eur J Epidemiol* 2011;26:789–796.
- 45 Hull HR, Thornton JC, Ji Y, Paley C, Rosenn B, Mathews P, Navder K, Yu A, Dorsey K, Gallagher D: Higher infant body fat with excessive gestational weight gain in overweight women. *Am J Obstet Gynecol* 2011;205:211.e1–7.
- 46 Gale C, Logan KM, Santhakumaran S, Parkinson JR, Hyde MJ, Modi N: Effect of breastfeeding compared with formula feeding on infant body composition: a systematic review and meta-analysis. *Am J Clin Nutr* 2012;95:656–669.
- 47 Andres A, Shankar K, Badger TM: Body fat mass of exclusively breastfed infants born to overweight mothers. *J Acad Nutr Diet* 2012;112:991–995.
- 48 Beltrand J, Soboleva TK, Shorten PR, Deraiik JG, Hofman P, Albertsson-Wikland K, Hochberg Z, Cutfield WS: Post-term birth is associated with greater risk of obesity in adolescent males. *J Pediatr* 2012;160:769–773.
- 49 Wiklund P, Xu L, Lyytikäinen A, Saltevo J, Wang Q, Volgyi E, Munukka E, Cheng S, Alen M, Keinanen-Kiukkaanniemi S: Prolonged breast-feeding protects mothers from later-life obesity and related cardio-metabolic disorders. *Public Health Nutr* 2012;15:67–74.
- 50 Disantis KI, Collins BN, Fisher JO, Davey A: Do infants fed directly from the breast have improved appetite regulation and slower growth during early childhood compared with infants fed from a bottle? *Int J Behav Nutr Phys Act* 2011;8:89.
- 51 Cnattingius S, Villamor E, Lagerros YT, Wikstrom AK, Granath F: High birth weight and obesity—a vicious circle across generations. *Int J Obes (Lond)* 2012;36:1320–1324.
- 52 Koletzko B, von Kries R, Closa R, Escribano J, Scaglioni S, Giovannini M, Beyer J, Demmelair H, Gruszfeld D, Dobrzanska A, Sengier A, Langhendries JP, Rolland Cachera MF, Grote V, European Childhood Obesity Trial Study Group: Lower protein in infant formula is associated with lower weight up to age 2 y: a randomized clinical trial. *Am J Clin Nutr* 2009;89:1836–1845.
- 53 Hawkes CP, Hourihane JO, Kenny LC, Irvine AD, Kiely M, Murray DM: Gender- and gestational age-specific body fat percentage at birth. *Pediatrics* 2011;128:e645–e651.
- 54 Wang Y, Beydoun MA, Li J, Liu Y, Moreno LA: Do children and their parents eat a similar diet? Resemblance in child and parental dietary intake: systematic review and meta-analysis. *J Epidemiol Community Health* 2011;65:177–189.
- 55 Coad J, Al-Rasasi B, Morgan J: Nutrient insult in early pregnancy. *Proc Nutr Soc* 2002;61:51–59.
- 56 Thornburg KL, O’Tierney PF, Louey S: Review: the placenta is a programming agent for cardiovascular disease. *Placenta* 2010;31:S54–S59.
- 57 Amusquivar E, Laws J, Clarke L, Herrera E: Fatty acid composition of the maternal diet during the first or the second half of gestation influences the fatty acid composition of sows’ milk and plasma, and plasma of their piglets. *Lipids* 2010;45:409–418.
- 58 Fields DA, Demerath EW: Relationship of insulin, glucose, leptin, IL-6 and TNF-alpha in human breast milk with infant growth and body composition. *Pediatr Obes* 2012;7:304–312.
- 59 Socha P, Grote V, Gruszfeld D, Janas R, Demmelair H, Closa-Monasterolo R, Subias JE, Scaglioni S, Verduci E, Dain E, Langhendries JP, Perrin E, Koletzko B: Milk protein intake, the metabolic-endocrine response, and growth in infancy: data from a randomized clinical trial. *Am J Clin Nutr* 2011;94:S1776–S1784.
- 60 McMillen IC, Robinson JS: Developmental origins of the metabolic syndrome: prediction, plasticity, and programming. *Physiol Rev* 2005;85:571–633.
- 61 Vrijheid M, Casas M, Bergstrom A, Carmichael A, Cordier S, Eggesbo M, Eller E, Fantini MP, Fernandez MF, Fernandez-Somoano A, Gehring U, Grazuleviciene R, Hohmann C, Karvonen AM, Keil T, Kogevinas M, Koppen G, Kramer U, Kuehni CE, Magnus P, Majewska R, Andersen AM, Patelarou E, Petersen MS, Pierik FH, Polanska K, Porta D, Richiardi L, Santos AC, Slama R, Sram RJ, Thijs C, Tischer C, Toft G, Trnovec T, Vandentorren S, Vrijkotte TG, Wilhelm M, Wright J, Nieuwenhuijsen M: European birth cohorts for environmental health research. *Environ Health Perspect* 2012;120:29–37.
- 62 Mendez MA, Kogevinas M: Alternative methods of accounting for underreporting and overreporting when measuring dietary intake-obesity relations. *Am J Clin Nutr* 2011;94:1993–1999.
- 63 Padmanabhan V, Veiga-Lopez A, Abbott DH, Recabarren SE, Herkimer C: Developmental programming: impact of prenatal testosterone excess and postnatal weight gain on insulin sensitivity index and transfer of traits to offspring of overweight females. *Endocrinology* 2010;151:595–605.
- 64 Freedman LS, Schatzkin A, Midthune D, Kipnis V: Dealing with dietary measurement error in nutritional cohort studies. *J Natl Cancer Inst* 2011;103:1086–1092.
- 65 Arab L, Tseng CH, Ang A, Jardack P: Validity of a multipass, web-based, 24 h self-administered recall for assessment of total energy intake in blacks and whites. *Am J Epidemiol* 2011;174:1256–1265.
- 66 Forner F, Kumar C, Luber CA, Fromme T, Klingenspor M, Mann M: Proteome differences between brown and white fat mitochondria reveal specialized metabolic functions. *Cell Metab* 2009;10:324–335.
- 67 Druet C, Stettler N, Sharp S, Simmons RK, Cooper C, Smith GD, Ekelund U, Levy-Marchal C, Jarvelin MR, Kuh D, Ong KK: Prediction of childhood obesity by infancy weight gain: an individual-level meta-analysis. *Paediatr Perinat Epidemiol* 2012;26:19–26.
- 68 Mendez MA, Popkin BM, Buckland G, Schroder H, Amiano P, Barricarte A, Huerta JM, Quiros JR, Sanchez MJ, Gonzalez CA: Alternative methods of accounting for underreporting and overreporting when measuring dietary intake-obesity relations. *Am J Epidemiol* 2011;173:448–458.
- 69 Thompson FE, Subar AF, Loria CM, Reedy JL, Baranowski T: Need for technological innovation in dietary assessment. *J Am Diet Assoc* 2010;110:48–51.
- 70 Painter RC, Roseboom TJ, Bleker OP: Prenatal exposure to the Dutch famine and disease in later life: an overview. *Reprod Toxicol* 2005;20:345–352.
- 71 Phelan S, Hart C, Phipps M, Abrams B, Schaffner A, Adams A, Wing R: Maternal behaviors during pregnancy impact offspring obesity risk. *Exp Diabetes Res* 2011;2011:985139.
- 72 Frayling TM, Timpson NJ, Weedon MN, Zeggini E, Freathy RM, Lindgren CM, Perry JR, Elliott KS, Lango H, Rayner NW, Shields B, Harries LW, Barrett JC, Ellard S, Groves CJ, Knight B, Patch AM, Ness AR, Ebrahim S, Lawlor DA, Ring SM, Ben-Shlomo Y, Jarvelin MR, Sovio U, Bennett AJ, Melzer D, Ferrucci L, Loos RJ, Barroso I, Wareham NJ, Karpe F, Owen KR, Cardon LR, Walker M, Hitman GA, Palmer CN, Doney AS, Morris AD, Smith GD, Hattersley AT, McCarthy MI: A common variant in the FTO gene is associated with body mass index and predisposes to childhood and adult obesity. *Science* 2007;316:889–894.
- 73 Godfrey KM, Gluckman PD, Hanson MA: Developmental origins of metabolic disease: life course and intergenerational perspectives. *Trends Endocrin Met* 2010;21:199–205.