



- Zitteliana B 32, 163 – 174
- München, 31.12.2014
- Manuscript received 14.04.2014; revision accepted 10.10.2014
- ISSN 1612 - 4138

## Intraspecific evolution within an ecologically diverse species can be used to examine evolutionary trends observed in ruminant morphology: a review of case studies of the Japanese sika deer

Mugino Ozaki Kubo

*The University Museum, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan*

*E-mail:* mugino@um.u-tokyo.ac.jp

### Abstract

This review addresses the question “Are interspecific and macroevolutionary trends observed in ruminant morphology found among populations of a single species?” Several case studies of the Japanese sika deer, *Cervus nippon*, are discussed. The Japanese sika deer is a suitable species for this purpose because it inhabits various environments from the northern subarctic forests to the southern subtropical forests. It shows conspicuous variations in feeding habits, body size, and other morphological and behavioral traits. It can be concluded that selective pressures governing interspecific variations among ruminants, e.g., tooth wear due to grazing or mountainous habitat, promote morphological adaptation of local sika deer populations. However, in some cases, genetic differentiation among local populations is relatively small, resulting in a small degree of change compared to that observed between species. Intraspecific comparison of an ecologically diverse species like the Japanese sika deer is useful for elucidating the evolutionary patterns within ruminants and the possible causes for those patterns.

**Key words:** Sika deer, *Cervus nippon*, Cervidae, evolution, morphology, hypsodonty, teeth, limb, body size, populations.


### 1. Introduction

#### 1.1 Comparative study of populations: less investigated issues in ruminant evolutionary research

Macroevolutionary trends in ruminant morphology have been examined in numerous studies. These include trends in the body size (e.g., Jarman 1974), skull and teeth, particularly cheek teeth (e.g., Janis 1995), digestive tract (e.g., Hofmann 1989), and cranial ornamentation (e.g., Davis et al. 2011). Based on the interspecific comparison of diverse extant and fossil species, these studies have not only provided a suite of morphological characteristics that are useful for the reconstruction of the phylogeny of ruminants but also indicated the possible selective pressures that drove morphological differentiation. Because ruminants are highly successful herbivores and show continuous feeding variation from grass-dominant diets to browse-dominant diets, morphological characteristics that are relevant to the feeding type are well investigated (Hofmann 1989; Janis 1995; Pérez-Barbería & Gordon 1999; Williams & Kay 2001; Clauss et al. 2003, 2008; Hofmann et al.

2008; Codron & Clauss 2010; Damuth & Janis 2011; Kaiser et al. 2013). Recent statistical advances in phylogenetical comparative methods (PCMs) and increased reliability of the phylogenetic relationships among Ruminantia have greatly contributed to an improved understanding of the relationship between feeding ecology and morphology.

Fewer studies have investigated morphological differentiation among ruminant populations based on the notion that selective pressures shaping macroevolutionary trends among species also effectively drive microevolution within a species (Foster & Endler 1999). This is because 1) ecological variation within a species is normally smaller than that found among the diverse species of ruminants, and 2) recent studies of the population ecology of ruminants pay close attention to what causes differences in specific characteristics of populations, without placing them in the context of macroevolutionary trends. A number of studies of ruminant populations have identified ecological/environmental factors associated with differences in body mass, fecundity, mortality, or recruitment among populations (Clutton-Brock et al. 1984; Gaillard et al. 1993; Clutton-Brock et al. 1997; Mysterud et al. 2001; Festa-Bian-



**Figure 1:** Geographical distribution of the sika deer in Japan, along with its latitudinal variation in diet. The dotted areas of the left map are sika deer habitats. The solid lines represent the boundary between the two phylogenetic lineages: the northern lineage inhabits the area from Hokkaido to the central Honshu, whereas the southern lineage inhabits the southwestern part of Honshu, Kyushu, and the islets surrounding Kyushu. A possible hybridization area is located in central-western parts of Honshu and Shikoku. The scatter plot on the right shows latitudinal variation in the diet (modified from Takatsuki 2009). The x-axis represents the percentage of graminoids in the diet determined using quantitative dietary analyses of rumen contents or feces.

chet et al. 2003; Martinez-Jauregui et al. 2009), but these studies can not necessarily determine if these ecological/environmental factors are important for macroevolutionary diversification of ruminants.

To this end, we have been interested in the relationship between microevolution (evolution occurring at the species level) and macroevolution (evolution occurring above the species level or higher taxa). There has been a lack of interaction between those studying the paleontology/macroevo-lution and others studying the current population ecology of ruminants. If interspecific variation of traits among ruminant species had been the result of natural selection, then the same selective pressure could have produced similar patterns among populations within a species. In this sense, studies of intraspecific variation can provide empirical evidence of recent evolutionary trends.

## 1.2 The Japanese sika deer as a model for examining macroevolutionary trends

Based on the above mentioned notion, morphological comparisons were made among Japanese sika deer populations (Ozaki et al. 2007, 2010; Kubo et al. 2011, 2013; Kubo & Yamada 2014). These studies

will be introduced in the subsequent section (Section 2), with the primary focus being on the relationship between feeding ecology and dentognathic morphology (Subsections 2.1 and 2.2).

The sika deer (*Cervus nippon*) is widespread throughout eastern and northeastern Asia, from the Ussuri region of Siberia to northern Vietnam, Taiwan, and Japan (Ohtaishi 1986; Geist 1999). Because the distribution of the sika deer in the Japanese archipelago extends from the northern to the southern islands (from 44°N to 26°N, Fig. 1), they inhabit a wide range of environments, from the northern, subarctic coniferous forests to the southern, temperate or even subtropical broad-leaved forests. Such variation in habitat has a strong influence on their ecology, such as feeding habits, seasonal migration, reproduction, and demography (McCullough et al. 2009). Information on the ecology of local populations has been accumulated for management purposes. Skeletal specimens are available for each population from these surveys (collection of animals that have died of natural causes in the field and animals harvested for nuisance culling or for hunting).

Among ecological variations found in sika deer populations, the most notable and informative variation in macroevolutionary research is variation in

**Table 1:** Sika deer populations used in analyses in Subsections 2.1 and 2.2.

Population	Latitude (°N)	Habitat type	Genetic lineage*	Percentage of grass in their diet**	Feeding type***
1. Eastern Hokkaido	43.4	Deciduous broad-leaved and mixed forest	N	46.3	G
2. Mt. Goyo	39.2	Coniferous or deciduous broad-leaved forest	N	52.5	G
3. Kinkazan Island	38.3	Deciduous broad-leaved forest with some open grasslands	N	65.8	G
4. Nikko	36.8	Deciduous broad-leaved forest	N	73.9	G
5. Ashio	36.7	Deciduous broad-leaved forest with some open grasslands	N	90.0	G
6. Okutama	35.9	Deciduous broad-leaved forest	N	32.1	IM
7. Yamanashi	35.8	Deciduous broad-leaved forest	N	42.2	IM
8. Tanzawa	35.4	Deciduous broad-leaved evergreen broad-leaved forest	N	—	IM
9. Boso Peninsula	35.2	Evergreen broad-leaved forest	N	45.2	IM
10. Izu Peninsula	35.0	Deciduous broad-leaved evergreen broad-leaved forests	N	30.5	IM
11. Nara Park	34.4	Open grassland with planted conifers and deciduous trees	N	81.4	G
12. Shimane	35.4	Evergreen broad-leaved forest	S	38.1	IM
13. Yamaguchi	34.2	Evergreen broad-leaved forest	S	6.5	B
14. Tsushima Island	34.1	Evergreen broad-leaved forest	S	3.4	B
15. Fukuoka	33.5	Evergreen broad-leaved forest	S	7.2	B
16. Mt. Shiraga	32.2	Evergreen broad-leaved forest	S	25.6	IM
17. Yakushima Island	30.3	Evergreen broad-leaved forest	S	4.4	B

\*The phylogenetic lineage based on mitochondrial DNA haplotype: N, northern lineage; S, southern lineage.


\*\*Percentage of graminoids in their diet was obtained from Kubo & Yamada (2014).

\*\*\*Categorical classification of the feeding type is based on data from Ozaki et al. (2007).

food habits (reviewed in Takatsuki 2009) (Fig. 1). Sika deer populations inhabiting the subarctic coniferous forests or the cool temperate deciduous broad-leaved forests subsist mostly on graminoids, particularly dwarf bamboo (e.g., *Sasa nipponica*), which is dominant within the understories of these forests. In some ways, dwarf bamboo is broadly equivalent to other grass species with respect to the relative amount of neutral detergent fiber (NDF; cellulose, hemicellulose, and lignin) and silica (2.8%–15.9% per dry matter) (Van Soest 1994; Motomura et al. 2000; Yokoyama et al. 2000). Thus, these northern populations can be categorized as “grazers” inhabiting closed habitats (Hofmann 1985). On the other hand, the sika deer that inhabit the temperate

evergreen broad-leaved forests feed mainly on the leaves of woody plants, and they are classified as either “intermediate feeders” or “browsers.” Most of the browse species have < 1% of silica concentration; for example, Japanese laurel (*Aucuba japonica*), which is a preferred diet for the sika deer in temperate evergreen forests, has 0.58% of silica in the leaves (Ma & Takahashi 2002). Previous studies of the food habits of the sika deer have provided quantitative and comparative data through analyses of rumen contents or feces.

The intraspecific phylogeny of the Japanese sika deer has been described. There are two genetically distinct groups of the sika deer: the northern and southern lineages (Tamate et al. 1998; Nagata et al.



**Figure 2:** The 15 measurements of mandibles of the sika deer (modified from Ozaki et al. 2007). (1) GL, greatest length of the mandible; (2) MAL, mandibular angle length; (3) CTRL, cheek tooth row length; (4) MRL, molar row length; (5) PMRL, premolar row length; (6) DL, diastema length; (7) MH\_M3, height of the mandible behind  $M_3$ ; (8) MH\_M1, height of the mandible in front of  $M_1$ ; (9) MH\_P2, height of the mandible in front of  $P_2$ ; (10) GOV\_NO, gonion ventrale to the deepest point of the mandibular notch; (11) M1B, breadth of  $M_1$ ; (12) M2B, breadth of  $M_2$ ; (13) M3B, breadth of  $M_3$ ; (14) M3L, length of  $M_3$ ; (15) M3H, height of unworn  $M_3$ ; and GOC, gonion caudale. The height of unworn  $M_3$  was measured from a cross-sectional image obtained using a microfocus X-ray CT scanning system (see Ozaki et al. 2007 for analytical details). Both scale bars are 1 cm.

1999; Goodman et al. 2001; Nagata 2009) (Fig. 1). The northern lineage includes populations in Hokkaido and most of Honshu, except for the southwestern region. The southern lineage includes populations in southwestern Honshu, Kyushu, and southwestern islands (e.g., Tsushima Island, Yakushima Island). It is believed that the separation of the two lineages occurred on the Asian continent ca. 0.3–0.5 mya, prior to their respective migration to the Japanese archipelago (Tamate et al. 1998; Nagata et al. 1999; Nagata 2009), although the timing and routes of the respective migration events have not been fully clarified.

Based on this information, there are two different levels at which microevolutionary hypotheses can be tested: between the two phylogenetic lineages and among populations within each lineage, the latter reflecting a more recent divergence than the former.


## 2. Case studies

### 2.1 Mandibular morphology

The morphology of sika deer mandibles has been examined in specimens collected from all over the Japanese archipelago (Ozaki et al. 2007). Based on prior interspecific comparative studies in ruminants, a number of morphological characteristics have been shown to correlate with grazing: e.g., broader snouts, larger areas for the insertion of the masseter muscle, longer diastemas, shorter premolar rows, and higher mandibular corpora (reviewed in Janis 1995). Some of these morphological characteristics were not statistically different after adjusting for phylogeny; however, molar row volume, molar hypsodonty, and masseter muscle mass were still significantly larger in the grazing species after adjustment (Pérez-Barbería & Gordon 1999; Williams &

**Table 2:** Results of regression analyses testing differences in mandibles of the sika deer. Significant differences between the northern and southern lineages or between the grazer and intermediate feeder populations of the northern lineage are in bold. \*In the case of the  $M_3$  hypsodonty index being the response variable, log-transformed SIZE was not included in the model because the  $M_3$  hypsodonty index was independent of the body size in the sika deer. Abbreviations of the mandibular measurements are the same as the legend of Fig. 2.

Response variable	Northern vs. southern			Grazer vs. intermediate feeder populations of northern lineage		
	Explanatory variable	Parameter estimate	P-value	Explanatory variable	Parameter estimate	P-value
Log (MAL)	Intercept	-0.194	0.462	Intercept	0.039	0.908
	Log (SIZE)	0.983	<0.001	Log (SIZE)	0.929	<0.001
	<b>Northern</b>	0.008	0.305	Grazer	0.004	0.698
Log (CTRL)	Intercept	3.445	<0.001	Intercept	3.361	<0.001
	Log (SIZE)	0.243	<0.001	Log (SIZE)	0.269	<0.001
	<b>Northern</b>	<b>0.025</b>	<b>0.026</b>	Grazer	0.009	0.310
Log (MRL)	Intercept	2.863	<0.001	Intercept	2.942	<0.001
	Log (SIZE)	0.268	<0.001	Log (SIZE)	0.256	<0.001
	<b>Northern</b>	<b>0.029</b>	<b>0.023</b>	Grazer	0.015	0.211
Log (PMRL)	Intercept	3.102	<0.001	Intercept	2.922	<0.001
	Log (SIZE)	0.096	0.069	Log (SIZE)	0.146	0.018
	<b>Northern</b>	<b>0.031</b>	<b>0.024</b>	Grazer	0.014	0.170
Log (M1B)	Intercept	1.497	<0.001	Intercept	1.565	<0.001
	Log (SIZE)	0.196	<0.001	Log (SIZE)	0.190	0.001
	<b>Northern</b>	<b>0.043</b>	<b>0.002</b>	Grazer	0.013	0.144
Log (M2B)	Intercept	0.449	0.025	Intercept	0.816	0.001
	Log (SIZE)	0.474	<0.001	Log (SIZE)	0.392	<0.001
	<b>Northern</b>	<b>0.019</b>	<b>0.039</b>	Grazer	0.008	0.384
Log (M3B)	Intercept	0.005	0.987	Intercept	0.511	0.116
	Log (SIZE)	0.579	<0.001	Log (SIZE)	0.461	<0.001
	<b>Northern</b>	0.006	0.378	Grazer	0.001	0.903
Log (M3L)	Intercept	0.719	0.001	Intercept	1.023	<0.001
	Log (SIZE)	0.557	<0.001	Log (SIZE)	0.491	<0.001
	<b>Northern</b>	<b>0.024</b>	<b>0.003</b>	Grazer	-0.002	0.810
Log (DL)	Intercept	-0.237	0.279	Intercept	0.074	0.788
	Log (SIZE)	0.991	<0.001	Log (SIZE)	0.911	<0.001
	<b>Northern</b>	<b>-0.026</b>	<b>0.002</b>	Grazer	-0.007	0.373
Log (GOV_NO)	Intercept	0.399	<0.001	Intercept	0.555	<0.001
	Log (SIZE)	0.915	<0.001	Log (SIZE)	0.878	<0.001
	<b>Northern</b>	0.000	0.947	<b>Grazer</b>	<b>0.013</b>	<b>0.009</b>
Log (MH_M3)	Intercept	0.356	0.117	Intercept	0.578	0.033
	Log (SIZE)	0.735	<0.001	Log (SIZE)	0.691	<0.001
	<b>Northern</b>	<b>0.034</b>	<b>0.002</b>	<b>Grazer</b>	<b>0.018</b>	<b>0.038</b>
Log (MH_M1)	Intercept	-3.104	<0.001	Intercept	-3.446	<0.001
	Log (SIZE)	1.476	<0.001	Log (SIZE)	1.554	<0.001
	<b>Northern</b>	-0.011	0.176	Grazer	-0.011	0.181
Log (MH_P2)	Intercept	-1.336	<0.001	Intercept	-1.460	<0.001
	Log (SIZE)	1.016	<0.001	Log (SIZE)	1.046	<0.001
	<b>Northern</b>	0.001	0.771	Grazer	0.006	0.285
$M_3$ hypsodonty index*	Intercept	1.789	<0.001	Intercept	1.833	<0.001
	<b>Northern</b>	<b>0.042</b>	<b>0.018</b>	Grazer	-0.026	0.104



**Figure 3:** Scatter plots of log-transformed mandibular measurements against those of the mandibular size (SIZE) of the sika deer: **(a)** cheek tooth row length, **(b)** M<sub>1</sub> breadth, **(c)** mandibular height behind M<sub>3</sub>, **(d)** diastema length, and **(e)** ascending ramus height (GOV\_NO). The blue (or black in black and white print) solid lines represent regression lines for the northern lineage, whereas the red (or gray in black and white print) ones represent those for the southern lineage, except for **(e)** in which the blue (or black) line represent those for the grazer population and the pale blue (or gray) one represent those for the intermediate feeder population. Significant differences between the two lineages (differences in the height of the intercepts) are found in the variables related to the cheek tooth size and the diastema length. Significant differences between grazer and intermediate feeder populations are found only in the ascending ramus height.


Kay 2001; Clauss et al. 2008). These morphological characteristics have been interpreted as adaptations to ingesting and masticating a relatively larger amount of less nutritious, more fibrous, and abrasive graminoids compared with browsers or intermediate feeders. Therefore, these morphological characteristics were compared between the two phylogenetic lineages as well as between grazer and intermediate feeder populations within the same lineage.

The data of Ozaki et al. (2007) was reanalyzed including additional data obtained after its initial publication. Mandibles of 623 female sika deer from 11 populations of the northern lineage and 6 populations from the southern lineage were used (Tab. 1). Regression analyses (mixed model) was conducted with each log-transformed mandibular measurement being the response variable. Log-transformed mandibular size (SIZE, a geometric mean of three measurements: GL, MH\_M1, and GOV\_NO; variable names defined in the legend to Fig. 2) and two categorical factors (lineage or feeding type) were explanatory variables. A population nested within a lineage or feeding type was assigned as a random effect in the mixed model. Among 15 measurements (Fig. 2), the measurements related to the cheek tooth size (lengths of the cheek tooth row, molar row, and premolar row; breadths of  $M_1$  and  $M_2$ ; length of  $M_3$  and  $M_3$  hypsodonty), mandibular height behind  $M_3$ , and diastema length were significantly different between the two lineages after adjusting for size ( $P < 0.05$ ). The northern lineage had larger and more hypodont cheek teeth, higher mandibular corpus, and shorter diastema than those of the southern lineage (Tab. 2, Figs 3a-d). On the other hand, these measurements were not significantly different between grazer and intermediate feeder populations of the same lineage, except for the mandibular corpus height behind  $M_3$ , which was larger in the grazers. In addition, a significant difference between grazer and intermediate feeder populations was observed in the ascending ramus height (Tab. 2, Fig. 3e).

Significant differences between the two lineages were observed in the cheek tooth size. The northern lineage had a buccolingually wider and mesiodistally more elongated cheek tooth row than those of the southern lineage. This result implies that the ancestral population of the northern lineage may have acquired a larger occlusal surface area of cheek teeth, which was adaptive for grazing. The shorter diastema of the northern lineage seems to be associated with the elongation of the cheek tooth row. The occlusal area of the cheek tooth row and  $M_3$  hypsodonty (which was represented by both the  $M_3$  hypsodonty index and mandibular corpus height) were significantly increased in the northern lineage. This result implies that the ancestral population of the northern lineage may have adapted to abrasive diets, possibly in a continuous or a recurrent harsh environment during the last glacial period, when possible colonization into Japan considered to have taken place (Ozaki et al. 2007). The evolution of hypsodonty will be discussed in depth later (Subsection 2.2).

On the other hand, the cheek tooth size (length, breadth, and height) was not significantly different between grazer and intermediate feeder populations of the northern lineage. This finding indicates that a further increment of the cheek tooth size may not have occurred in grazer populations. The ascending ramus height was significantly greater in grazer populations. The higher ascending ramus may be associated with a larger area for the insertion of the masseter muscle, which was adaptive for masticating fibrous graminoids.

Two hypotheses may explain the difference between the results of between-lineage comparison and between-feeding type comparison. One is that the ancestral population of the northern lineage may have acquired both larger cheek teeth and a higher ascending ramus, but under the current condition of relaxed selective pressure, the ascending ramus of intermediate feeder populations has shrunk, whereas the cheek tooth size has not changed yet (i.e., a



**Figure 4:** Correlations between the percentage of graminoids in the diet and dental variables (**a**) mesowear score, (**b**) molar wear rate) and correlations between the molar wear rate and the  $M_3$  hypsodonty index in the sika deer (**c**). The solid lines represent linear regression lines. Populations of the northern lineage are indicated by black closed circles and populations of the southern lineage are indicated by gray circles. These scatter plots are modified from Kubo & Yamada (2014).



**Figure 5:** A young female sika deer fed on dwarf bamboo at Mt. Ohdaigahara, central Japan. Note the abundance of dwarf bamboo in the forest floor and the height of the bamboo. Feeding near the ground rarely occurs. (Photograph by Sha Tech).

slower evolutionary response of cheek teeth). Another is that the ancestral population of the northern lineage may have acquired a larger cheek tooth size but not a higher ascending ramus. In populations of the northern lineage, the significant difference in the ascending ramus between grazer and intermediate feeder populations merely reflects a phenotypic (i.e., not evolutionary) response to a diet comprised of a larger amount of fibrous graminoids, which is known based on developmental experiments on laboratory animals reared on hard diets (Ito et al. 1988).

## 2.2 Molar hypsodonty and the molar wear rate

Among the morphological traits characterizing grazing ruminants, increased hypsodonty is the most remarkable and well-studied trait because of its importance for paleodietary and paleoenvironmental inferences. It is generally accepted that increased hypsodonty in herbivorous mammals is an adaptation to an abrasive diet, whereas the source and relative contribution of abrasive materials, i.e., intrinsic grass phytoliths or extrinsic abrasive materials (dust, grit, or sand), are still disputed (Sansou et al. 2007; Damuth & Janis 2011; Jardine et al. 2012; Kaiser et al. 2013; Lucas et al. 2013; Schulz et al. 2013; Stroemberg et al. 2013). The sika deer can provide a rare opportunity to answer this question because it inhabits forest environments; thus, a confounding habitat effect can be excluded. In a recently published paper (Kubo & Yamada 2014), the data of Ozaki et al. (2007, 2010) were reanalyzed with additional populations (16 populations in total, Tab. 1) to quantitatively evaluate the influence of dietary and environmental factors on 3 dental variables: mesowear score (MS), molar wear rate, and  $M_3$  hypsodonty index.

We found that the percentage of graminoids in diet of the sika deer significantly correlated with MS and the molar wear rate (regression slope of molar height against age determined using the cementum

annuli of an incisor root). A larger amount of graminoids in the diet resulted in a rounded molar cusp and a faster molar wear rate (Fig. 4a and 4b). Annual precipitation in the habitat and types of grass species (effect of feeding height) did not influence molar wear. The observed correlation between molar wear rate and proportion of graminoids in diet was not considered to be a by-product of feeding near the ground and ingesting more soil when deer crop grass. This is because the height of dwarf bamboo, on which the northern sika deer predominantly rely, is 40–60 cm; feeding near the ground rarely occurs (Fig. 5). We also found a significant correlation between estimated silica content (%) in deer diet and molar wear rate (see the supplementary analysis of Kubo & Yamada 2014 and discussion therein). Therefore, grass phytoliths can be abrasive and possibly serve as effective selective pressure for increased hypsodonty of cheek teeth, at least for Japanese sika deer. Additional analysis on the amount of soil/grit in deer diet is needed to evaluate the relative importance of grass phytoliths against that of the external abrasives. We also found a positive correlation between the molar wear rate and the  $M_3$  hypsodonty index (Fig. 4c). These data indicated an evolutionary increment in the molar height corresponding to the molar wear rate (selective pressure). However, we have some reservations regarding this conclusion because the degree of hypsodonty differed significantly between the northern and southern lineages (Tab. 2). In Fig. 4c, the hypsodonty index increased with the accelerating molar wear rate in the southern lineage but not in the northern lineage. Genetic evidence indicated that the greatest genetic differentiation was found among populations of the southern lineage, whereas among populations of the northern lineage, this differentiation occurred more recently because of habitat fragmentation. Therefore, according to the findings of comparative analyses of mandibles (Subsection 2.1), we hypothesize that the ancestral population of the northern lineage acquired more hypsodont molars in the past, possibly under strong environmental pressure that promoted molar wear. We also hypothesize that after subdivision into local populations, molar height did not change measurably because of relaxed selection and/or insufficient duration of selection, particularly in populations of the northern lineage. In the paleontological, macroevolutionary context, it has been discussed once evolution of increased hypsodonty occurred, it does not easily reverse because cost of maintaining hypsodonty may be low (see discussion in Fortelius 1985, Hummel et al. 2010). This interpretation can be concordant with our observation that intermediate feeder populations of the northern lineage have the equivalent degree of hypsodonty to those of the grazer populations.

### 2.3 Limb morphology

Interspecific variation in the limb length in ungulates has been investigated from both aspects of biomechanics (McMahon 1973, 1975; Alexander et al. 1979) and locomotory adaptation to specific habitats (Scott 1985, 1987; Garland & Janis 1993; Janis et al. 2012). The former predicts an allometric relationship between body mass and the limb length; the limb length is proportional to  $(\text{body mass})^{0.25}$ , whereas the latter focuses on the departure from the predictions with reference to ecological characteristics. It is of great interest if general biomechanical predictions are also applicable even at the species level, and if this body size-limb length relationship is related to some ecological or behavioral characteristics.

In the sika deer, postcranial morphological traits have been less investigated because most specimens of the sika deer housed in museums are limited to skulls and mandibles. Based on measurements of the culled sika deer of a northern population, Takatsuki (1992) showed that the sika deer cannot live in areas with heavy snowfall, where snow accumulates at a level higher than the limb length of the sika deer. Ochiai & Asada (1995) reported a growth pattern of the body size of the sika deer from a population in a temperate region (Boso Peninsula) and found that the hind limb length was relatively (compared with shoulder height) shorter than that of other northern populations. This finding implies that the amount of snowfall or habitat closedness may influence variation in the hind limb length. In the subsequent paper, Ochiai & Asada (1997) compared the size of the skull and limb bones of the Boso population with those of a northern population (Nikko). It was shown that the limb length was proportional to  $(\text{body mass})^{0.24}$ , thus confirming the biomechanical prediction of McMahon (1973). These data also implied that the positive deviation of the Nikko deer from the allometric prediction line can be explained by snow depth of the habitat. Recently, Terada et al. (2012) showed morphological differentiation in limbs among southern island populations of the sika deer. They used 8 populations belonging to the southern lineage and measured both the skull size [condylobasal length (CBL)] and the length of limb bones (scapula, humerus, radius, metacarpus, femur, tibia, and metatarsus). They showed that there was significant variation in the measurements among the populations, particularly in CBL and the metacarpal length. In analyses, they regressed the log-transformed metacarpal length against log-transformed CBL (a proxy of the body size) and obtained the regression slope and intercept for each population. The slope did not differ significantly among the populations and had a value of 0.74, which was significantly smaller than 1 (i.e., negative allometry). This value is close to the biomechanical prediction as well as to the empirical regression slope obtained for other sika deer populations (Ochiai & Asada 1997), provided that CBL is

isometric to body mass ( $\text{CBL}^3 = \text{body mass}$ ). On the other hand, the regression intercept (relative carpal length adjusted for the body size) negatively correlated with habitat topology. This was quantitatively evaluated using a geographical information system (GIS); the steeper the habitat area, the shorter the metacarpal length. As Terada et al. (2012) discussed in their paper, this result is in agreement with the interspecific pattern of bovid species; species in mountainous habitats have shorter limbs than those in open habitats (Scott 1985, 1987). Considering the fact that annual snowfall in less than 5cm for the studied habitats of sika deer, Terada et al. (2012) discussed that shorter metacarpal may be favored because it can increase stability in rugged terrain. The relationship between habitat topology and limb morphology has not been studied in other populations, particularly in those of the northern lineage. Together with the observations of Takatsuki (1992) and Ochiai & Asada (1997), it is expected that limb morphology of the northern sika deer is affected by both habitat topology and the amount of snowfall.

Terada et al. (2012) have also convincingly demonstrated that habitat topology can be quantitatively assessed using GIS; this will of great use to interspecific comparison with quantitative categorization of habitat types.

### 2.4 Body size

Evolutionary changes in the body size is one of the most studied issues in evolutionary biology because it affects so many aspects of the animal's biology (Schmidt-Nielsen 1984). Among ruminant species, the evolution of body size has been assessed with reference to ecological diversification (Bell 1970; Jarman 1974; Geist 1999; Pérez-Barbería et al. 2002). On the other hand, changes in body size within a species are confounded by phenotypic plasticity due to environmental factors. Body size (mass) can be affected greatly by nutritional conditions. To separate genetic and environmental effects, "common garden" type experiments or reciprocal transplants of target populations must be used. These can be very difficult to conduct for large, wild mammals (see Pelletier et al. 2007; Wilson et al. 2007 for studies of the genetic basis of the body size of wild sheep populations). Therefore, intraspecific studies of body size variation in mammals have focused on identifying an empirical relationship between the body size and environmental factors (Gay & Best 1996; Wigginton & Dobson 1999; Smith et al. 2002; Wolverton et al. 2009; Gür 2010). In many cases, this relationship was discussed from the perspective of Bergmann's rule, which describes an increase in the body size of endotherms at high latitudes (Bergmann 1847).

Because the Japanese sika deer shows considerable latitudinal variation in the body size among local populations, it has been considered as an ex-

ample that follows Bergmann's rule (Ohtaishi 1986). Body mass of adult males during the winter season varies with latitude; 120 kg in the northernmost population (Hokkaido) but only 30 kg in the southernmost population (Kerama Islands) (Ohtaishi 1986). Most mainland (Honshu) populations show intermediate values (60 kg in the central area to 100 kg in the northern area). However, quantitative measurements of the body size of local populations were limited at the time of publication of the report by Ohtaishi (1986). Therefore, further accumulation of body size data is needed to properly evaluate Bergmann's rule for Japanese sika deer.

There have been several reports on geographic variation in the skull size among sika deer populations. Imaizumi (1970) described a latitudinal cline in CBL for the purpose of species classification in what he termed the subgenus *Sika*. His taxonomic splitting of the sika deer into several species is not generally accepted (Ohtaishi 1986; Endo 1996). Imaizumi (1970) reported that "*C. nippon*" (most Japanese populations, except for Hokkaido, Yakushima Island, Kerama Islands, and Tsushima Islands) showed a cline in CBL that is larger in the north and smaller in the south. This regression slope was based on only 7 individual males. Thus, further data are needed to clarify latitudinal variation in the skull size. Shiroma & Ohta (1996) compared skull morphology among 8 populations (Hokkaido, Nikko, Kagoshima, Nozakijima Island, Tsushima Island, Mageshima Island, Yakushima Island, and Kerama Islands) and showed that the northernmost population (Hokkaido) had the largest skull size followed by Nikko, Kagoshima, and other island populations. Because the sampled populations were biased to island populations, it is still unclear whether the latitudinal cline followed Bergmann's rule. Terada et al. (2012) found that CBL was negatively related to temperature and latitude; these findings are congruent with Bergmann's rule among 8 southern island populations.

In summary, although geographical variation in the body size is conspicuous in the Japanese sika deer, there has been no comprehensive study of quantitative data of populations from all over the Japanese archipelago. Recently, both body mass and skull measurements were collected for more than 30 sika deer populations by us. These data will be reported in the near future.

### 3. Conclusions

The focus of this review was the question "Are interspecific and macroevolutionary trends in ruminant morphology also found among populations of a single species?" Several case studies of the Japanese sika deer were conducted by our group and other researchers. Several conclusions can be drawn from these studies: 1) selective pressures, such as accelerated tooth wear associated with grass consump-

on or topographical features of a habitat that require climbing ability, promote local adaptation of populations within an ecologically diverse species, similar to the situations in which these same selective pressures govern interspecific variation among ruminant species; 2) in some cases, genetic divergence among local populations is insufficient in duration, resulting in a smaller extent of morphological change compared with that in subspecific lineages or closely related species; and 3) a phenotypic change resulting from nutritional conditions affects intraspecific comparison to some extent, depending on traits. This may confound the conclusion that the observed morphological differentiation is caused by adaptation. To confirm the third conclusion, some genetic models of traits are needed that involve pedigree analysis or "common garden" experiments, both of which are difficult to conduct for wild mammals.

There are still several traits, not only morphological but also physiological and behavioral, to be investigated for intraspecific variations or trends in the sika deer. For example, geographical variation in the size and morphology of antlers is an attractive topic with reference to the evolution of antlers in Cervidae (Gould 1974; Geist 1999). Miura (1984) reported geographical variation in mating behavior among local populations. This implies that habitat closedness may have a strong impact on the degree of aggregation and thus the mating system of the sika deer. This is in agreement with the evolutionary trend observed across ruminant species (Jarman 1974; Pérez-Barbería et al. 2002). Whether this variation in the mating system is related to antler variation among local deer populations is another intriguing question.

### Acknowledgements

The author would like to thank G. Rößner and co-organizers of the conference for the opportunity to present my results. The guest editor W. Silvia, M. Fortelius and J. Damuth were greatly appreciated for their constructive comments. The author is grateful to G. Suwa and S. Takatsuki for their support throughout her doctoral research. She is grateful to the following people for their permission to use the sika deer specimens in their charge: N. Ohtaishi, M. Eda, K. Kaji, T. Yoshida, T. Hayashi, H. Endo, H. Taru, K. Ochiai, and T. Koizumi. The author thanks all the people who helped her to collect sika deer specimens housed in The University Museum, The University of Tokyo. This study was supported by a Grant-in-Aid for JSPS Fellows (25-10506).

### 4. References

Alexander RM, Jayes AS, Maloiy GMO, Wathuta EM. 1979. Allometry of the limb bones of mammals from shrews (*Sorex*) to elephant (*Loxodonta*). Journal of Zoology 189, 305-314.

Bell R. 1970. The use of the herb layer by grazing ungulates in the Serengeti. In: A Watson (Ed.), *Animal populations in relation to their food resources*. Oxford, Blackwell Scientific, 111-124.

Bergmann C. 1847. Über die Verhältnisse der Wärmeökonomie der Thiere zu ihrer Grösse. *Göttinger Studien* 1, 595-708.

Clauss M, Lechner-Doll M, Streich WJ. 2003. Ruminant diversification as an adaptation to the physicomechanical characteristics of forage. A reevaluation of an old debate and a new hypothesis. *Oikos* 102, 253-262.

Clauss M, Hofmann RR, Streich WJ, Fickel J, Hummel J. 2008. Higher masseter muscle mass in grazing than in browsing ruminants. *Oecologia* 157, 377-385.

Clutton-Brock TH, Guinness FE, Albon SD. 1984. Individuals and populations: the effects of social behaviour on population dynamics in deer. *Proceedings of the Royal Society of Edinburgh B* 82, 275-290.

Clutton-Brock TH, Illius AW, Wilson K, Grenfell BT, MacColl ADC, Albon SD. 1997. Stability and instability in ungulate populations: An empirical analysis. *American Naturalist* 149, 195-219.

Codron D, Clauss M. 2010. Rumen physiology constrains diet niche: linking digestive physiology and food selection across wild ruminant species. *Canadian Journal of Zoology-Revue Canadienne De Zoologie* 88, 1129-1138.

Damuth J, Janis CM. 2011. On the relationship between hypsodonty and feeding ecology in ungulate mammals, and its utility in palaeoecology. *Biological Reviews* 86, 733-758.

Davis EB, Brakora KA, Lee AH. 2011. Evolution of ruminant headgear: a review. *Proceedings of the Royal Society B* 278, 2857-2865.

Endo H. 1996. Scientific and Japanese names of artiodactyls of Japan. *Honyurui Kagaku (Mammalian Science)* 35, 203-209 (in Japanese with English abstract).

Festa-Bianchet M, Gaillard JM, Cote SD. 2003. Variable age structure and apparent density dependence in survival of adult ungulates. *Journal of Animal Ecology* 72, 640-649.

Fortelius M. 1985. Ungulate cheek teeth: developmental, functional, and evolutionary interrelations. *Acta Zoologica Fennica* 180, 1-76.

Foster SA, Endler JA eds. 1999. *Geographic variation in behavior: perspectives on evolutionary mechanisms*. New York, Oxford University Press, 336 p.

Gür H. 2010. Why do Anatolian ground squirrels exhibit a Bergmannian size pattern? A phylogenetic comparative analysis of geographic variation in body size. *Biological Journal of the Linnean Society* 100, 695-710.

Gaillard JM, Delorme D, Boutin JM, Vanlaere G, Boisaubert B, Pradel R. 1993. Roe deer survival patterns - a comparative analysis of contrasting populations. *Journal of Animal Ecology* 62, 778-791.

Garland T, Janis CM. 1993. Does metatarsal femur ratio predict maximal running speed in cursorial mammals. *Journal of Zoology* 229, 133-151.

Gay SW, Best TL. 1996. Relationships between abiotic variables and geographic variation in skulls of pumas (*Puma concolor*: Mammalia, Felidae) in North and South America. *Zoological Journal of the Linnean Society* 117, 259-282.

Geist V. 1999. *Deer of the World*. Shrewsbury, Swan Hill Press, 421 p.

Goodman SJ, Tamate HB, Wilson R, Nagata J, Tatsuzawa S, Swanson GM, Pemberton JM, McCullough DR. 2001. Bottlenecks, drift and differentiation: the population structure and demographic history of sika deer (*Cervus nippon*) in the Japanese archipelago. *Molecular Ecology* 10, 1357-1370.

Gould SJ. 1974. Origin and function of bizarre structures - antler size and skull size in Irish elk, *Megaceros giganteus*. *Evolution* 28, 191-220.

Hlusko LJ, Weiss KM, Mahaney MC. 2002. Statistical genetic comparison of two techniques for assessing molar crown size in pedigree baboons. *American Journal of Physical Anthropology* 117, 182-189.

Hofmann RR. 1985. Digestive physiology of the deer - their morphophysiological specialisation and adaptation. In: Fennessy PF, Drew KR (Eds), *Biology of Deer Production*, 393-407.

Hofmann RR. 1989. Evolutionary steps of ecophysiological adaptation and diversification of ruminants - a comparative view of their digestive-system. *Oecologia* 78, 443-457.

Hofmann RR, Streich WJ, Fickel J, Hummel J, Clauss M. 2008. Convergent evolution in feeding types: salivary gland mass differences in wild ruminant species. *Journal of Morphology* 269, 240-257.

Hummel J, Findeisen E, Suedekum K-H, Ruf I, Kaiser TM, Bucher M, Clauss M, Codron D. 2011. Another one bites the dust: faecal silica levels in large herbivores correlate with high-crowned teeth. *Proceedings of the Royal Society B* 278, 1742-1747.

Imaizumi Y. 1970. Description of a new species of *Cervus* from the Tushima Islands, Japan, with a revision of the subgenus *Sika* based on clinal analysis. *Bulletin of National Science Museum*, Tokyo 13, 185-194.

Ito G, Mitani S, Kim JH. 1988. Effect of soft diets on craniofacial growth in mice. *Anatomischer Anzeiger* 165, 151-166.

Janis CM. 1995. Correlations between craniodontal morphology and feeding behavior in ungulates: reciprocal illumination between living and fossil taxa. In: Thomason JJ (Ed) *Functional morphology in vertebrate paleontology*. Cambridge, Cambridge University Press, 76-98.

Janis CM, Shoshitaishvili B, Kambic R, Figueirido B. 2012. On their knees: distal femur asymmetry in ungulates and its relationship to body size and locomotion. *Journal of Vertebrate Paleontology* 32, 433-445.

Jardine PE, Janis CM, Sahney S, Benton MJ. 2012. Grit not grass: concordant patterns of early origin of hypsodonty in Great Plains ungulates and Glires. *Palaeogeography Palaeoclimatology, Palaeoecology* 365, 1-10.

Jarman PJ. 1974. Social-organization of antelope in relation to their ecology. *Behaviour* 48, 215-266.

Kaiser TM, Mueller DWH, Fortelius M, Schulz E, Codron D, Clauss M. 2013. Hypsodonty and tooth facet development in relation to diet and habitat in herbivorous ungulates: implications for understanding tooth wear. *Mammal Review* 43, 34-46.

Kubo MO, Yamada E. 2014. The inter-relationship between dietary and environmental properties and tooth wear: comparisons of mesowear, molar wear rate, and hypsodonty index of extant sika deer populations. *PLoS ONE* 9, e90745.

Kubo MO, Kaji K, Ohba T, Hosoi E, Koizumi T, Takatsuki S. 2011. Compensatory response of molar eruption for environment-mediated tooth wear in sika deer. *Journal of Mammalogy* 92, 1407-1417.

Kubo MO, Minami M, Higuchi N, Ohnishi N, Okada A, Kaji K, Ohba T, Hosoi E, Koizumi T, Takatsuki S. 2013. Female sika deer have evolved larger incisors than males under relaxed selection against rapid tooth wear. *Biological Journal of the Linnean Society* 110, 384-397.

Lucas PW, Omar R, Al-Fadhalah K, Almusallam AS, Henry AG, Michael S, Thai LA, Watzke J, Strait DS, Atkins AG. 2013. Mechanisms and causes of wear in tooth enamel: implications for hominin diets. *Journal of the Royal Society Interface* 10.

Ma JF, Takahashi E. 2002. *Soil, Fertilizer, and Plant Silicon Research in Japan*. Amsterdam, Elsevier Science B, V, 294 p.

Martinez-Jauregui M, San Miguel-Ayanz A, Mysterud A, Rodriguez-Vigal C, Clutton-Brock T, Langvatn R, Coulson T. 2009. Are local weather, NDVI and NAO consistent determinants of red deer weight across three contrasting European countries? *Global Change Biology* 15, 1727-1738.

McCullough DR, Takatsuki S, Kaji K (Eds). 2009. *Sika Deer: Biology and Management of Native and Introduced Populations*. Tokyo, Springer, 666 p.

McMahon T. 1973. Size and shape in biology. *Science* 179, 1201-1204.

McMahon T. 1975. Allometry and biomechanics - limb bones in adult ungulates. *American Naturalist* 109, 547-563.

Miura S. 1984. Social behavior and territoriality in male sika deer (*Cervus nippon* Temminck 1838) during the rut. *Zeitschrift für Tierpsychologie* 64, 33-73.

Motomura H, Fujii T, Suzuki M. 2000. Distribution of silicified cells

in the leaf blades of *Pleioblastus chino* (Franchet et Savatier) Makino (Bambooideae). *Annals of Botany* 85, 751-757.

Mysterud A, Yoccoz NG, Stenseth NC, Langvatn R. 2001. Effects of age, sex and density on body weight of Norwegian red deer: evidence of density-dependent senescence. *Proceedings of the Royal Society of London Series B* 268, 911-919.

Nagata J. 2009. Two genetically distinct lineages of the Japanese sika deer based on mitochondrial control regions. In: DR McCullough, S Takatsuki, K Kaji (Eds), *Sika Deer: Biology and Management of Native and Introduced Populations*. Tokyo, Springer, 27-41.

Nagata J, Masuda R, Tamate HB, Hamasaki S, Ochiai K, Asada M, Tatsuzawa S, Suda K, Tado H, Yoshida MC. 1999. Two genetically distinct lineages of the sika deer, *Cervus nippon*, in Japanese islands: comparison of mitochondrial D-loop region sequences. *Molecular Phylogenetics and Evolution* 13, 511-519.

Ochiai K, Asada M. 1995. Growth in the body size of sika deer (*Cervus nippon*) on the Boso Peninsula, central Japan. *Journal of the Natural History Museum and Institute, Chiba* 3, 223-232 (in Japanese with English abstract).

Ochiai K, Asada M. 1997. Growth pattern of the skull and limb bone size of sika deer, *Cervus nippon*, on the Boso Peninsula, central Japan. *Journal of the Natural History Museum and Institute, Chiba* 4, 159-172 (in Japanese with English abstract).

Ohtaishi N. 1986. Preliminary memorandum of classification, distribution and geographic variation on Sika deer. *Honyurui Kagaku (Mammalian Science)* 53, 13-17 (in Japanese with English abstract).

Ozaki M, Suwa G, Kaji K, Ohba T, Hosoi E, Koizumi T, Takatsuki S. 2007. Correlations between feeding type and mandibular morphology in the sika deer. *Journal of Zoology (London)* 272, 244-257.

Ozaki M, Kaji K, Matsuda N, Ochiai K, Asada M, Ohba T, Hosoi E, Tado H, Koizumi T, Suwa G, Takatsuki S. 2010. The relationship between food habits, molar wear and life expectancy in wild sika deer populations. *Journal of Zoology* 280, 202-212.

Pérez-Barbería FJ, Gordon IJ. 1999. The functional relationship between feeding type and jaw and cranial morphology in ungulates. *Oecologia* 118, 157-165.

Pérez-Barbería FJ, Gordon IJ, Pagel M. 2002. The origins of sexual dimorphism in body size in ungulates. *Evolution* 56, 1276-1285.

Pelletier F, Reale D, Garant D, Coltman DW, Festa-Bianchet M. 2007. Selection on heritable seasonal phenotypic plasticity of body mass. *Evolution* 61, 1969-1979.

Sanson GD, Kerr SA, Gross KA. 2007. Do silica phytoliths really wear mammalian teeth? *Journal of Archaeological Science* 34, 526-531.

Schmidt-Nielsen K. 1984. *Scaling: why is animal size so important?* Cambridge, Cambridge University Press, 256 p.

Schulz E, Piotrowski V, Clauss M, Mau M, Merceron G, Kaiser TM. 2013. Dietary abrasiveness is associated with variability of microwear and dental surface texture in rabbits. *PLoS ONE* 8, e56167.

Scott KM. 1985. Allometric trends and locomotor adaptations in the Bovidae. *Bulletin of the American Museum of Natural History* 179, 197-288.

Scott KM. 1987. Allometry and habitat-related adaptations in the postcranial skeleton of Cervidae. In: CM Wemmer (Ed.), *Biology and management of the Cervidae*, Washington D.C., Smithsonian Institution Press, 65-80.

Shiroma T, Ohta H. 1996. Geographical variation in cranial morphology in sika deer with a special reference on Kerama deer. In: Okinawa Prefectural Board of Education (Ed.) *A report of conservation and management of sika deer in Kerama Islands*. Okinawa, Okinawa Prefectural Board of Education, 13-55.

Smith PA, Schaefer JA, Patterson BR. 2002. Variation at high latitudes: the geography of body size and cranial morphology of the muskox, *Ovibos moschatus*. *Journal of Biogeography* 29, 1089-1094.

Stroemberg CAE, Dunn RE, Madden RH, Kohn MJ, Carlini AA. 2013. Decoupling the spread of grasslands from the evolution of grazer-type herbivores in South America. *Nature Communications* 4, 1478.

Takatsuki S. 1992. Foot morphology and distribution of sika deer in relation to snow depth in Japan. *Ecological Research* 7, 19-23.

Takatsuki S. 2009. Geographical variations in food habits of sika deer: the northern grazer vs. the southern browser. In: McCullough DR, Takatsuki S, Kaji K (Eds), *Sika Deer: Biology and Management of Native and Introduced Populations*. Tokyo, Springer, 231-238.

Tamate HB, Tatsuzawa S, Suda K, Izawa M, Doi T, Sunagawa K, Miyahira F, Tado H. 1998. Mitochondrial DNA variations in local populations of the Japanese sika deer, *Cervus nippon*. *Journal of Mammalogy* 79, 1396-1403.

Terada C, Tatsuzawa S, Saitoh T. 2012. Ecological correlates and determinants in the geographical variation of deer morphology. *Oecologia* 169, 981-994.

Van Soest PJ. 1994. *Nutritional ecology of the ruminant*. New York, Cornell University Press, 488 p.

Wigginton JD, Dobson FS. 1999. Environmental influences on geographic variation in body size of western bobcats. *Canadian Journal of Zoology-Revue Canadienne De Zoologie* 77, 802-813.

Williams SH, Kay RF. 2001. A comparative test of adaptive explanations for hypsodonty in ungulates and rodents. *Journal of Mammalian Evolution* 8, 207-229.

Wilson AJ, Pemberton JM, Pilkington JG, Clutton-Brock TH, Coltman DW, Kruuk LEB. 2007. Quantitative genetics of growth and cryptic evolution of body size in an island population. *Evolutionary Ecology* 21, 337-356.

Wolverton S, Huston MA, Kennedy JH, Cagle K, Cornelius JD. 2009. Conformation to Bergmann's rule in white-tailed deer can be explained by food availability. *American Midland Naturalist* 162, 403-417.

Yokoyama M, Kaji K, Suzuki M. 2000. Food habits of sika deer and nutritional value of sika deer diets in eastern Hokkaido, Japan. *Ecological Research* 15, 345-355.