

COMMENTARY

Open Access

Apoptosis induction and tumor cell repopulation: The yin and yang of radiotherapy

Kirsten Lauber^{1*}, Luis E Munoz², Christian Berens³, Verena Jendrosseck⁴, Claus Belka¹ and Martin Herrmann²

The induction of tumor cell death is a central goal of radiotherapy. Surprisingly, a recent study sheds new light on this process, and the results presented by Huang et al. strongly question the benefit of radiation-induced apoptosis for the outcome of cancer radiotherapy.

In their study, Huang and coworkers describe that induction of apoptosis by radiotherapy stimulates rapid tumor cell repopulation - a process crucially dependent on caspase 3 activity [1]. These findings are of immense relevance for the clinical use of radiotherapy, particularly when combined with targeted agents aiming at radiosensibilization and enhanced apoptosis induction [2-8]. In contrast to the results presented by Huang et al., previous work has convincingly demonstrated that the increased induction of apoptotic cell death, for example by the combination of agonistic TRAIL antibodies with radiation, results in a pronounced benefit for long term tumor control in a colorectal xenograft model [9-11]. Parallel *in vitro* studies have revealed that enhanced caspase-mediated apoptosis is the underlying mechanism for the improved eradication of clonogenic tumor cells [10,12]. Thus, the effect described by Huang and coworkers should be considered as a repopulation mechanism, which is of importance under specific, currently unknown circumstances. In this regard, it can be speculated that the balance between the apoptotic net cell kill, and the PGE₂-driven tumor cell survival and repopulation accounts for the reported discrepancies.

Nevertheless, the study by Huang and coworkers is highly interesting, in particular because of the elegant experiments, with which the signaling cascade of apoptosis-induced tumor cell repopulation was unraveled. The downstream mechanisms identified by Huang et al. involve the caspase 3-dependent cleavage and activation of iPLA₂ and the subsequent production of PGE₂. During radiation-induced cell death *in vitro* PGE₂ was shown to be released by tumor cells as well as by


fibroblasts. *In vivo* (in experimental mouse models), both tumor and tumor stroma cells reportedly contributed to rapid tumor cell repopulation by few residual tumor cells in response to radiotherapy.

We would like to point out that the tumor stroma contains a highly interesting cell population, which might contribute to or even dominate the tumor-growth-stimulating PGE₂ production: macrophages that govern the elimination of apoptosing cells and instigate tissue healing by producing a clearance-related cytokine milieu, including PGE₂ [13-15]. Of note, Huang and coworkers observed that more macrophages were present in irradiated (apoptotic) tumors than in non-irradiated ones. These phagocytes have presumably been recruited by apoptotic cell-derived find-me signals, such as nucleotides and lysophosphatidylcholine, which - akin to PGE₂ - are released in a caspase 3- or caspase 3- plus iPLA₂-dependent manner, respectively [16-19]. Caspase 3 apparently is a key player in this context. So it should be taken into consideration that caspase 3 controls more processes than the release of PGE₂ or phagocyte-recruiting attraction signals by apoptotic cells. Caspase 3 also shifts the balance between apoptosis, necrosis and autophagy as described by Huang et al., and orchestrates the central features of apoptosis, which have profound impact on macrophage activation and cytokine production after the engulfment of apoptotic cells. As such, externalization of phosphatidylserine, bleb formation and internucleosomal DNA fragmentation, known to be crucial for the subsequent anti-inflammatory cytokine production by macrophages [20-22], have been reported to depend on caspase 3 activity during apoptosis [23-27]. Thus, caspase 3-positive apoptosing cells recruit more macrophages, are more efficiently phagocytosed, and induce a stronger anti-inflammatory, wound-healing and growth promoting phagocyte response, including PGE₂ production, than their caspase 3-negative counterparts. This might contribute or translate to the clinical observation by Huang et al. that elevated expression levels of activated caspase-3 were associated with a poor

* Correspondence: kirsten.lauber@med.uni-muenchen.de

¹Department of Radiation Oncology, LMU Munich, Germany

Full list of author information is available at the end of the article



outcome in two patient cohorts with head and neck carcinoma or with advanced stage breast carcinoma.

Overall, Huang and coworkers suggest a scenario, in which the caspase 3-driven iPLA₂-dependent PGE₂ production by irradiated tumor and tumor stroma cells plays a pivotal role for radiation-induced tumor cell repopulation and for poor therapeutic outcome. We would like to add the clearance of apoptosing cells by macrophages, and the subsequently produced clearance-related anti-inflammatory milieu, including PGE₂, to this model (Figure 1). Intriguingly, in their final step both processes rely on cyclooxygenase activity, thus re-opening the therapeutic perspective of cautious cyclooxygenase inhibition as an adjuvant to radiotherapy [28-30]. Up to now several clinical trials have documented that a safe combination of cyclooxygenase inhibitors (celecoxib) and radiotherapy is feasible, yet most of the trials were not adequately powered to detect meaningful differences in tumor control [31,32]. Thus, further clinical trials, in particular phase III studies, are required to shed light onto this issue. In the same line, it should be addressed, whether caspase inhibition (upstream of PGE₂ production) in combination with radiotherapy displays a benefit for the overall therapeutic outcome - provided that tumor cell systems utilizing the caspase 3-

dependent, PGE₂-driven tumor cell repopulation can reliably be identified.

Author details

¹Department of Radiation Oncology, LMU Munich, Germany. ²Department of Internal Medicine 3, Friedrich-Alexander-University Erlangen-Nuremberg, Germany. ³Department of Biology, Friedrich-Alexander-University Erlangen-Nuremberg, Germany. ⁴Institute of Cell biology, University Hospital Essen, Germany.

Received: 8 December 2011 Accepted: 19 December 2011

Published: 19 December 2011

References

- Huang Q, Li F, Liu X, Li W, Shi W, Liu FF, O'Sullivan B, He Z, Peng Y, Tan AC, et al: Caspase 3-mediated stimulation of tumor cell repopulation during cancer radiotherapy. *Nat Med* 2011, **17**(7):860-866.
- Connell PP, Weichselbaum RR: A downside to apoptosis in cancer therapy? *Nat Med* 2011, **17**(7):780-782.
- Itani W, Geara F, Haykal J, Haddadin M, Gali-Muhtasib H: Radiosensitization by 2-benzoyl-3-phenyl-6,7-dichloroquinoxaline 1,4-dioxide under oxia and hypoxia in human colon cancer cells. *Radiat Oncol* 2007, **2**:1.
- Liao X, Che X, Zhao W, Zhang D, Long H, Chaudhary P, Li H: Effects of propranolol in combination with radiation on apoptosis and survival of gastric cancer cells in vitro. *Radiat Oncol* 2010, **5**:98.
- Rubel A, Handrick R, Lindner LH, Steiger M, Eibl H, Budach W, Belka C, Jendrossek V: The membrane targeted apoptosis modulators erucylphosphocholine and erucylphosphohomocholine increase the radiation response of human glioblastoma cell lines in vitro. *Radiat Oncol* 2006, **1**:6.
- Rudner J, Ruiner CE, Handrick R, Eibl HJ, Belka C, Jendrossek V: The Akt-inhibitor Erufosine induces apoptotic cell death in prostate cancer cells

- and increases the short term effects of ionizing radiation. *Radiat Oncol* 2010, **5**:108.
7. Welsh JW, Mahadevan D, Ellsworth R, Cooke L, Bearss D, Stea B: **The c-Met receptor tyrosine kinase inhibitor MP470 radiosensitizes glioblastoma cells.** *Radiat Oncol* 2009, **4**:69.
 8. Zerp SF, Stoter R, Kuipers G, Yang D, Lippman ME, van Blitterswijk WJ, Bartelink H, Rooswinkel R, Lafleur V, Verheij M: **AT-101, a small molecule inhibitor of anti-apoptotic Bcl-2 family members, activates the SAPK/JNK pathway and enhances radiation-induced apoptosis.** *Radiat Oncol* 2009, **4**:47.
 9. Marini P, Budach W, Niyazi M, Junginger D, Stickl S, Jendrossek V, Belka C: **Combination of the pro-apoptotic TRAIL-receptor antibody mapatumumab with ionizing radiation strongly increases long-term tumor control under ambient and hypoxic conditions.** *Int J Radiat Oncol Biol Phys* 2009, **75**(1):198-202.
 10. Marini P, Denzinger S, Schiller D, Kauder S, Welz S, Humphreys R, Daniel PT, Jendrossek V, Budach W, Belka C: **Combined treatment of colorectal tumours with agonistic TRAIL receptor antibodies HGS-ETR1 and HGS-ETR2 and radiotherapy: enhanced effects in vitro and dose-dependent growth delay in vivo.** *Oncogene* 2006, **25**(37):5145-5154.
 11. Marini P, Junginger D, Stickl S, Budach W, Niyazi M, Belka C: **Combined treatment with lexatumumab and irradiation leads to strongly increased long term tumour control under normoxic and hypoxic conditions.** *Radiat Oncol* 2009, **4**:49.
 12. Marini P, Jendrossek V, Durand E, Gruber C, Budach W, Belka C: **Molecular requirements for the combined effects of TRAIL and ionising radiation.** *Radiother Oncol* 2003, **68**(2):189-198.
 13. Fadok VA, Bratton DL, Konowal A, Freed PW, Westcott JY, Henson PM: **Macrophages that have ingested apoptotic cells in vitro inhibit proinflammatory cytokine production through autocrine/paracrine mechanisms involving TGF-beta, PGE2, and PAF.** *J Clin Invest* 1998, **101**(4):890-898.
 14. Voll RE, Herrmann M, Roth EA, Stach C, Kalden JR, Girkontaite I: **Immunosuppressive effects of apoptotic cells.** *Nature* 1997, **390**(6658):350-351.
 15. Mosser DM, Edwards JP: **Exploring the full spectrum of macrophage activation.** *Nat Rev Immunol* 2008, **8**(12):958-969.
 16. Elliott MR, Chekeni FB, Trampont PC, Lazarowski ER, Kadl A, Walk SF, Park D, Woodson RI, Ostankovich M, Sharma P, *et al*: **Nucleotides released by apoptotic cells act as a find-me signal to promote phagocytic clearance.** *Nature* 2009, **461**(7261):282-286.
 17. Lauber K, Bohn E, Krober SM, Xiao YJ, Blumenthal SG, Lindemann RK, Marini P, Wiedig C, Zobywalski A, Baksh S, *et al*: **Apoptotic cells induce migration of phagocytes via caspase-3-mediated release of a lipid attraction signal.** *Cell* 2003, **113**(6):717-730.
 18. Mueller RB, Sheriff A, Gaipl US, Wesselborg S, Lauber K: **Attraction of phagocytes by apoptotic cells is mediated by lysophosphatidylcholine.** *Autoimmunity* 2007, **40**(4):342-344.
 19. Peter C, Waibel M, Radu CG, Yang LV, Witte ON, Schulze-Osthoff K, Wesselborg S, Lauber K: **Migration to apoptotic "find-me" signals is mediated via the phagocyte receptor G2A.** *J Biol Chem* 2008, **283**(9):5296-5305.
 20. Huynh ML, Fadok VA, Henson PM: **Phosphatidylserine-dependent ingestion of apoptotic cells promotes TGF-beta1 secretion and the resolution of inflammation.** *J Clin Invest* 2002, **109**(1):41-50.
 21. Stadler K, Frey B, Munoz LE, Finzel S, Rech J, Fietkau R, Herrmann M, Hueber A, Gaipl US: **Photopheresis with UV-A light and 8-methoxypsoralen leads to cell death and to release of blebs with anti-inflammatory phenotype in activated and non-activated lymphocytes.** *Biochem Biophys Res Commun* 2009, **386**(1):71-76.
 22. Nagata S: **DNA degradation in development and programmed cell death.** *Annu Rev Immunol* 2005, **23**:853-875.
 23. Coleman ML, Sahai EA, Yeo M, Bosch M, Dewar A, Olson MF: **Membrane blebbing during apoptosis results from caspase-mediated activation of ROCK I.** *Nat Cell Biol* 2001, **3**(4):339-345.
 24. Sebbagh M, Renvoize C, Hamelin J, Riche N, Bertoglio J, Breard J: **Caspase-3-mediated cleavage of ROCK I induces MLC phosphorylation and apoptotic membrane blebbing.** *Nat Cell Biol* 2001, **3**(4):346-352.
 25. Enari M, Sakahira H, Yokoyama H, Okawa K, Iwamatsu A, Nagata S: **A caspase-activated DNase that degrades DNA during apoptosis, and its inhibitor ICAD.** *Nature* 1998, **391**(6662):43-50.
 26. Sakahira H, Enari M, Nagata S: **Cleavage of CAD inhibitor in CAD activation and DNA degradation during apoptosis.** *Nature* 1998, **391**(6662):96-99.
 27. Janicke RU, Sprengart ML, Wati MR, Porter AG: **Caspase-3 is required for DNA fragmentation and morphological changes associated with apoptosis.** *J Biol Chem* 1998, **273**(16):9357-9360.
 28. Mazhar D, Ang R, Waxman J: **COX inhibitors and breast cancer.** *Br J Cancer* 2006, **94**(3):346-350.
 29. Solomon SD, McMurray JJ, Pfeffer MA, Wittes J, Fowler R, Finn P, Anderson WF, Zauber A, Hawk E, Bertagnoli M: **Cardiovascular risk associated with celecoxib in a clinical trial for colorectal adenoma prevention.** *N Engl J Med* 2005, **352**(11):1071-1080.
 30. Jendrossek V: **Targeting apoptosis pathways by Celecoxib in cancer.** *Cancer Lett* 2011.
 31. Gore E, Bae K, Langer C, Extermann M, Movsas B, Okunieff P, Videtic G, Choy H: **Phase I/II trial of a COX-2 inhibitor with limited field radiation for intermediate prognosis patients who have locally advanced non-small-cell lung cancer: radiation therapy oncology group 0213.** *Clin Lung Cancer* 2011, **12**(2):125-130.
 32. De Ruysscher D, Bussink J, Rodrigus P, Kessels A, Dirx M, Houben R, Wanders R: **Concurrent celecoxib versus placebo in patients with stage II-III non-small cell lung cancer: a randomised phase II trial.** *Radiother Oncol* 2007, **84**(1):23-25.

doi:10.1186/1748-717X-6-176

Cite this article as: Lauber *et al*: Apoptosis induction and tumor cell repopulation: The yin and yang of radiotherapy. *Radiation Oncology* 2011 **6**:176.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

