

TWO DIFFERENT TYPES OF PHOTOCHEMISTRY IN PHYCOERYTHROCYANIN α -SUBUNIT*

QIANG HONG†, KAI-HONG ZHAO AND HUGO SCHEER‡

Botanisches Institut der Universität München, Menzinger Str. 67, D-80638 München 19, Germany

(Received 18 February 1993; accepted 13 July 1993)

Abstract—The photochemical activities of phycoerythrocyanin α -subunits from *Mastigocladus laminosus* separated by isoelectric focusing were tested by irradiating at 500, 550, 577 and 600 nm. Two types of photoreversible photochromic responses have been characterized by absorption and absorption difference spectroscopy. Type I is the well-known absorption shift from 571 to 506 nm. Type II is a new response characterized by a line-broadening of the 570 nm absorption.

INTRODUCTION

Phycoerythrocyanin (PEC)§ shows, unlike other phycobiliproteins,¹ a photoreversible photochromism reminiscent of that of the plant photoreceptor phytochrome, but with orange/green rather than red/far red action spectra.^{2–8} The photochemical activity is due to the phycoviolobin chromophore ($\lambda_{\max} \approx 570$ nm as isolated) located on the α -subunit.^{9,10} Upon irradiation with orange light, a new absorption arises at ≈ 500 nm (P506), which is converted back reversibly to the 570 nm form (P571) upon irradiation with green light. There are several lines of evidence indicating that this photochemical response involves, as in the case of phytochrome, a reversible Z/E-isomerization at the double bond between rings C and D of the linear tetrapyrrole.⁵

Phycoerythrocyanin is commonly considered an antenna pigment, located together with phycocyanin and allophycocyanin in the phycobilisomes of certain cyanobacteria.^{11–13} There is presently no obvious physiological function for its photochromic behavior, although an involvement of PEC in photomorphogenetic responses of cyanobacteria cannot be ruled out.^{14,15} Even so, it represents an interesting case with a transitory behavior between that of the highly fluorescent but photochemically inactive phycobiliproteins, and the photochemically active phytochrome.¹⁶

In previous investigations of PEC, large quantitative variations in the photochemistry of PEC and its α -subunit were noticed, which are (among others) related to the aggregation state² and to the status of SH-groups located on the same subunit.¹⁷ The results presented here show that there exist also qualitative variations in the photochemical signal, and that (at least) two different reactivities are involved in the reversible photochemistry of the PEC α -subunit.

MATERIALS AND METHODS

Phycoerythrocyanin and its subunits were isolated from *Mastigocladus laminosus* as before.^{2,12}

Absorption spectra were recorded in a lambda 2 spectrophotometer (Perkin-Elmer). Irradiations were done with a cold-light source (model 150, Volpi) equipped with fiber optics and suitable interference filters (10 nm bandwidth). The orange/green response (577 and 500 nm filters, respectively) was quantified in $\Delta\Delta A_1$ units. It corresponds (as the $\Delta\Delta A$ used before²) to the amplitude difference (maximum *minus* minimum) of the light-induced difference spectrum with extrema at ≈ 506 and ≈ 571 nm (Fig. 1A), divided by the maximum absorption of the 570 nm form. The type II photochemistry (see text) was defined similarly by $\Delta\Delta A_{II}$. It corresponds to the amplitude difference (maximum *minus* minimum) of the light-induced difference spectrum with extrema at ≈ 595 and ≈ 571 nm (Figs. 1B,2), divided by the maximum absorption of the 570 nm form.

RESULTS

Several separations of PEC under apparently the same conditions yielded preparations of the PEC α -subunit with rather different photochemical activities. The orange/red response (measured in $\Delta\Delta A_1$ units²) varied between 12% and 92%, and was generally on the high side. These variations are even more extreme than the reported values of 38%² and 50%⁴ and are mainly related to the status of SH-groups.¹⁷

The variations in photochemical activity of the α -PEC samples of different photochemistry were accompanied by a shift in the absorption maximum of the red-absorbing form (enriched after 500 nm irradiation) from 561 (lowest) to 566 nm (highest photochemistry). Published spectra also follow this correlation: Siebzehnriibl *et al.*² report $\lambda_{\max} = 562$ nm for a sample with $\Delta\Delta A_1 = 38\%$, and Kufer and Björn⁴ $\lambda_{\max} = 564$ nm for a sample with $\Delta\Delta A_1 = 50\%$. Furthermore, the low photochemistry sample showed a broader absorption and a less obvious shoulder after irradiation at 500 nm than the high photochemistry one (Figs. 1 and 2).

The difference spectra showed even more distinct changes, depending on the irradiation program. After subsequent irradiations at 500 and 570 nm, the difference spectra of high-photochemistry samples were always dominated by the well-known s-shaped type I signal (Fig. 1A). A rather different signal which has a w-shape typical for line broadening and a much smaller amplitude, was most clearly seen after subsequent irradiations at 577 and 600 nm (Fig. 1B). To distin-

*Dedicated to Prof. Dr. Wolfhart Rüdiger on the occasion of his 60th birthday.

†Present address: Center for Photomolecular Sciences, Imperial College, Prince Consort Road, London SW7 2BB, United Kingdom

‡To whom correspondence should be addressed.

§Abbreviations: PEC, phycoerythrocyanin; P_r and P_{fr}, phytochrome in the red and far-red absorbing form respectively; $\Delta\Delta A$, difference amplitude of the light-induced absorption differences (see Materials and Methods section).

guish it from the classical type I response, it shall be called type II response. For quantifying this response, we define the $\Delta\Delta A_{II}$ as for the type I response, but with the extrema at ≈ 570 and ≈ 595 as the reference wavelengths (see Materials and Methods).

The type II response was more pronounced in the low-photochemistry samples. Subsequent irradiations with 500 and 600 nm light resulted in a low amplitude ($\Delta\Delta A_I \approx 3.8\%$), somewhat distorted type I-signal ($\lambda_{max} \approx 500$ and 570 nm, Fig. 2). Subsequent irradiations at other wavelength pairs (577/600 or 500/577) resulted, in contrast, mainly in w-shaped absorption differences characteristic of the type II response (Fig. 2), with $\Delta\Delta A_{II}$ values $> 10\%$. Obviously, both signals are present in all samples, but the relatively small w-shaped one is prominent only when the "classical" photochemistry of PEC is low, either by internal factors¹⁷ or by external ones (*viz.* the irradiation program).

DISCUSSION

The photochemistry of PEC α -subunit was discovered by G.S. Björn³ during the search for cyanobacterial photochromic pigments (phycochromes) and has since been studied in considerable detail.^{2-8,17} Irrespective of quantitative variations,¹⁷ this phototransformation of the form absorbing at ≈ 570 nm (P571) to that absorbing at ≈ 500 nm (P506), is characterized by (i) a red shift of the long-wavelength absorption, (ii) a decrease in the near-UV absorption,¹⁶ (iii) a drop in fluorescence by a factor ≥ 5 ,⁵ (iv) a quantum yield of 10%,⁷ (v) an inversion of the circular dichroism signal,⁵ and (vi) only minor changes in the linear dichroism.⁸ Many of these features are similar to the ones observed during the phytochrome photoconversion from P_r to P_{fr} . Since the phytochrome conversion involves most likely a Z-E isomerization of the chromophore at the 15,16-double bond between rings C and D,¹⁶ a similar isomerization has been suggested to occur during this type I photochemistry in the α -84 phycoviolobin chromophore of PEC.^{5,7,17} The two distinct differences between the photochemistry of both pigments are also compatible with this interpretation: The absorption of the Z-isomer is bathochromically shifted in phytochrome and hypsochromically shifted in α -PEC, but it has been realized (albeit not understood on a molecular basis) for a long time that the bathochromic shift in phytochrome is unusual, whereas the hypsochromic shift is common for 15Z-15E isomerizations in bile pigments.¹⁸ The other difference is the small (apparent) reorientation of the chromophore in PEC, as determined by linear dichroism of the immobilized biliproteins.⁸ However, the change in orientation of the transition moment depends on many factors including the rigidity of the apoprotein, which can be very different for the two pigments, and the chromophore structure, whose π -system is considerably shorter in PEC than in phytochrome.

The molecular origin of the type II photochemistry described in this communication is presently unknown. Currently, we are investigating three possibilities: The first one is a Z/E-isomerization at the methine bridge between rings B and C. Although such isomers are unstable at ambient temperatures in solution,^{18,19} they might be stabilized by the protein. The second possibility is a Z/E-isomerization at the methine bridge between rings C and D, *e.g.* the same process




Figure 1. A) Type I photochemistry of PEC α -subunit ("high-photochemistry sample", $\Delta\Delta A_I = 92\%$). Absorption spectra after saturating preirradiations with 500 and 577 nm light (see label). The upper curve is the difference spectrum on the same scale (577 nm irradiated minus 500 nm irradiated). B) Type II photochemistry of PEC α -subunit ("high-photochemistry sample"): Absorption difference spectra of three different isolates, which were first irradiated with 577 nm light, then with 600 nm light, and the spectra subtracted in that order. $\Delta\Delta A_{II}$ of the type I ($\Delta\Delta A_{II}$ of the type II) responses of the samples were a: 89 (9.1), b: 91 (5.3) and c: 49 (7.8%).

as in the type I reaction. However, due to a modification in the protein,¹⁷ there may be a different interaction of the E-isomer with the environment. Since the spectral features of biliproteins are mainly determined by such interactions,^{16,19-22} this can lead to rather different spectra. An example for such interactions is P_{fr} , which as an E-isomer absorbs at longer wavelengths than the Z-isomer.²⁶ A third, more exotic possibility would be the formation of another form of the Z-isomer by a process reminiscent of photophysical hole-burning. The Z-E photoisomerization requires a considerable reaction volume and probably a certain amount of flexibility in the protein,^{16,19,20} and the quantitative variations of the type I photochemical signal have been related to changes in the rigidity of the apoprotein. The data presented here indicate that the type II photochemistry is not decreased in parallel, but rather remains unchanged or is even increased,¹⁷ thus indicating that the rigidity of the protein may play a




Figure 2. Photochemistry of PEC α -subunit ("low-photochemistry sample"). A) Absorption spectra after saturating pre-irradiations with 500 and 577 nm light (see label). B) Absorption difference spectra for the following irradiation programs: a) 500 minus 577 nm, b) 500 minus 600 nm, c) 577 minus 600 nm. Relative to the absorption spectra (A), the y-axis is expanded five fold.

lesser role. Line broadening is typical for photophysical hole-burning, a process which is generally observed at low temperatures only.²³ However, a protein has some structural features which might allow similar processes at ambient temperatures by locking the chromophores in local minima of the conformational space.

Acknowledgements—This work was supported by the Deutsche Forschungsgemeinschaft, Bonn (SFB 143, project A 1). K.-H.Z. acknowledges a postdoctoral stipend from the Alexander-von-Humboldt foundation, Bonn.

REFERENCES

- MacColl, R. and D. Guard-Friar (1987) *Phycobiliproteins*. CRC Press, Boca Raton, FL.
- Siebzehnriibl, S., R. Fischer, W. Kufer and H. Scheer (1989) Photochemistry of photobiliproteins: reciprocity of reversible photochemistry and aggregation in phycoerythrocyanin from *Mastigocladus laminosus*. *Photochem. Photobiol.* **49**, 753–761.
- Björn, G. S. (1979) Action spectra for *in vivo* and *in vitro* conversions of phycochrome b, a reversibly photochromic pigment in a blue-green alga, and its separation from other pigments. *Physiol. Plant* **46**, 281–286.
- Kufer, W. and G. S. Björn (1989) Photochromism of the cyanobacterial light harvesting biliprotein phycoerythrocyanin. *Physiol. Plant* **75**, 389–394.
- Maruthi Sai, P. S., S. Siebzehnriibl, S. Mahajan and H. Scheer (1992) Phycoerythrocyanins from *Westiellopsis prolifica* and *Nostoc rivulare*: Characterization of the phycoviolobin chromophore in both states. *Photochem. Photobiol.* **55**, 119–124.
- Kufer, W. (1988) Concerning the relationship of light harvesting biliproteins to phycochromes in cyanobacteria. In *Photosynthetic Light Harvesting Systems* (Edited by H. Scheer and S. Schneider), pp. 89–93. W. De Gruyter, Berlin.
- Siebzehnriibl, S. (1990) *Chromophorzuordnung und reversible Photochemie von C-Phycocyaninen und Phycoerythrocyaninen*, Dissertation, Universität München.
- Björn, G. S., N. Ekelund and L. O. Björn (1984) Light-induced linear dichroism in photoreversibly photochromic sensor pigments. 4. Lack of chromophore rotation in phycochrome-B immobilized *in vitro*. *Physiol. Plant* **60**, 253–256.
- Bishop, J. E., H. Rapoport, A. V. Klotz, C. F. Chan, A. N. Glazer, P. Füglistaller and H. Zuber (1987) Chromopeptides from phycoerythrocyanin. Structure and linkage of the three bilin groups. *J. Am. Chem. Soc.* **109**, 875–881.
- Düring, M., R. Huber, W. Bode, R. Rumbeli and H. Zuber (1990) Refined three-dimensional structure of phycoerythrocyanin from the cyanobacterium *Mastigocladus laminosus* at 2.7 Å. *J. Mol. Biol.* **211**, 633–644.
- Bryant, D. A. (1982) Phycoerythrocyanin and phycoerythrin—properties and occurrence in cyanobacteria. *J. Gen. Microbiol.* **128**, 835–844.
- Füglistaller, P., F. Suter and H. Zuber (1983) The complete amino-acid-sequence of both subunits of phycoerythrocyanin from the thermophilic cyanobacterium *Mastigocladus laminosus*. Hoppe Seyler's *Z. Physiol. Chem.* **364**, 691–712.
- Gantt, E. (1986) Phycobilisomes. In *Photosynthesis III—Photosynthetic Membranes and Light Harvesting Systems* (Edited by L. A. Staehelin and C. J. Arntzen), pp. 260–268. Springer, Berlin.
- Braune, W., T. Wilczok and R. Waclawek (1988) Indications for photoreversible reactions in the range of Phycochrome b absorption obtained by automated microscopic image analysis of germinating *Anabaena* akinetes. *Cytobios.* **54**, 39–48.
- Björn, L. O. (1979) Photoreversibly photochromic pigments in organism: properties and role in physiological light perception. *Quart. Rev. Biophys.* **12**, 1–25.
- Rüdiger, W. and F. Thümmel (1991) Phytochrome, the visual pigment of plants. *Angew. Chem. Int. Ed. Engl.* **30**, 1216–1228.
- Zhao, K.-H., Q. Hong, S. Siebzehnriibl and H. Scheer (1993) Phycoerythrocyanin: A pigment with two faces. In *Frontiers of Photobiology* (Edited by A. Shima, M. Ichihashi, Y. Fujiwara and H. Takebe), pp. 31–36. Elsevier, Amsterdam.
- Falk, H. (1989) *The Chemistry of Linear Oligopyrroles and Bile Pigments*. Wien, Springer, New York.
- Braslavsky, S. E., A. R. Holzwarth and K. Schaffner (1983) Konformationsanalyse, Photochemie und Photochemie der Gallenpigmente; Bilirubin- und Biliverdinmethylester und verwandte lineare Tetrapyrrole. *Angew. Chem.* **91**, 670–689; *Angew. Chem. Int. Ed. Engl.* **22**, 656–674.
- Scheer, H. (1981) Biliproteine. *Angew. Chem.* **93**, 230–250; *Angew. Chem. Int. Ed. Engl.* **20**, 241–261.
- Scheer, H. (1982) Light reaction path of photosynthesis. In *Phycobiliproteins: Molecular Aspects of Photosynthetic Antenna Systems* (Edited by F. K. Fong), pp. 7–45. Springer, Berlin.
- Huber, R., M. Schneider, O. Epp, I. Mayr, A. Messerschmidt, J. Pflugrath and H. Kayser (1987) Crystallization, crystal structure analysis and preliminary molecular model of the bilin binding protein from the insect *Pieris brassicae*. *J. Mol. Biol.* **195**, 423–434.
- Friedrich, J. and D. Haarer (1984) Photochemical hole-burning and optical relaxation spectroscopy in polymers and glasses. *Angew. Chem.* **96**, 96–123; *Angew. Chem. Int. Ed. Engl.* **23**, 113–140.