

Invited Trends Article

- Genetic Engineering of Disease and Pest Resistance in Plants: Present State of the Art
G. STRITTMATTER and D. WEGENER 673

Original Communications

- Epicuticular Leaf Waxes of the Hop (*Humulus lupulus*). Chemical Composition and Surface Structures
P.-G. GÜLZ, E. MÜLLER, T. HERRMANN, and P. LÖSEL 689
- Comparative FTIR and ¹³C CP/MAS NMR Spectroscopic Investigations on Sporopollenin of Different Systematic Origins
S. WILMESMEIER, ST. STEUERNAGEL, and R. WIERMANN 697
- Quinolizidine Alkaloid Profiles of South American Lupins: *Lupinus linearis* and the *Lupinus gibertianus* Complex
A. PLANCHUELO-RAVELO, L. WITTE, and M. WINK 702
- Fructose 1,6-Bisphosphatase in *Chlorella kessleri* Grown in Red or Blue Light
N. GROTJOHANN 707
- 3 α -Hydroxysteroid-5 β -oxidoreductase in Tissue Cultures of *Digitalis lanata*
U. STUHLEMMER, W. HAUSSMANN, F. MILEK, W. KREIS, and E. REINHARD 713
- Effect of Different Condensed Tannins on *Trichoplusia ni* Performance
A. GONZÁLEZ-COLOMA, CH. S. WISDOM, and PH. W. RUNDEL 722
- 6 β -Hydroxylation of 17 α -Acetoxycortexone by *Flavobacterium*
G. SPASSOV and W.-R. ABRAHAM 727
- Study of Canadian Propolis by GC-MS and HPLC
C. GARCÍA-VIGUERA, F. FERRERES, and F. A. TOMÁS-BARBERÁN 731

- Glandular Trichomes and the Volatiles Obtained by Steam Distillation of *Quercus robur* Leaves
R. ENGEL, P.-G. GÜLZ, TH. HERRMANN, and A. NAHRSTEDT 736

- The Unexpected Reduction of the Vinyl Group of Chlorophyll *b* by Sodium Borohydride in Methanolic Extracts of Maize Leaves and Its Inhibition by 8-Hydroxyquinoline
R. J. PORRA, W. SCHÄFER, E. CMIEL, I. KATHEIDER, and H. SCHEER 745

- Metabolic Reduction of Phenylpropanoid Compounds in Primary Leaves of Rye (*Secale cereale* L.) Leads to Increased UV-B Sensitivity of Photosynthesis
S. REUBER, J. LEITSCH, G. H. KRAUSE, and G. WEISSENBÖCK 749

- Naturally Occurring Antidotes against Benzimidazole Fungicides
S. TAHARA, Y. MATSUKURA, H. KATSUTA, and J. MIZUTANI 757

- Oxidative Stress of Crops Monitored by EPR
H. B. STEGMANN, P. SCHULER, ST. WESTPHAL, and E. WAGNER 766

- Molecular Modelling of the Interaction of Cyanoacrylate Inhibitors with Photosystem II. Part 1. The Effect of Hydrophobicity of Inhibitor Binding
S. P. MACKAY and P. J. O'MALLEY 773

- Molecular Modelling of the Interaction of Cyanoacrylate Inhibitors with Photosystem II. Part 2. The Effect of Stereochemistry of Inhibitor Binding
S. P. MACKAY and P. J. O'MALLEY 782

- Photoregulation of Nitrogen Metabolism and Protein Accumulation in the Red Alga *Corallina elongata* Ellis et Soland
F. L. FIGUEROA 788

- Translocation of Adenine Nucleotides in the Mitochondria of Male Sterile and Male Fertile Sorghum
J. ARORA, P. NATH, and P. V. SANE 795

Continued overleaf

Complete Sequence of One Copy of the *psbA* Gene from the Thermophilic Cyanobacterium *Synechococcus elongatus*

R. KLOOS, E. STEVENS, and W. OETTMEIER 799

Linear Free Energy Relationships for N(7)-Substituted Guanosines as Substrates of Calf Spleen Purine Nucleoside Phosphorylase. Possible Role of N(7)-Protonation as an Intermediary in Phosphorolysis

A. BZOWSKA, E. KULIKOWSKA, and D. SHUGAR 803

Notes

Isolation and Properties of Trimethylamine N-Oxide/Dimethylsulfoxide Reductase from the Purple Bacterium *Rhodospirillum rubrum*

P. SAJITZ, J.-H. KLEMME, H.-G. KOCH, and M. MOLITOR 812

Inhibition of Dihydrofolate Reductase by Mofebutazon

E. STREHL, I. SCHNEIDER, and E. F. ELSTNER 815

Antitumoral Effect of Bleomycin + Dolomite Combination Treatment, in Mice Bearing Ehrlich Ascites Carcinoma

S. SCHELLER, W. KROL, K. SKIRMUNTT, G. ZYDOWICZ, and J. SHANI 818

The Unexpected Reduction of the Vinyl Group of Chlorophyll *b* by Sodium Borohydride in Methanolic Extracts of Maize Leaves and Its Inhibition by 8-Hydroxyquinoline

R. J. Porra^{a,*}, W. Schäfer^b, E. Cmiel^c, Ingrid Katheder^a, and H. Scheer^a

^a Botanisches Institut der Universität, D-80638 München

^b Max-Planck-Institut für Biochemie, D-82152 Martinsried

^c Institut für Physikalische Chemie, Technische Universität München, D-85748 Garching

Z. Naturforsch. **48c**, 745–748 (1993); received May 14/July 9, 1993


Photosynthesis, Chlorophyll *b*, Reductive Extraction, Borohydride, 8-Hydroxyquinoline

During rapid extraction of chlorophylls from maize leaves under reducing conditions with methanol containing NaBH₄, chlorophyll *a* remained unchanged but chlorophyll *b* yielded [7-hydroxymethyl]-chlorophyll *b*. The 3-vinyl group of chlorophyll *b* was also reduced forming significant amounts, up to 60%, of [3-ethyl]-[7-hydroxymethyl]-chlorophyll *b*. This was unexpected since this reduction of the 3-vinyl group does not occur when isolated chlorophyll *b* is treated in an identical manner with methanolic borohydride. The vinyl-group of chlorophyll *a* is not reduced during the same extraction conditions suggesting that the presence of a formyl or hydroxyethyl group at C-7 is necessary. The presence of 8-hydroxyquinoline and NaBH₄ in equimolar (16.5 mM) concentrations strongly inhibits the reduction of the 3-vinyl group of chlorophyll *b* in leaf extracts.

Introduction

The reduction of pure Chl *a* (Fig. 1; structure **I**) and Chl *b* (**II**), and of bacteriochlorophylls by NaBH₄ has been extensively studied [1–6]. In the plant chlorophylls, the 7-formyl group of Chl *b* is rapidly reduced within seconds to form [7-hydroxymethyl]-Chl *b* (**III**): the reduction of the 13¹-oxo group of both Chls *a* and *b* is much slower taking several hours at room temperature for completion [6]. No concomitant reduction of conjugated double bonds, a well known side-reaction of borohydrides [7], has been reported so far. We have now found that as much as 60% of the Chl *b* was reduced to [3-ethyl]-[7-hydroxymethyl]-Chl *b* (**IV**) during extraction of the newly-formed chlorophylls in greening maize seedlings with methanol containing 16.5 mM NaBH₄: this reductant was employed to convert the formyl group to a hydroxymethyl group in biosynthetic ¹⁸O-incorpora-

tion studies which showed that the precursor of the formyl-group oxygen of Chl *b* is molecular oxygen [8].



Abbreviations: Chl, chlorophyll; DEAE-cellulose, diethylaminoethyl-cellulose; EDTA, ethylenediamine-tetra-acetic acid; NMR, nuclear magnetic resonance.

* **Permanent address:** CSIRO-Division of Plant Industry, GPO Box 1600, Canberra ACT 2601, Australia.

Reprint requests to Hugo Scheer, Botanisches Institut, Universität München, Menzinger Straße 67, D-80638 München, Germany.

Verlag der Zeitschrift für Naturforschung,
D-72072 Tübingen
0939-5075/93/0900-0745 \$ 01.30/0

Fig. 1. The structures of chlorophyll derivatives referred to in this paper. The IUB-IUPAC approved numbering system for tetrapyrroles [14] has been used with bracket [] nomenclature for substitutions. [7-hydroxymethyl]-Chl *b* (**III**) is identical to [7-hydroxymethyl]-Chl *a* but will be referred to here as a Chl *b* derivative to indicate its origin.

Experimental

Chemicals

NaBH_4 and 8-hydroxyquinoline (technical grade) were obtained from Merck-Schuchardt, Darmstadt, Germany. NaB^2H_4 was obtained from Cambridge Isotope Laboratories, Cambridge, MA., U.S.A. Solvents and other chemicals were analytical reagent grade or purified by standard techniques. DEAE-cellulose (DE 52), supplied by Whatman Laboratory Division, Maidstone, England, was prepared as a methanolic suspension [9] which was then equilibrated with CHCl_3 . Pure Chls *a* and *b* were prepared from green maize leaves as previously described [10].

Organisms, growth and greening conditions

Etiolated maize seedlings (*Zea mays* hybrid var. Dekalb XL689) were grown in the dark at 18 °C for 18 days [11]. Etiolated leaves, excised from these seedlings, were then placed in H_2O and greened by illumination ($50\text{--}60 \mu\text{E} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$) with white light (Philips TLD 18 W/84 tubes) for up to 26 h at 27 °C [12].

Extraction of chlorophylls from greened maize leaves

Leaves were finely chopped with scissors into a mortar and extracted by grinding with a pestle to a translucent pulp in a freshly prepared solution (14 ml/g fresh wt. of leaves) of NaBH_4 (16.5 mM) in methanol. Where specified in the text, the extraction was carried out with methanol containing either 16.5 mM NaB^2H_4 or equimolar (16.5 mM) concentrations of NaBH_4 and 8-hydroxyquinoline. Grinding was completed in three minutes when an excess of glucose (150 mg) was added to remove unspent NaBH_4 . The clear methanolic supernatant contained Chl *a* and reduced derivatives of Chl *b*.

Column chromatography of extracted chlorophyll derivatives

The chlorophylls in the methanolic supernatant were transferred to diethylether by adding large volumes of saturated brine. The ether solution was then washed with a further large volume of saturated brine, dried over solid NaCl and evaporated to dryness at approximately 40 °C under vacuum

in a rotary evaporator. The dry chlorophylls were redissolved in a minimum quantity of CHCl_3 and applied to a 4×65 mm high column of DEAE-cellulose equilibrated with CHCl_3 (see above). The carotenoids and chlorophylls were eluted with CHCl_3 or CHCl_3 containing either 2 or 10% methanol as described in the Results section. The elution of the chlorophylls was monitored by adsorption spectroscopy of each ml of chlorophyll-containing eluant between 680–630 nm and 480–400 nm.

Spectroscopy

All absorption spectroscopy was performed in quartz cuvettes (1 ml capacity and 1 cm light path) using a Shimadzu UV 1202 Spectrophotometer. Mass spectra were obtained by fast atom bombardment ionization (*m*-nitro-benzylic alcohol matrix) with a MAT 900 Mass Spectrometer (Finnigan MAT, Bremen, Germany). ^1H NMR spectra were recorded in $^2\text{H}_5$ -pyridine with a model AM 360 MHz instrument (Bruker, Karlsruhe, Germany).

Results and Discussion

The chlorophylls present in methanolic- NaBH_4 extracts of greened dark-grown maize leaves as determined by chromatography and absorption spectroscopy

When the pigments extracted in methanol containing 16.5 mM NaBH_4 were applied to a DEAE-cellulose column and developed with CHCl_3 , a yellow carotenoid fraction ($\lambda_{\text{max}} = 481, 454$ and 426 nm) was followed by three blue chlorophyll bands. The first contained Chl *a* (**I**) ($\lambda_{\text{max}} = 666$ nm). The second, split band contained two pigments with similar absorption spectra (Fig. 2). The first one ($\lambda_{\text{max}} = 659$ and 428 nm, spectrum 2) contained an unknown pigment, the second half ($\lambda_{\text{max}} = 659$ and 434 nm, spectrum 3), eluting only slowly with CHCl_3 , and better with 2% methanol added, contained the expected C-7¹ reduction product of Chl *b*, viz. [7-hydroxymethyl]-Chl *b* (**III**) [cf. 6]. The slowest-moving third blue band ($\lambda_{\text{max}} = 653$ and 415 nm), which eluted with CHCl_3 containing 10% methanol, contained [7-hydroxymethyl]-[13¹-hydroxy]-Chl *b* (**V**), which is additionally reduced at the 13¹-oxo group [cf. 6].




Fig. 2. UV-Vis absorption spectra of Chl *b* and its reduction products in chloroform. Chl *b* (**II**) (---, spectrum 1); [3-ethyl]-[7-hydroxymethyl]-Chl *b* (**IV**) (—, spectrum 2); and [7-hydroxymethyl]-Chl *b* (**III**) (-·-·-, spectrum 3). Peak positions of the main bands (nm) are indicated.

Consistent with previous studies [2–4, 6], we showed that pure Chl *b* (**II**), when treated rapidly with NaBH₄ in methanol, was all reduced to [7-hydroxymethyl]-Chl *b* (**III**). We speculated that the additional band in extracts of leaves ($\lambda_{\max} = 659$ and 428 nm) was [3-ethyl]-[7-hydroxymethyl]-Chl *b* (**IV**) in which both the 3-vinyl and 7-formyl groups were reduced: both the more rapid elution of the 428 nm-absorbing material from the column and the shift of the Soret peak by 6 nm to shorter wavelengths relative to that of [7-hydroxymethyl]-Chl *b* (**III**) are consistent with the reduction of a vinyl group. Assuming equal absorption coefficients of **III** and **IV** at 659 nm, the latter comprised up to 60% of reduction products.

Product identification

The ¹H NMR spectrum of [7-hydroxymethyl]-Chl *b* (**III**) agreed with the structure. The 3-vinyl signals occur at 8.06 (H_x), 6.38 (H_A) and 6.06 ppm (H_B). The OH signal, which is somewhat variable and known to be very solvent dependent, occurs at approximately 7.4 ppm and was identified by ¹H/²H exchange with ²H₂O. The mass spectrum showed both M⁺ (base at 906 *m/z*) and (M + H)⁺ ions (base at 907 *m/z*) and the corresponding isotope peaks in an approximately 4:1 ratio. Upon reductive extraction with NaB²H₄ in methanol,

there was an increase in mass by 1 *m/z*, and a decrease in the intensity of the 7-CH₂ signal at 6.08 ppm by 50% in the ¹H NMR spectrum.

In the ¹H NMR spectrum of [3-ethyl]-[7-hydroxymethyl]-Chl *b* (**IV**), the ring current was slightly reduced, and the vinyl signals were no longer present. The mass spectrum showed again M⁺ (base at 908 *m/z*) and (M + H)⁺ ions (base at 909 *m/z*) in an approximately 4:1 ratio. Reductive extraction with NaB²H₄ in methanol increased the mass by 2 *m/z* indicating the incorporation of one ²H atom into the hydroxymethyl group and also the 3-ethyl group.

Investigation of the reduction of the vinyl group of Chl *b* in methanolic-borohydride extracts of maize leaves

Consideration of which components of the leaf extract initiated reduction of the vinyl group of Chl *b* led to investigation of the effect of 10, 20 and 40% water on the reaction of pure Chl *b* with methanolic NaBH₄. It was always completely reduced to the [7-hydroxymethyl]-Chl *b* (**III**) with Soret absorption at 434 nm: no [3-ethyl]-[7-hydroxymethyl]-Chl *b* (**IV**), the “3,8-diethyl” derivative with Soret absorption at 428 nm was detected.

To investigate the possibility that metal ions in the cell sap of the maize leaves are involved in the vinyl group reduction, the effect of three metal chelating agents on ethyl group formation during chlorophyll extraction with methanolic NaBH₄ were investigated: 8-hydroxyquinoline, 8-hydroxyquinoline-5-sulphonic acid (sodium salt) or EDTA (tetra-sodium salt) were added in equimolar concentrations with the NaBH₄. The two 8-hydroxyquinoline derivatives, but not the EDTA, strongly inhibited the formation of the 428 nm-absorbing “3,8-diethyl” derivative (**IV**). However, when the concentration of the NaBH₄ exceeded that of the chelating agent, the “3,8-diethyl” compound was again formed. Further, metal ions such as Fe²⁺, Mg²⁺, Mn²⁺, Co²⁺, Cu²⁺ and Ca²⁺ did not induce the reduction of the vinyl group of pure Chl *b*. We concluded, therefore, that the inhibition is not due to chelation of an activating metal ion; rather, the 8-hydroxyquinoline derivatives, in equimolar proportions with NaBH₄, were complexing the reductant and modifying its reducing properties. Indeed the chelator prevented also reduction of the

13^1 -oxo group. In the presence of equimolar chelator, the NaBH_4 formed a third slow-moving chlorophyll band but it no longer had principal peaks of the doubly-reduced Chl *b* (V) (see above) but absorbed at 652.5, 612.5 and 431.5 nm with a shoulder at 411 nm and moved considerably faster during chromatography.

Concluding remarks

The presence of a formyl- or hydroxymethyl-group at C-7 appears to be needed for the reduction of the nearby 3-vinyl group as no evidence could be found for the reduction of the 3-vinyl group of Chl *a* during extraction. Since the 7-formyl and the 3-vinyl substituents of Chl *b* are close to each other and conjugated, a concerted mechanism is feasible. However, the identity of the agents in these extracts which catalyze the reduction are unknown.

Because the above ^1H NMR investigations have shown that ^2H from NaB^2H_4 was incorporated not

only into the 7-hydroxymethyl group but also into the 3-ethyl group, it is clear that the majority (if not all) of the "3,8-diethyl" derivative arose by borohydride reduction during extraction and was not originally present in the leaves. This fact, coupled with the failure to find any "3,8-diethyl" derivative of Chl *a* in extracts, supports the view of Rebeiz *et al.* [13] that the natural heterogeneity of Chls *a* and *b* in nature, which includes the so-called "3-mono-vinyl-" and "3,8-di-vinyl-" derivatives, does not extend to "3,8-diethyl" forms.

Acknowledgements

One of us (RJP) thanks the Deutscher Akademischer Austauschdienst (DAAD) for financial support during his visit to the Botanisches Institut der Universität München to complete this work. This work was also in part supported by the Deutsche Forschungsgemeinschaft (SFB 143, Projekt A9) and the CSIRO-Division of Plant Industry, Australia.

- [1] S. L. Ditson, R. C. Davis, and R. M. Pearlstein, *Biochim. Biophys. Acta* **766**, 623–629 (1984).
- [2] A. S. Holt, *Plant Physiol.* **34**, 310–314 (1959).
- [3] H. Wolf and H. Scheer, *Tetrahedron Lett.* **28**, 5839–5867 (1972).
- [4] P. H. Hynninen, *J. Chromatogr.* **175**, 89–104 (1979).
- [5] R. J. Porra, *Eur. J. Biochem.* **156**, 111–121 (1986).
- [6] H. Scheer, R. J. Porra, and J. M. Anderson, *Photochem. Photobiol.* **50**, 403–412 (1989).
- [7] C. F. Lane, *Synthesis* **1975**, 135–146.
- [8] R. J. Porra, W. Schäfer, E. Cmiel, I. Katheder, and H. Scheer, *FEBS Lett.* **323**, 31–34 (1993).
- [9] N. Sato and N. Murata, *Biochim. Biophys. Acta* **501**, 103–111 (1978).
- [10] R. J. Porra, *Biochim. Biophys. Acta* **1015**, 493–502 (1990).
- [11] O. Klein and R. J. Porra, *Hoppe-Seyler's Z. Physiol. Chem.* **363**, 551–562 (1982).
- [12] R. J. Porra, O. Klein, and P. E. Wright, *Eur. J. Biochem.* **130**, 509–516 (1983).
- [13] C. A. Rebeiz, S. M. Wu, M. Kuhadja, H. Daniell, and E. J. Perkins, *Mol. Cell Biochem.* **57**, 97–125 (1983).
- [14] J. E. Merritt and K. L. Loening, *Pure Appl. Chem.* **51**, 2251–2304 (1979).