

Localization of the genes for tumor necrosis factor and lymphotoxin between the *HLA* class I and III regions by field inversion gel electrophoresis

J. Ragoussis¹, K. Bloemer², E. H. Weiss², and A. Ziegler¹

¹ Medizinische Klinik und Medizinisch-Naturwissenschaftliches Forschungszentrum, University of Tübingen, Ob dem Himmelreich 7, Federal Republic of Germany

² Institute of Immunology, University of Munich, Goethestrasse 31, D-8000 Munich 2, Federal Republic of Germany

The human major histocompatibility (*HLA*) complex is located on the short arm of chromosome 6 in the *6p21.31-6p21.33* region (Spring et al. 1985, Ziegler et al. 1985a). The physical length of the entire *HLA* complex is unknown so far, but our estimate based on the separation of DNA fragments containing *HLA* genes by pulsed-field gel electrophoresis (Schwartz and Cantor 1984, Carle and Olson 1984) indicates that it encompasses at least 2500 kb pairs (Ragoussis et al. 1986). This estimate has recently been confirmed (Lawrance et al. 1987). Apart from the highly polymorphic class I and II loci, the *HLA* complex contains the genes for several complement components and 21-steroid hydroxylase (class III region) (Lamm and Olaisen 1985). In addition, the loci for tumor necrosis factor (*TNFA*) as well as lymphotoxin (*TNFB*) are also near or even within the *HLA* region (Nedwin et al. 1985). Spies and co-workers (1986) demonstrated that the *TNF* genes map either centromeric to *HLA-DP* or telomeric to the class II region, although apparently not in the vicinity of any known class I or III genes. The recent demonstration that the *TNF* loci are situated 70 kb upstream of the *H-2D* gene in the BALB/c mouse between the class III and class I regions (Müller et al. 1987a, b) suggested an analogous location in man, because the genetic organization of the major histocompatibility complexes (MHC) of both species is very similar. To clarify the position of the *TNFA* and *TNFB* genes on the *HLA* map, we have assigned *TNFA* to large DNA restriction fragments separated by field inversion gel electrophoresis (FIGE) (Carle et al. 1986), which hybridize with either class III- or class I-specific probes as well. These results prove that the *TNFA* locus is localized between the *HLA* class III region and the *HLA-B* locus.

To avoid interpretative difficulties which might arise from haplotype-specific restriction fragment length polymorphisms, mutant human cell lines with monosomy 6 or *HLA* hemizyosity were employed. All mutants were derived from BJAB-B95.8.6 lymphoma cells with the *HLA* haplotypes *A1, Cw4, B35* and *A2, C-, B13* (Spring et

al. 1985). Mutant BM 19.7 is a monosomy 6 mutant cell line retaining the *A2* haplotype (Ziegler et al. 1985b). BM 28.7 also exhibits monosomy 6, but with loss of the chromosome bearing the *A2* haplotype (Ragoussis et al. 1986). In the interstitial deletion mutant BM 2.2.3, the class I region of the *A2* haplotype has been deleted, but the class II and III regions of this haplotype are still present, as is the chromosome 6 carrying the *A1* haplotype (Ziegler et al. 1985a).

Large genomic DNA fragments were generated with various restriction enzymes, separated by FIGE, and analyzed by hybridization to Southern blots (Southern 1975) under stringent conditions. In the *Bss* HII digests (Fig. 1), a 370 kb DNA fragment was detected which hybridized both to the *HLA-B* locus-specific probe that had been derived 29 kb 5' of the *HLA-B* gene (Fig. 1b) and to the *TNFA* probe (Fig. 1c), indicating linkage of *TNFA* with the *HLA-B* locus. The class III loci were present on another *Bss* HII fragment 125 kb in length (Fig. 1d). The 370 kb *Bss* HII fragment did not contain any class I or class II genes (Fig. 1e and not shown).

Hybridizations with the same probes to blots of *Nru* I-digested DNA (Fig. 2) gave evidence for the linkage of *TNFA* to the class III genes. Two *Nru* I fragments of 710 and 690 kb were detected which contained the class III genes (Fig. 2a and not shown) and *TNFA* (Fig. 2b). It is likely that partial digestion of an *Nru* I site is responsible for the generation of two fragments of similar size carrying the class III and *TNFA* genes. The class I region probes hybridized to a distinct fragment of about 800 kb (Fig. 2c). An even larger *Nru* I fragment of more than 1000 kb carried class II sequences (not shown).

To prove linkage, double digests with *Nru* I and *Bss* HII were performed as well (Fig. 3). The hybridization pattern obtained with the class III probes showed that the 125 kb *Bss* HII fragment carrying the class III region is contained within the *Nru* I fragment (Fig. 3a). However, the *Bss* HII fragment hybridizing with the *TNF* probe is reduced to a length of only about 50–90 kb (Fig. 3b). Con-




Fig. 1a-e. Southern blot hybridization of human DNA digested with *Bss* HII. Low melting point (LMP)-agarose blocks containing cells from the cell lines BM19.7, BJAB-B95.8.6, BM28.7, and BM2.2.3 (10^6 cells per block) were prepared according to the method of van der Bliek and co-workers (1986), slightly modified for FIGE: 2×10^7 cultured cells were pelleted and resuspended in PBS (10 mM sodium phosphate, 150 mM sodium chloride, pH 8), pelleted again, and resuspended in 1 ml PBS. One percent of LMP agarose (BRL ultra pure) in PBS was melted, equilibrated at 37 °C, and mixed well with an equal volume of cell suspension. The agarose/cell mixture was poured into a series of slots in a perpex mold. After solidification, the agarose blocks were deposited in a lysis mix [0.5 M ethylenediaminetetraacetate (EDTA), pH 9.5, 1% sodium lauryl sarcosine, 1 mg/ml proteinase K] at 1 ml per block and incubated for 48 h at 50 °C. The lysis mix was washed off overnight with TE buffer [10 mM Tris/HCl, pH 8, 10 mM EDTA, 0.1 mM phenylmethylsulfonylfluoride (PMSF)], 20 ml/block. Prior to digestion, the blocks were washed for 1 h in restriction enzyme buffer containing 0.1 mM PMSF (RE) (10 ml/block). One hundred microliters RE buffer was mixed with 25 units of enzyme in an Eppendorf tube; a 1/2 block was added and incubated for 6 h at 37 °C. The reaction was stopped with 1 ml TE containing 0.1% NaDoSO₄. Half a block was loaded in a 17 × 15 × 0.8 cm 1% agarose gel (Seakem GTG) in 0.25 × TBE (22.5 mM Tris borate, 0.5 mM EDTA, pH 8) and placed in an electrophoresis chamber. FIGE was carried out for 24 h at 150 V or 189 V constant, 15 °C. The voltage gradient was inverted periodically with a time ratio of 3 : 1 and the longer time increasing from 10 to 60 or 120 s. We used a computer-controlled power supply (CS 130, version 2.1) supplied by the EMBL, Heidelberg. After a run, the gel was stained in ethidium bromide (1 µg/ml) for 30 min, destained for 30 min in H₂O, and photographed under 302 nm UV light (a). The DNA was then blotted to Amersham Hybond membranes, baked for 2 h at 80 °C, and hybridized sequentially with various DNA probes. Prehybridization and hybridization solutions were as follows: 3 × SSC (1 × SSC is 0.15 M NaCl/0.015 M sodium citrate, pH 7.4) containing 5 × Denhardt's solution (0.1% Ficoll, 0.1% polyvinylpyrrolidone, 0.1% bovine serum albumin), 10% dextran sulfate, 0.5% NaDodSO₄, and 100 µg/ml salmon sperm DNA. Prehybridization was performed for 3 h at 65 °C and hybridization overnight at 65 °C with ³²P dCTP-labeled DNA probe (Feinberg and Vogelstein 1983) at 10⁶ cpm/ml. Washing was performed in four steps: 10 min in 2 × SSC, 0.1% NaDodSO₄ at room temperature; 30 min in 1 × SSC, 0.1% NaDodSO₄ at 65 °C; 30 min in 0.3 × SSC, 0.1% NaDodSO₄ at 65 °C and 30 min in 0.1 × SSC, 0.1% NaDodSO₄ at 65 °C. The filters were exposed for 2–7 days to Kodak XAR-5 film at –70 °C. Before rehybridization, the hybridized DNA probe was removed in bidistilled H₂O at 70 °C and the result checked by autoradiography. An *HLA-B* locus-specific probe, a 5.0 kb *Kpn* I/*Hind* III fragment, was isolated which is located 29 kb 5' of the *HLA-B* gene (Weiss et al. 1987). Another *B* locus-specific probe, a 2.2 kb *Pst* I/*Bam* HI fragment, was isolated 26 kb 3' of the *B* gene. Probes for C4 and C2 (class III region) were kindly provided by D. Campbell (Belt et al. 1984) and P. Schneider. The *TNFA* probe was a *Hind* III/*Sal* I fragment derived 20 kb 5' of the *TNFA* gene (Kioussis et al. 1987). As molecular size markers, the yeast strain way 5-4A prepared as described (Schwartz and Cantor 1984) and λ concatemers prepared as described (Van Ommen and Verkerk 1986) were used. The *HLA-B* 5' locus probe detected a 370 kb fragment (b) which was also recognized by the *TNF* probe (c) but not by either the C4 (d) or the C2 probe (not shown), which hybridize to a 125 kb *Bss* HII fragment, or by the *HLA-B* 3' probe, which hybridized to a 270 kb fragment (e)




Fig. 2a-c. Southern blot hybridization of human DNA digested with *Nru* I. DNA from the cell line BM28.7 was digested with *Nru* I. The conditions for FIGE were as in Figure 1, except that the voltage was increased to 189 V. Southern blotting and hybridization conditions were identical with those employed for *Bss* HII-digested DNA. The C2 probe detected two bands of slightly larger or smaller sizes than 700 kb (a) and these were also hybridizing to the *TNFA* probe (b). The *HLA-B5'* locus-specific probe detected another fragment of 800 kb (c) which contains also the *HLA-B* gene (not shown)

sequently, the *HLA-B* locus-specific probe was expected to hybridize with a different band of about 300 kb, and indeed, a new band of this approximate size (280 kb) was found (Fig. 3c). This result proves that the *Nru* I and *Bss* HII fragments overlap by 50–90 kb, and the *TNF* genes must be localized within this region (Fig. 4).

Since an *HLA-B* gene probe itself and a fragment, isolated 26 kb 3' of this gene, hybridized to a different *Bss* HII fragment (270 kb) (Fig. 1e), it is possible to give a rather precise estimate of the distance between *TNFA* and *HLA-B*. The *TNFA* locus and the closely linked




Fig. 3a-c. Southern blot hybridization of human DNA doubly digested with *Nru* I and *Bss* HII. DNA from the cell line BM19.7 was doubly digested with *Nru* I and *Bss* HII (left) or *Bss* HII alone (right). FIGE conditions were as in Figure 1 except that the longer time increased linearly from 10 to 120 s. Hybridization was carried out sequentially (as described in Fig. 1) with C4 (a), *TNFA* (b), and the *HLA-B5'* locus probe (c)

(Nedospasov et al. 1986) *TNFB* gene are therefore located about 300 kb centromeric to the *HLA-B* gene. Preliminary experiments employing double digests with *Nru* I and Not I indicate a distance of *TNF* to the class III region of not more than 300–400 kb. Since the *TNFA*/class III region-carrying *Nru* I fragment does not contain class II genes, the minimal distance between the class II loci and the class I region must be about 1000 kb.

These data are in agreement with the mapping of the *TNF* genes in the mouse, where they are located 70 kb proximal to the *H-2D* gene. However, it is very likely that the physical distance between *HLA-B* and *TNF* is considerably larger in the human genome than in the mouse.

Since the association of particular *HLA* class I and II alleles with a variety of diseases (Tiwari and Terasaki 1985) is still an enigma, the mapping of loci in the vicinity of these genes might provide further insight into the underlying mechanisms. Therefore the location of *TNFA* and *TNFB* by FIGE or related techniques in the *HLA* region may contribute to an understanding of *HLA*-disease associations. Since the region occupied by the *TNFA* and *TNFB* genes is only about 7 kb (Nedospasov et al. 1986),




Fig. 4. Physical map of the central part of the *HLA* complex based on separation of large DNA restriction fragments by FIGE. \dashv , the existence of additional *Nru* I fragments between the ones detected cannot be excluded

it is to be expected that several additional loci will be found between the class I and III regions. The preparative application of FIGE in conjunction with cloning techniques (Michiels et al. 1987) will help to bridge the long chromosomal distance between the different class I genes and the class III region.

Acknowledgments. This work was supported by the Deutsche Forschungsgemeinschaft through SFB 120 (project A4) and the Genzentrum München. We are grateful to Ms. Andrea Schenk for her valuable technical assistance and to Ms. Gudrun Szalay for her help in the preparation of the molecular size markers. We also thank Dr. Denise Barlow, Mr. Carol Stettner, and Mr. Alfons Riedinger (EMBL, Heidelberg) for help in setting up field inversion gel electrophoresis.

Note added in proof:

Improved separation conditions in FIGE suggest that the Nru I fragment containing class I genes may be even larger than 800 kb (Fig. 2c), possibly about 1000 kb.

References

- Belt, K. T., Carroll, M. C., and Porter, R. R.: Multiple forms of human complement component. *Cell* 36: 907-914, 1984
- Carle, G. F. and Olson, M. V.: Separation of chromosomal DNA molecules from yeast by orthogonal-field-alternation gel electrophoresis. *Nucleic Acids Res.* 12: 5647-5664, 1984
- Carle, G. F., Frank, M., and Olson, M. V.: Electrophoretic separations of large DNA molecules in one-dimensional electric fields that are periodically inverted. *Science* 232: 65-68, 1986
- Feinberg, A. P. and Vogelstein, B.: A technique for radiolabeling DNA restriction endonuclease fragments to high specific activity. *Anal. Biochem.* 132: 6-13, 1983
- Kioussis, D., Wilson, F., Daniels, C., Leveton, C., Taverne, J., and Playfair, H. L.: Expression and rescuing of a cloned human tumor necrosis factor gene using an EBV-based shuttle cosmid vector. *EMBO J.* 6: 355-361, 1987
- Lamm, L. U. and Olaisen, B.: Report of the Committee on the Genetic Constitution of Chromosomes 5 and 6. Eighth International Workshop on Human Gene Mapping. *Cytogenet. Cell Genet.* 40: 128-155, 1985
- Lawrance, S. K., Smith, C. L., Srivastava, R., Cantor, C. R., and Weissman, S. M.: Megabase-scale mapping of the HLA gene complex by pulsed-field gel electrophoresis. *Science* 235: 1387-1390, 1987
- Michiels, F., Burmeister, M., and Lehrach, H.: Derivation of clones close to *met* by preparative field-inversion gel electrophoresis. *Science* 236: 1305-1308, 1987
- Müller, U., Jongeneel, V., Nedospasov, S. A., Fischer Lindahl, K., and Steinmetz, M.: Tumor necrosis factor and lymphotoxin map close to H-2D in the mouse major histocompatibility complex. *Nature* 325: 265-267, 1987a
- Müller, U., Stephan, D., Philippsen, P., and Steinmetz, M.: Orientation and molecular map position of the complement genes in the mouse MHC. *EMBO J.* 6: 369-373, 1987b
- Nedospasov, S. A., Shakov, A. N., Turetskaya, R. L., Mett, V. A., Azizov, M. M., Georgiev, G. P., Korobko, V. G., Dobrynin, V. N., Filippov, S. A., Bystrov, E. F., Boldyreva, E. F., Chuvpilo, S. A., Chumakov, A. M., Shingarova, L. N., and Ovchinnikov, Y. A.: Tandem arrangement of genes coding for tumor necrosis factor (TNF- α) and lymphotoxin (TNF- β) in the human genome. *Cold Spring Harbor Symp. Quant. Biol.* 51: 611-624, 1986
- Nedwin, G. E., Naylor, S. L., Sakaguchi, A. Y., Smith, D., Jarrett-Nedwin, J., Pennica, D., Goeddel, D. V., and Gray, P. W.: Human lymphotoxin and tumor necrosis factor genes: structure, homology and chromosomal localisation. *Nucleic Acids Res.* 13: 6361-6373, 1985
- Ragoussis, J., van der Blik, A., Trowsdale, J., and Ziegler, A.: Mapping of HLA genes using pulsed field gradient electrophoresis. *FEBS Lett.* 204: 1-4, 1986
- Schwartz, D. C. and Cantor, C. R.: Separation of yeast chromosome-sized DNAs by pulsed field gradient gel electrophoresis. *Cell* 37: 67-75, 1984
- Southern, E. M.: Detection of specific sequences among DNA fragments separated by gel electrophoresis. *J. Mol. Biol.* 98: 503-517, 1975
- Spies, T., Morton, C. C., Nedospasov, S. A., Fiers, W., Pious, D., and Strominger, J. L.: Genes for the tumor necrosis factors α and β are linked to the human major histocompatibility complex. *Proc. Natl. Acad. Sci. U.S.A.* 83: 8699-8702, 1986
- Spring, B., Fonatsch, C., Müller, C., Pawelec, G., Kömpf, J., Wernet, P., and Ziegler, A.: Refinement of HLA gene mapping with induced B-cell line mutants. *Immunogenetics* 21: 277-291, 1985
- Tiwari, L. J. and Terasaki, P. I.: *HLA and Disease Associations*. Springer Verlag, Berlin, 1985
- van der Blik, A. M., van der Velde-Koerts, T., Ling, V., and Borst, P.: Overexpression and amplification of five genes in a multi-drug resistant Chinese hamster ovary cell line. *Mol. Cell. Biol.* 6: 1671-1678, 1986
- van Ommen, G. J. B. and Verkerk, J. M. H.: Restriction analysis of chromosomal DNA in a size range up to two million base pairs by pulse field gradient electrophoresis. In K. E. Davies (ed.): *Human Genetic Diseases: A Practical Approach*, pp. 113-133, IRL Press, Oxford, 1986
- Weiss, E. H., Bloemer, K., Doerner, C., Kuon, W., Lang, M., Pohla, H., Schattenkirchner, M., and Riethmüller, G.: Molecular biology of the HLA-B27 locus. *Brit. J. Rheumatol.*, in press, 1987
- Ziegler, A., Zimmer, F.-J., Fonatsch, C., So, A., and Trowsdale, J.: Mapping of HLA class I and class II genes using HLA deletion mutant cell lines. Eighth International Workshop on Human Gene Mapping. *Cytogenet. Cell Genet.* 40: 788, 1985a
- Ziegler, A., Müller, C., Heinig, J., Radka, S. F., Kömpf, J., and Fonatsch, C.: Monosomy 6 in a human lymphoma cell line induced by selection with a monoclonal antibody. *Immunobiology* 169: 455-460, 1985b

Received July 17, 1987; revised version received September 1, 1987