

Kinetics of Hydride Transfer Reactions from Hydrosilanes to Carbenium Ions. Substituent Effects in Silicenium Ions^{†,‡}

Herbert Mayr,^{*,§} Nils Basso, and Gisela Hagen[§]

Contribution from the Institut für Chemie der Medizinischen Universität zu Lübeck, D-2400 Lübeck, Germany. Received August 20, 1991


Abstract: Rates of hydride transfer from hydrosilanes $\text{HSiR}^1\text{R}^2\text{R}^3$ with widely varying substitution to para-substituted diarylcarbenium ions have been measured in dichloromethane solution. Generally the reactions follow a second-order rate law, $-\text{d}[\text{Ar}_2\text{CH}^+]/\text{dt} = k_2[\text{Ar}_2\text{CH}^+][\text{HSiR}^1\text{R}^2\text{R}^3]$, and k_2 is independent of the degree of ion-pairing and the nature of the counterion (exceptions are reported). The reaction rates are almost independent of solvent polarity. Kinetic isotope effects exclude an SET-type mechanism and are in accord with a polar mechanism with rate-determining formation of silenium ions. The reactivities of para-substituted aryldimethylsilanes are linearly correlated with σ_p ($\rho = -2.46$), not with σ_p^+ . In the series H_3SiHex , H_2SiHex_2 , HSiHex_3 , the relative reactivities are 1.00:155:7890, and in the corresponding phenyl series the reactivity increase is much smaller ($\text{H}_3\text{SiPh}:\text{H}_2\text{SiPh}_2:\text{HSiPh}_3 = 1.00:17.2:119$). As a consequence, trihexylsilane is approximately two orders of magnitude more reactive than triphenylsilane though hexylsilane and phenylsilane show similar reactivities. Tris(trimethylsilyl)silane is just slightly more reactive than trimethylsilane. Replacement of hydrogen by chlorine reduces the reactivity by one order of magnitude. Variation of the electrophilicities of the hydride abstractors does not affect the relative reactivities of the silanes, i.e., constant selectivity (Ritchie-type) relationships are encountered. Correlation equations are given, which permit the calculation of hydride transfer rates from hydrosilanes to any carbenium ion on the basis of $\text{p}K_{\text{R}^+}$ values or the ethanolsis rate constants of the corresponding alkyl chlorides.

I. Introduction

Hydrosilanes are well-known reducing agents for a variety of functional groups.¹ When these reactions are performed in Brønsted or Lewis acidic media, hydride transfer from silicon to positively charged carbon with formation of silenium ions takes place (Scheme I).¹ Despite their thermodynamic stability, silenium ions are very short lived reaction intermediates, and reports on the observation of persistent silenium ions in the condensed phase² have been rejected.³ Information on the stabilities of silenium ions has been obtained from quantum chemical calculations,⁴ gas-phase studies,^{5a-c} electrochemical experiments,^{5f} and kinetic investigations of hydride abstractions from hydrosilanes.⁶

A qualitative order of hydride-donating abilities of hydrosilanes ($\text{HSiEt}_3 > \text{HSi}^i\text{Oct}_3 \approx \text{H}_2\text{SiEt}_2 > \text{H}_2\text{SiPh}_2 \approx \text{HSiPh}_3 > \text{H}_3\text{SiPh}$) was reported by Carey in 1968.⁷ Kinetic investigations of the hydride transfer reactions from triarylsilanes and aryldimethylsilanes to the tris(2,6-dimethoxyphenyl)carbenium ion in acetic acid provided ρ -values of -1.84 and -1.01 , respectively.⁷ The small magnitude of these reaction constants was explained by nucleophilic solvent assistance in replacement of hydride from silicon.⁷

Extensive kinetic studies on hydride transfer reactions from trisubstituted silanes to the trityl cation and tropylium ion have been performed by Chojnowski.^{6a,b} According to Figure 1, replacement of methyl in HSiMe_3 by either Et, Ph, EtS, or Cl reduces the reactivity toward the trityl cation to a various extent. Since the nature of the negative counterion (SbF_6^- , AsF_6^- , PF_6^- , BF_4^- , SbCl_6^- , FeCl_4^-) hardly affected the rates in dichloromethane, an $\text{S}_{\text{N}}2$ -Si pathway has been excluded.^{6a} The structural effects on reactivity and the kinetic isotope effects ($k(\text{HSiR}_3)/k(\text{DSiR}_3) = 1.4-1.5$) have been interpreted in terms of the mechanism in eq 1, with rate-limiting single-electron transfer, followed by a rapid hydrogen shift.^{6a}



[†] Dedicated to Professor F. W. Lichtenthaler on the occasion of his 60th birthday.

[‡] In part presented at the European Symposium on Organic Reactivity, ESOR III, in Göteborg, Sweden, July, 7-12, 1991.

[§] New address: Institut für Organische Chemie und Biochemie, Petersenstrasse 22, D-6100 Darmstadt, Germany.

Scheme I




Table I. Second-Order Rate Constants for the Reaction of the Bis(*p*-anisyl)carbenium Ion (An_2CH^+ , $0.11 \text{ mmol}\cdot\text{L}^{-1}$)^a with Dimethylphenylsilane in the Presence of Various Lewis Acids (CH_2Cl_2 , -70°C)

Lewis acid	$[\text{Lewis acid}]_0$, $\text{mmol}\cdot\text{L}^{-1}$	$[\text{HSiMe}_2\text{Ph}]_0$, $\text{mmol}\cdot\text{L}^{-1}$	k_2 , $\text{L}\cdot\text{mol}^{-1}\cdot\text{s}^{-1}$
TiCl_4	3.26	0.432	1.20
	7.61	0.436	1.15
	16.1	0.432	1.12
Me_3SiOTf	1.11	7.14	1.11
	1.11	17.8	1.10
BCl_3	3.04	0.432	169 ^b
	8.69	0.872	470 ^b
	43.5	0.432	1121 ^b

^a Counterion = TiCl_4^- , OTf^- , or BCl_4^- , depending on Lewis acid employed. ^b Pseudo-second-order rate constant.

In a series of papers, we have reported that the reactivities of various classes of nucleophiles toward para-substituted diaryl-

(1) Reviews: (a) Nagai, Y. *Org. Prep. Proced. Int.* **1980**, *12*, 13. (b) Kursanov, D. N.; Parnes, Z. N.; Loim, N. M. *Synthesis* **1974**, 633. (c) Carey, F. A. *Intra-science Chem. Rept.* **1973**, *7*, 55. (d) Kursanov, D. N.; Parnes, Z. N. *Russ. Chem. Rev.* **1969**, *38*, 812. (e) Keinan, E. *Pure Appl. Chem.* **1989**, *61*, 1737. (f) Hajós, A. In *Houben-Weyl, Methoden der Organischen Chemie*, 4th ed.; Georg Thieme Verlag: Stuttgart, 1981; Vol. IV/1d, pp 1-487. (g) Pawlenko, S. In *Houben-Weyl, Methoden der Organischen Chemie*, 4th ed.; Georg Thieme Verlag: Stuttgart, 1980; Vol. XIII/5, pp 350-361. (h) Colvin, E. W. *Silicon Reagents in Organic Synthesis*; Academic Press: London, 1988.

(2) (a) Lambert, J. B.; McConnell, J. A.; Schulz, W. J., Jr. *J. Am. Chem. Soc.* **1986**, *108*, 2482. (b) Lambert, J. B.; Schulz, W. J., Jr. *J. Am. Chem. Soc.* **1983**, *105*, 1671. (c) Lambert, J. B.; Schulz, W. J., Jr.; McConnell, J. A.; Schilf, W. *J. Am. Chem. Soc.* **1988**, *110*, 2201. (d) Lambert, J. B.; McConnell, J. A.; Schilf, W.; Schulz, W. J., Jr. *J. Chem. Soc., Chem. Commun.* **1988**, 455. (e) Lambert, J. B.; Schilf, W. *J. Am. Chem. Soc.* **1988**, *110*, 6364.






Figure 1. Rate constants for the hydride transfer reactions from $\text{HSiMe}_{3-n}\text{R}_n$ to Ph_3C^+ (CH_2Cl_2 , 25 °C).^{6a}

Scheme II

Kinetic Method



Abbreviations



carbenium ions follow well-behaved linear free energy relationships, which allow the construction of a nucleophilicity scale

(3) (a) Prakash, G. K. S.; Keyaniyan, S.; Aniszfeld, R.; Heiliger, L.; Olah, G. A.; Stevens, R. C.; Choi, H.-K.; Bau, R. *J. Am. Chem. Soc.* **1987**, *109*, 5123. (b) Olah, G. A.; Heiliger, L.; Li, X.-Y.; Prakash, G. K. S. *J. Am. Chem. Soc.* **1990**, *112*, 5991. (c) Wang, N.; Hwu, J. R.; White, E. H. *J. Org. Chem.* **1991**, *56*, 471. (d) Eaborn, C. *J. Organomet. Chem.* **1991**, *405*, 173.

(4) (a) Hopkinson, A. C.; Lien, M. H. *Can. J. Chem.* **1989**, *67*, 991. (b) Apeloig, Y. In *Heteroatom Chemistry*; Block, E., Ed.; VCH-Verlagsgesellschaft: Weinheim, 1990; p 27. (c) Apeloig, Y. In *The Chemistry of Organic Silicon Compounds*; Patai, S., Rappoport, Z., Eds.; Wiley Interscience: New York, 1989; Chapter 2 and literature cited therein. (d) Hopkinson, A. C.; Lien, M. H. *J. Org. Chem.* **1981**, *46*, 998. (e) Apeloig, Y.; Schleyer, P. v. R. *Tetrahedron Lett.* **1977**, *18*, 4647. (f) Apeloig, Y.; Godleski, S. A.; Heacock, D. J.; McKelvey, J. M. *Tetrahedron Lett.* **1981**, *22*, 3297. (g) Godleski, S. A.; Heacock, D. J.; McKelvey, J. M. *Tetrahedron Lett.* **1982**, *23*, 4453. (h) Truong, T.; Gordon, M. S.; Boudjouk, P. *Organometallics* **1984**, *3*, 484.

(5) (a) Angelini, G.; Keheyani, Y.; Laguzzi, G.; Lilla, G. *Tetrahedron Lett.* **1988**, *29*, 4159. (b) Shin, S. K.; Beauchamp, J. L. *J. Am. Chem. Soc.* **1989**, *111*, 900. (c) Li, X.; Stone, J. A. *J. Am. Chem. Soc.* **1989**, *111*, 5586. (d) Drewello, T.; Burgers, P. C.; Zummack, W.; Apeloig, Y.; Schwarz, H. *Organometallics* **1990**, *9*, 1161. (e) Schwarz, H. In *The Chemistry of Organic Silicon Compounds*; Patai, S., Rappoport, Z., Eds.; Wiley Interscience: New York, 1989; Chapter 7 and literature cited therein. (f) Okano, M.; Mochida, K. *Chem. Lett.* **1991**, 819.

Scheme III. Relative Reactivities toward AnPhCH^+ (-70 °C)

HSiAlk_3	k_{rel}
HSiMe_3	≈ 1.00
HSiEt_3	1.95
$\text{HSi}^{(n)\text{Pr}}_3$	3.53
$\text{HSi}^{(n)\text{Bu}}_3$	6.04
$\text{HSi}^{(n)\text{Hex}}_3$	5.93

including compounds of widely differing reactivity.⁸ In order to include hydrosilanes into this nucleophilicity scale, we have systematically studied the influence of the structure on the reactivities of carbenium ions and hydrosilanes, and we will present correlation equations that allow the prediction of rates for a great variety of carbenium ion hydrosilane combinations. We will report the first rate constants for mono- and disubstituted hydrosilanes and, furthermore, present evidence that requires revision of the mechanism outlined in eq 1.

II. Method—The Role of the Negative Counterion

The method recently described for the determination of reactivities of carbenium ions toward alkenes^{8a,h} has analogously been used for this investigation. As previously reported, diarylmethyl halides are added to dichloromethane solutions of Lewis acids that are sufficiently strong to warrant complete ionization. In this way, colored, electrically conducting solutions of $\text{Ar}_2\text{CH}^+\text{MY}_n\text{X}^-$ are produced (Scheme II). When the hydrosilanes are added, covalent products are formed (Scheme II), and the disappearance of the carbenium ions can be monitored photometrically and conductometrically.

Generally, the rate was found to depend linearly on carbenium ion and on hydrosilane concentration, as indicated by eq 2. In accord with

$$-d[\text{Ar}_2\text{CH}^+]/dt = k_2[\text{Ar}_2\text{CH}^+][\text{HSiR}_3] \quad (2)$$

previous investigations,^{6a} the degree of ion-pairing was found not to affect the reaction rates. Identical reactivities of paired and nonpaired carbenium ions have also been observed in reactions with alkenes, and a rationalization has been given.^{8k,9} Table I shows that identical rates were observed for the reaction of HSiMe_2Ph with $\text{An}_2\text{CH}^+\text{TiCl}_5^-$ or $\text{An}_2\text{CH}^+\text{OTf}^-$. This observation corroborates Chojnowski's conclusion that the counterion does not affect the rate-determining step.^{6a}

Table I furthermore shows that excess TiCl_4 or Me_3SiOTf , which is often necessary to achieve complete ionization of the benzhydryl chlorides,¹⁰ does not affect the rates of the reaction. When BCl_3 was utilized for the ionization of An_2CHCl , however, the reaction was found to be 100 to 1000 times faster, depending on the concentration of BCl_3 in the solution. This observation is attributed to the rapid intermediate formation of HBCl_2 ,¹¹ which acts as an efficient hydride donor in the presence of BCl_4^- .

¹¹B NMR spectroscopy (64.1 MHz) supports this interpretation: The singlet of BCl_3 (δ 46.6 with respect to external $\text{BF}_3\cdot\text{OEt}_2$)¹² is instant-

(6) (a) Chojnowski, J.; Fortuniak, W.; Stanczyk, W. *J. Am. Chem. Soc.* **1987**, *109*, 7776. (b) Chojnowski, J.; Wilczek, L.; Fortuniak, W. *J. Organomet. Chem.* **1977**, *135*, 13. (c) Carey, F. A.; Wang Hsu, C.-L. *J. Organomet. Chem.* **1969**, *19*, 29. (d) Traylor, T. G.; Koermer, G. S. *J. Org. Chem.* **1981**, *46*, 3651.

(7) Carey, F. A.; Tremper, H. S. *J. Am. Chem. Soc.* **1968**, *90*, 2578.

(8) (a) Schneider, R.; Grabis, U.; Mayr, H. *Angew. Chem.* **1986**, *98*, 94; *Angew. Chem., Int. Ed. Engl.* **1986**, *25*, 89. (b) Mayr, H.; Pock, R. *Tetrahedron* **1986**, *42*, 4211. (c) Mayr, H.; Pock, R. *Chem. Ber.* **1986**, *119*, 2473. (d) Mayr, H.; Pock, R. *Chem. Ber.* **1986**, *119*, 2497. (e) Schneider, R.; Mayr, H. *Angew. Chem.* **1986**, *98*, 1033; *Angew. Chem., Int. Ed. Engl.* **1986**, *25*, 1016. (f) Mayr, H.; Schneider, R.; Grabis, U. *Angew. Chem.* **1986**, *98*, 1034; *Angew. Chem., Int. Ed. Engl.* **1986**, *25*, 1017. (g) Mayr, H.; Hagen, G. *J. Chem. Soc., Chem. Commun.* **1989**, 91. (h) Mayr, H.; Schneider, R.; Schade, C.; Bartl, J.; Bederke, R. *J. Am. Chem. Soc.* **1990**, *112*, 4446. (i) Mayr, H.; Schneider, R.; Irrgang, B.; Schade, C. *J. Am. Chem. Soc.* **1990**, *112*, 4454. (j) Mayr, H.; Schneider, R.; Grabis, U. *J. Am. Chem. Soc.* **1990**, *112*, 4460. (k) Mayr, H. *Angew. Chem.* **1990**, *102*, 1415; *Angew. Chem., Int. Ed. Engl.* **1990**, *29*, 1371. (l) Irrgang, B.; Mayr, H. *Tetrahedron* **1991**, *47*, 219. (m) Hagen, G.; Mayr, H. *J. Am. Chem. Soc.* **1991**, *113*, 4954. (n) Bartl, J.; Steenken, S.; Mayr, H. *J. Am. Chem. Soc.* **1991**, *113*, 7710.

(9) Mayr, H.; Schneider, R.; Schade, C. *Makromol. Chem. Macromol. Symp.* **1988**, *13/14*, 43.

(10) Schneider, R.; Mayr, H.; Plesch, P. H. *Ber. Bunsenges. Phys. Chem.* **1987**, *91*, 1369.

(11) Soundararajan, R.; Matteson, D. S. *J. Org. Chem.* **1990**, *55*, 2274.

Table II. Second-Order Rate Constants for the Reactions of Diarylcarbenium Ions^a with Silanes in CH₂Cl₂ at -70 °C

silanes	X, Y of (<i>p</i> -XC ₆ H ₄)(<i>p</i> -YC ₆ H ₄)CH ⁺	<i>k</i> ₂ (-70 °C), L·mol ⁻¹ ·s ⁻¹
HSiMe ₃	OMe, H	6.37 × 10 ¹
HSiMe ₂ Et	OMe, H	9.06 × 10 ¹
HSiMeEt ₂	OMe, H	1.16 × 10 ²
HSiEt ₃	OMe, H	1.24 × 10 ² ^b
HSi(ⁿ Pr) ₃	OMe, H	2.25 × 10 ²
HSi(ⁱ Pr) ₃	OMe, H	3.67 × 10 ¹
HSi(ⁿ Bu) ₃	OMe, OMe	4.30
	OMe, OPh	2.15 × 10 ¹
	OMe, Me	1.22 × 10 ²
	OMe, H	3.85 × 10 ²
H ₃ Si(ⁿ Hex)	OMe, H	4.79 × 10 ⁻²
H ₂ Si(ⁿ Hex) ₂	OMe, H	7.43
HSi(ⁿ Hex) ₃	OMe, H	3.78 × 10 ²
H ₃ SiPh	OMe, H	6.96 × 10 ⁻²
	OPh, H	3.61 × 10 ⁻¹
	Me, Me	3.37
	Me, H	1.44 × 10 ¹
H ₂ SiPh ₂	OMe, H	1.20
HSiPh ₃	OMe, OMe	6.52 × 10 ⁻²
	OMe, Me	2.22
	OMe, H	8.27 ^c
	OPh, H	4.52 × 10 ¹
	Me, Me	3.47 × 10 ²
H ₂ SiMePh	OMe, H	5.33
HSiMe ₂ Ph	OMe, OMe	1.12 ^d
	OMe, Me	3.16 × 10 ¹
	OMe, H	1.49 × 10 ² ^e
	OPh, H	6.92 × 10 ²
	Me, Me	4.24 × 10 ³
HSiMe ₂ Tol	OMe, H	3.30 × 10 ²
HSiMe ₂ An	OMe, H	8.71 × 10 ²
HSiMe ₂ (<i>p</i> -Cl-C ₆ H ₄)	OMe, H	4.96 × 10 ¹
HSiMePh ₂	OMe, H	2.24 × 10 ¹
HSiMe ₂ CH ₂ Ph	OMe, H	2.58 × 10 ¹
HSiMe ₂ CH ₂ Cl	OMe, H	2.11 × 10 ⁻¹
HSiMe ₂ Cl	OMe, H	2.1 × 10 ⁻¹
HSiMeCl ₂	Me, Me	≈ 5 × 10 ⁻⁴
HSi(SiMe ₃) ₃	OMe, H	4.57 × 10 ²
HSiMe ₂ (OSiMe ₃)	OMe, H	5.86 × 10 ¹

^a Counterion = TiCl₅⁻ or OTf⁻, depending on the Lewis acid employed. ^b Δ*H*[‡] = (18.6 ± 0.4) kJ·mol⁻¹; Δ*S*[‡] = (-110 ± 2) J·mol⁻¹·K⁻¹. ^c Δ*H*[‡] = (22.2 ± 0.6) kJ·mol⁻¹; Δ*S*[‡] = (-115 ± 3) J·mol⁻¹·K⁻¹. ^d Δ*H*[‡] = (30.1 ± 0.3) kJ·mol⁻¹; Δ*S*[‡] = (-92.5 ± 1.2) J·mol⁻¹·K⁻¹. ^e Δ*H*[‡] = (20.0 ± 0.5) kJ·mol⁻¹; Δ*S*[‡] = (-102 ± 3) J·mol⁻¹·K⁻¹.


neously replaced by a doublet (δ 57.5, *J* = 200 Hz), when HSiMe₂Ph is added to the solution of BCl₃ in CH₂Cl₂ at -70 °C. When the NMR tube is opened at ambient temperature, a volatile compound is released which immediately catches fire in the air; therefore tetrachloroborates cannot be employed as counterions for kinetic investigations with hydrosilanes.

Control experiments showed that the ¹H NMR spectrum of HSiMe₂Ph in CD₂Cl₂ is not altered, when TiCl₄ or Me₃SiOTf was added, indicating that these Lewis acids do not react with ordinary silanes at low temperature. Accordingly, the presence of variable amounts of these Lewis acids does not affect the reaction rates (see above). As similar observations were made for other silanes (see Supplementary Material), we can conclude that the rate constants measured in the presence of TiCl₅⁻ or F₃CSO₃⁻ (Table II) reflect the rate of the hydride transfer step from silicon to carbon.

III. Variation of the Silanes

1. Trialkylsilanes. The successive replacement of methyl by ethyl in HSiMe₃ has been reported to cause a slight, but steady decrease of the hydride transfer rate to the trityl cation (Figure 1, *k*(HSiMe₃)/*k*(HSiEt₃) = 1.6).^{6a} Entries 1–4 of Table II show the opposite effect on the rates of hydride abstraction by AnPhCH⁺. As the effects are small in both series, one can assume that the reactivity order toward the diarylcarbenium ion is con-

Scheme IV

**Scheme V.** Relative Reactivities toward AnPhCH⁺ (-70 °C)

	<i>k</i> _{rel} (R = ⁿ Hex)	<i>k</i> _{rel} (R = Ph)
H ₃ SiR	1.00	1.00
H ₂ SiR ₂	155	17.2
HSiR ₃	7890	119

Scheme VI. Relative Reactivities of HSiMe₂Ph_{3-x} toward AnPhCH⁺ (-70 °C)

	HSiPh ₃	HSiMePh ₂	HSiMe ₂ Ph	HSiMe ₃
<i>k</i> _{rel}	1.00	2.71	18.0	7.70

trolled by inductive effects ($\sigma_1(\text{Et}) = -0.055$, $\sigma_1(\text{Me}) = -0.046$),¹³ which are slightly overcompensated by steric effects when the bulky triphenylcarbenium ion acts as the hydride acceptor.

Accordingly, Scheme III illustrates that hydride abstractions from tri(*n*-alkyl)silanes by AnPhCH⁺ are controlled by inductive rather than steric effects. While the increase of the chain length of the *n*-alkyl groups causes an increase of the hydride transfer rate, branching reduces the rate constants. Though isopropyl ($\sigma_1 = -0.064$)¹³ is a stronger inductive donor than *n*-propyl ($\sigma_1 = -0.057$),¹³ HSi(ⁱPr)₃ is approximately 6 times less reactive than HSi(ⁿPr)₃ (Table II).

2. Para-Substituted Aryldimethylsilanes: Hammett Analysis. Substituents at the aromatic ring have a relatively small effect on the rate of formation of silicenium ions. Scheme IV shows the widely differing effects which *p*-methoxy groups exert on the transition states that lead to silicenium (via hydride abstraction) and analogous carbenium ions (via S_N1 reaction). The rate constants measured for the reaction of AnPhCH⁺ with four aryldimethylsilanes correlate only poorly with σ_p^+ ($r = 0.95$) but give a good correlation with σ_p ($\rho = -2.46$, $r = 0.996$, Figure 2), indicating that mesomeric effects play a minor role for stabilizing silicenium ions.

3. Comparison of Alkyl and Phenyl Groups. Since exchange of hydrogen by chlorine considerably weakens the reactivity of the remaining Si–H groups (see below), it is possible to investigate the selective abstraction of only one hydride from dihydro- and trihydrosilanes (eq 3). According to Table II, phenylsilane and H₃SiAlkyl + ArAr'CH⁺ TiCl₅⁻ → H₂SiClAlkyl + ArAr'CH₂ + TiCl₄ (3)

n-hexylsilane hardly differ in reactivity toward AnPhCH⁺ (*k*(H₃SiPh)/*k*(H₃SiHex) = 1.5). Successive replacement of hydrogen by hexyl (H₃SiHex → H₂SiHex₂ → HSiHex₃) increases the reactivity by two orders of magnitude per hexyl group while in the analogous phenyl series (H₃SiPh → H₂SiPh₂ → HSiPh₃) a reactivity increase of only one order of magnitude is observed (Scheme V), with the result that the reactivity of HSiHex₃ exceeds that of HSiPh₃ by almost two orders of magnitude (Table II).

With the assumption that the hydride transfer rates reflect the stabilization of the resulting silicenium ions, one would deduce that trialkyl-substituted silicenium ions are stabilized to a greater

(12) Wrackmeyer, B.; Köster, R. NMR-Spektroskopie von Organo-Bor-Verbindungen, Organobor-Verbindungen III. In *Houben-Weyl, Methoden der Organischen Chemie*, 4th ed.; Georg Thieme Verlag: Stuttgart, 1984; Vol. XIII/3c, p 399.

(13) (a) Taft, R. W., Jr.; Lewis, I. C. *J. Am. Chem. Soc.* **1958**, *80*, 2436. (b) Taft, R. W., Jr.; Lewis, I. C. *Tetrahedron* **1959**, *5*, 210. (c) Levitt, L. S.; Levitt, B. W. *Chem. Ind. (London)* **1970**, 990. (d) Levitt, L. S.; Widing, H. F. *Prog. Phys. Org. Chem.* **1976**, *12*, 119. (14) Brown, H. C.; Okamoto, Y. *J. Am. Chem. Soc.* **1958**, *80*, 4979.




Figure 2. Correlation between the reactivities of the aryltrimethylsilanes toward AnPhCH^+ (CH_2Cl_2 , -70°C) with σ_p ($\rho = -2.46$).

Table III. Calculated Stabilization Energies for Silicenium and Carbenium Ions ($\text{kcal}\cdot\text{mol}^{-1}$, $\text{MP3}/6\text{-}31\text{G}^*/6\text{-}31\text{G}^*$)^{4b}

R	ΔE	
	eq 4	eq 5
CH_3	15.1	34.1
NH_2	36.8	97.8
OH	17.9	62.7
F	-2.3	21.5
SiH_3	12.6	17.7
PH_2	13.5	60.0
SH	18.4	60.9
Cl	2.0	26.6

extent than the triphenyl-substituted analogues—in contrast to the substituent effects known for carbenium ions.¹⁵ The minor importance of π -conjugation for the stabilization of silenium ions, which is also reflected by the Hammett correlation depicted in Figure 2, has previously been deduced from ab initio MO calculations on HO- and HS-substituted silenium ions.^{4b}

Combination of the effects responsible for the reactivity orders $\text{H}_3\text{SiAlk} \lesssim \text{H}_3\text{SiPh}$ and $\text{HSiAlk}_3 \gg \text{HSiPh}_3$ leads to a shallow reactivity maximum for HSiMe_2Ph in the $\text{HSiMe}_x\text{Ph}_{3-x}$ series (Scheme VI).


4. Heteroatom-Substituted Silanes. According to ab initio MO calculations, substituent effects are generally greater in carbenium ions (eq 5) than in silenium ions (eq 4, Table III).^{4b}



Oxygen Substitution. The small effect of π -conjugation on silenium ion stabilization, which has been discussed for aryl-substituted silenium ions (see above), can also be derived from Table III. While OH stabilizes carbenium ions twice as much as CH_3 , these two substituents have a similar effect on silenium ions. Accordingly, Chojnowski reported comparable hydride transfer rates from HSiMe_3 and $\text{HSiMe}_2(\text{O}^i\text{Pr})$ to the trityl cation.^{6b}

While we were able to reproduce these results at 25°C , we did not find second-order kinetics for the reaction of $\text{HSiMe}_2(\text{OEt})$ with $\text{Ph}_3\text{C}^+ \text{OTf}^-$ or with diarylcarbenium triflates at low temperatures (e.g. -70°C). Our hypothesis that reversible oxonium ion formation (eq 6) was responsible for the unusual kinetic behavior of alkoxyhydrosilanes was examined by combining a tetrasubstituted silane with a carbenium salt: When Me_3SiOMe was added to a solution of $\text{An}_2\text{CH}^+ \text{OTf}^- / \text{Me}_3\text{SiOTf}$ (excess) at -70°C , the carbenium ion absorbance disappeared immediately

(<5 s) but was slowly restored to its initial value (≈ 0.5 h).¹⁶ Hydride transfer from alkoxyhydrosilanes, therefore, may follow a complex mechanism (eq 6), and we have excluded alkoxyhydrosilanes from this investigation.



In contrast, the reaction of pentamethyldisiloxane [$\text{HSiMe}_2\text{OSiMe}_3$] and AnPhCH^+ at -70°C follows clean second-order kinetics. Product analysis shows the formation of the corresponding diarylmethane AnPhCH_2 . The rate constant, which is identical for the counterions TiCl_5^- and OTf^- , is very similar to that determined for hydride transfer from trimethylsilane, indicating comparable activating effects of Me and OSiMe_3 .

Amino Substitution. Analogous to the situation with alkoxyhydrosilanes, we were unable to measure hydride transfer rates from amino-substituted silanes. When $\text{HSiMe}_2(\text{NMe}_2)$ was added to a solution of $\text{An}_2\text{CH}^+ \text{OTf}^-$ at -70°C , a rapid decrease of absorbance was observed, but product analysis showed that amino group transfer instead of hydride transfer had taken place (eq 7).



Silyl Groups. From the calculations quoted in Table III it has been deduced that CH_3 stabilizes carbenium ions considerably better than SiH_3 , while both groups exert a similar effect on silenium ions. Accordingly, the superiority of CH_3 over SiMe_3 to stabilize carbenium ions has been observed in solvolysis reactions.¹⁷ Equal stabilizing effects of alkyl and SiMe_3 on silenium ions might now be derived from the almost identical rates of hydride transfer from $\text{HSi}(\text{SiMe}_3)_3$ and $\text{HSi}(\text{tBu})_3$ to AnPhCH^+ (Table II). It has to be noted, however, that the Si-H bond in $\text{HSi}(\text{SiMe}_3)_3$ is $88 \text{ kJ}\cdot\text{mol}^{-1}$ weaker than that in HSiEt_3 ¹⁸ so that ground-state effects cannot be ignored. One should, therefore, more precisely conclude that the relative stabilization of silanes and silenium ions by alkyl and trialkylsilyl groups is identical. As a consequence, the large reactivity differences between trialkylsilanes and tris(trimethylsilyl)silane that have been encountered in radical reactions^{18,19} are not observed in ionic hydride transfer processes.

Chlorine. Whereas chlorine has been calculated to slightly stabilize silenium ions relative to hydrogen (Table III), hydride abstraction from chlorodimethylsilane is one order of magnitude slower than that from dimethylsilane (eq 8). The rate constant

$$\frac{k(\text{HSiMe}_2\text{Cl})}{k(\text{H}_2\text{SiMe}_2)} \approx 0.09 \quad (8)$$

[for AnPhCH^+ , -70°C]

for H_2SiMe_2 ($k \approx 2.2$), necessary for this comparison, has been estimated from $k(\text{H}_2\text{SiHex}_2)$, assuming a methyl/hexyl ratio of 0.55 per alkyl group exchange (from $\text{HSiMe}_3/\text{HSiHex}_3$).

Analogously, $k(\text{H}_3\text{SiMe}) \approx 0.026$ is estimated from the value of $k(\text{H}_3\text{SiHex})$ and linear free energy relationships (see below) were used to approximate a rate constant for $\text{HSiMeCl}_2 + \text{AnPhCH}^+$ (1.6×10^{-5} , -70°C) from the measured rate constant for $\text{HSiMeCl}_2 + \text{ToI}_2\text{CH}^+$ (Table II). Equation 9 shows that

$$\frac{k(\text{HSiMeCl}_2)}{k(\text{H}_3\text{SiMe})} \approx 0.0006 \quad [\text{for } \text{AnPhCH}^+, -70^\circ\text{C}] \quad (9)$$

replacement of two hydrogens by chlorine atoms reduces the rate of hydride transfer by more than three orders of magnitude. Smaller deactivating effects by chlorine have been reported for hydride abstractions with Ph_3C^+ at 25°C .^{6a} Accidentally, the

(16) Details of this reaction sequence have not been investigated. It is assumed, however, that the rapidly formed oxonium triflate reacts with the excess Me_3SiOTf to restore the dianisylcarbenium triflate.

(17) Review: Lambert, J. B. *Tetrahedron* **1990**, *46*, 2677.

(18) Kanabus-Kaminska, J. M.; Hawari, J. A.; Griller, D.; Chatgililoglu, C. *J. Am. Chem. Soc.* **1987**, *109*, 5267.

(19) (a) Ballestri, M.; Chatgililoglu, C.; Clark, K. B.; Griller, D.; Giese, B.; Kopping, B. *J. Org. Chem.* **1991**, *56*, 678. (b) Arya, P.; Samson, C.; Lesage, M.; Griller, D. *J. Org. Chem.* **1990**, *55*, 6248 and references cited therein.

(15) Streitwieser, A., Jr. *Solvolytic Displacement Reactions*; McGraw Hill: New York, 1962; p 43.




Figure 3. Correlation between the reactivities of the silanes $\text{HSiMe}_2\text{CH}_2\text{X}$ (AnPhCH^+ , CH_2Cl_2 , -70°C) with the σ_1 -values of X ($\rho = -5.21$).

reactivities of HSiMe_2Cl and $\text{HSiMe}_2(\text{CH}_2\text{Cl})$ toward AnPhCH^+ are identical, i.e. the stronger ($-$)-effect of chlorine that is operating in the first compound, where chlorine is directly attached to silicon, must be compensated by the π -donor effect.

Figure 3 shows a linear correlation of the reactivities of the silanes $\text{HSiMe}_2(\text{CH}_2\text{X})$ toward AnPhCH^+ with the inductive substituent constants σ_1 of X. The positive σ_1 values of Ph and Cl explain that benzyltrimethylsilane is 2.5 times and (chloromethyl)dimethylsilane 300 times less reactive than trimethylsilane.

IV. Variation of the Hydride Acceptors

In previous work, we found fair correlations ($r = 0.97$ – 0.991) between the reactivities of para-substituted diarylcarbenium ions toward π -nucleophiles (e.g. alkenes^{8j,k} or allyl-element compounds^{8m}) and their $\text{p}K_{\text{R}^+}$ -values²⁰ or the $\text{S}_{\text{N}}1$ reactivities of the corresponding benzhydryl chlorides.²¹ The correlations between the rate constants of these reaction series are of higher quality ($r = 0.994$ – 0.9998), however. Equation 10 has been used to

$$\log k_2 = s \log k_0 + c \quad (10)$$

calculate rate constants (k_2 , -70°C , CH_2Cl_2) for the reactions of benzhydryl cations with π -nucleophiles from the $\log k_0$ values of the benzhydryl cations (reactivity toward the reference nucleophile 2-methyl-1-pentene at -70°C in CH_2Cl_2) and the s and c parameters, which have been determined for several π -nucleophiles.^{8j,k,m}

Accidentally, the rate constants for the attack of diarylcarbenium ions at 2-methyl-1-pentene and the silanes listed in Table II are of comparable magnitude. Correlations can, therefore, be based on the same reference electrophiles ($\log k_0$ as in eq 10). Figure 4 and Table IV show well-behaved linear correlations between the reactivities of diarylcarbenium ions toward hydrosilanes and 2-methyl-1-pentene. The almost parallel lines in Figure 4 indicate that the relative reactivities of the hydrosilanes are independent of the electrophilicities of the hydride abstractors, i.e., selectivity is independent of reactivity. Constant selectivity relationships (Ritchie-type behavior),²² as previously reported for the reactions of benzhydryl cations with terminal vinyl compounds, are thus also observed for hydride transfer reactions. The smaller values of s for the hydride transfer reactions are indicative of transition states in which charge transfer is less advanced than in the reactions of benzhydryl cations with π -nucleophiles.

(20) (a) Mindl, J.; Vecera, M. *Collect. Czech. Chem. Commun.* **1971**, *36*, 3621. (b) Deno, N. C.; Jaruzelski, J.; Schriesheim, A. *J. Am. Chem. Soc.* **1955**, *77*, 3044. (c) Deno, N. C.; Schriesheim, A. *J. Am. Chem. Soc.* **1955**, *77*, 3051.

(21) Schade, C.; Mayr, H. *Tetrahedron* **1988**, *44*, 5761.

(22) (a) Ritchie, C. D. *Acc. Chem. Res.* **1972**, *5*, 348. (b) Ritchie, C. D. *Can. J. Chem.* **1986**, *64*, 2239.

Table IV. Correlation between the Reactivities of Diarylcarbenium Ions ($\text{ArAr}'\text{CH}^+$) toward Hydrosilanes with the Corresponding Reactivities toward 2-Methyl-1-pentene (-70°C , CH_2Cl_2)^a

silane	s	c	r	no. of points
$\text{H}_2\text{C}=\text{C}(\text{CH}_3)\text{C}_3\text{H}_7$	1.00	0.00	reference	
H_3SiPh	0.762	-2.25	0.998	4
HSiPh_3	0.727	-7.45×10^{-2}	0.9995	5
HSiMe_2Ph	0.705	+1.14	0.9998	5
$\text{HSi}(\text{tBu})_3$	0.648	+1.72	0.994	4

^a Correlation according to eq 10; $\log k_0 = -1.535$ (An_2CH^+), -0.7721 ($\text{An}(\text{PhOP})\text{CH}^+$), 0.5289 (AnTolCH^+), 1.412 (AnPhCH^+), 2.456 ($(\text{PhOP})\text{PhCH}^+$), 3.531 (Tol_2CH^+), ~ 4.52 (TolPhCH^+) extrapolated from reactivity of compound **2j** in ref 8j.




Figure 4. Correlation of the reactivities of the hydrosilanes toward diarylcarbenium ions (para substituents in figure) with the corresponding reactivities toward 2-methyl-1-pentene (CH_2Cl_2 , -70°C).

Since further hydrosilanes can be expected to possess similar s -values, the average value $s = 0.711$ can be used to calculate c for those silanes, for which only a single rate constant (usually toward AnPhCH^+ , $\log k_0 = 1.412$) is listed in Table II. Insertion of $s = 0.711$ and $\log k_0 = 1.412$ in eq 10 yields

$$c = \log k_2(\text{AnPhCH}^+) - 1.004 \quad (11)$$

Now, rate constants for all combinations of diarylcarbenium ions (with known reactivity toward 2-methyl-1-pentene, $\log k_0$) with hydrosilanes (with known reactivity toward one carbenium ion) can be estimated.

As hydrosilanes follow the same linear free energy relationships as π -nucleophiles, the previously derived eqs 12 and 13 can also be used for obtaining a rough approximation for the reactivity of any diarylcarbenium ion (characterized by $\text{p}K_{\text{R}^+}$ or ethanolysis rate constant) toward a hydrosilane (characterized by s and c).

$$\log k_2(\text{CH}_2\text{Cl}_2, -70^\circ\text{C}) = -7.42s - 1.07s \text{p}K_{\text{R}^+} + c \quad (12)$$

$$\log k_2(\text{CH}_2\text{Cl}_2, -70^\circ\text{C}) = 0.90s - 1.26s \log k_{\text{sol}} + c \quad (13)$$

Though eqs 12 and 13 have been derived for benzhydryl cations,^{8j} their validity is not necessarily restricted to this class of electrophiles, and there is reason to believe that eqs 12 and 13 also hold for other types of carbenium ions,^{8k} if systems with large steric requirements are excluded. When eq 12 was applied to reactions of trityl cations with π -nucleophiles, rate constants were calculated that are 3–4 orders of magnitude greater than the experimentally determined values.²³ It has been concluded, therefore, that the

(23) Schade, C.; Mayr, H. *Makromol. Chem., Rapid Commun.* **1988**, *9*, 477.

Table V. Calculated (eq 12) and Experimental Rate Constants for the Hydride Abstraction by the Trityl Cation (Ph_3C^+ , -70°C , CH_2Cl_2)

silane	$k_2(\text{calc})^a$	$k_2(\text{exp})$
HSiEt_3	7.24 ^b	0.268 ^c
HSiMe_2Ph	8.7	0.175
HSiMe_2Cl	0.012 ^b	0.0031

^a $\text{p}K_{\text{a}}^+ = -6.63$ (ref 20). ^b $s = 0.710$ (average). ^c $\Delta H^\ddagger = 31.8 \pm 0.5$ $\text{kJ}\cdot\text{mol}^{-1}$, $\Delta S^\ddagger = -96.1 \pm 2.1$ $\text{J}\cdot\text{mol}^{-1}\cdot\text{K}^{-1}$.

electrophilicity of trityl cations is strongly reduced by steric factors.

Table V shows that steric effects play a minor role in hydride transfer reactions. The experimentally determined rate constants for hydride abstractions by Ph_3C^+ are just 1–2 orders of magnitude smaller than those calculated on the basis of eq 12. As trityl cations are the carbenium ions that showed the greatest deviations from linear free energy relationships for additions to π -systems,^{8k} it can be expected that eqs 12 and 13 provide good estimates for the rates of hydride abstractions by most types of carbenium ions.

V. Reaction Mechanism. Evidence against SET Processes

In accord with a rate-determining step, in which charge separation is neither created nor deleted, the rate of hydride transfer from HSiMe_2Ph to An_2CH^+ is almost equal in solvents of different polarity, ranging from dichloromethane to nitromethane (Table VI). This behavior is in agreement with the polar as well as the SET mechanism.


Kinetic isotope effects should be able to differentiate these two possibilities, because only in the polar mechanism does cleavage of the Si–H bond occur in the rate-determining step. The first four entries of Table VII show $k_{\text{H}}/k_{\text{D}}$ ratios of approximately 2 (-70°C), in accordance with Chojnowski's data ($k_{\text{H}}/k_{\text{D}} = 1.4$ – 1.5),^{6a} which refer to 25°C . Only small, unsystematic changes of $k_{\text{H}}/k_{\text{D}}$ were observed, when the electrophilicity of the hydride abstractor was varied by four orders of magnitude (see hydride abstractions from HSiPh_3 , Table VII).

Chojnowski interpreted the observed $k_{\text{H}}/k_{\text{D}}$ ratios as secondary deuterium isotope effects and considered the independence of the isotope effects of the reactivity as support for the suggested SET mechanism.^{6a} Since we doubted this interpretation, we have synthesized mono- and dideuteriodiphenylsilanes, in order to identify the origin of the isotope effects.

If the observed isotope effects were secondary effects, $k(\text{H}_2\text{SiPh}_2)/k(\text{HDSiPh}_2)$ should be similar to that observed for compounds $\text{HSiR}_3/\text{DSiR}_3$, and $k(\text{H}_2\text{SiPh}_2)/k(\text{D}_2\text{SiPh}_2)$ should be the square of this ratio (i.e. 3 to 4.5). Table VII shows that this is not the case and that the last two entries show much smaller isotope effects than expected on the basis of this hypothesis.

A consistent scheme arises, however, if rate-determining Si–H cleavage is assumed. In this case, the product ratio obtained by reaction of HDSiPh_2 with AnPhCH^+ must reflect the ratio of the kinetically determined rate constants.

When HDSiPh_2 was treated with AnPhCH^+ , hydride and deuteride transfer took place in a 1.82:1 ratio (eq 14). This ratio



is used to split the kinetically determined rate constant for HDSiPh_2 (Table VII) into the partial rate constants shown in Scheme VII. The ratios of the upper right and the lower left partial rate constants of Scheme VII (2.06 ± 0.05) now correspond to the primary isotope effect, and the ratios of the upper left and lower right rate constants (1.13 ± 0.03) correspond to the secondary isotope effect. As expected for a mechanism with rate-determining Si–H cleavage, the primary kinetic isotope effect thus derived is of similar size as the $k_{\text{H}}/k_{\text{D}}$ ratio for trisubstituted silanes.

(24) Reichardt, C. *Solvents and Solvent Effects in Organic Chemistry*, 2nd ed.; VCH-Verlagsgesellschaft: Weinheim, 1988.

(25) (a) Collins, C. J.; Bowman, N. S., Eds. *Isotope Effects in Chemical Reactions*; ACS Monograph 167; American Chemical Society: Washington, DC, 1970. (b) Scheppele, S. E. *Chem. Rev.* **1972**, *72*, 511.





Figure 5. Comparison of hydride transfer rates from hydrosilanes and some hydrocarbons to AnPhCH^+ (CH_2Cl_2 , -70°C). Superscript *a* designates data estimated from LFER in ref 29 assuming $\Delta S^\ddagger = -110$ $\text{J}\cdot\text{mol}^{-1}\cdot\text{K}^{-1}$. Superscript *b* designates data estimated from LFER in ref 30 assuming $\Delta S^\ddagger = -110$ $\text{J}\cdot\text{mol}^{-1}\cdot\text{K}^{-1}$.

Scheme VII. Partial Rate Constants (k_2 , $\text{L}\cdot\text{mol}^{-1}\cdot\text{s}^{-1}$) for the Reactions of AnPhCH^+ with Mono-, Di-, and Nondeuteriated Diphenylsilane (CH_2Cl_2 , -70°C)



If one were to adhere to the mechanism with rate-determining single-electron transfer and successive hydron shift (eq 1), one would not only have to explain the enormous size of the secondary isotope effect but also the different magnitude of $k_{\text{H}}/k_{\text{D}}$ for di- and trisubstituted silanes. Furthermore, one would have to postulate that the coincidence of the $k_{\text{H}}/k_{\text{D}}$ ratios determined by rate (Table VII) and by product studies (eq 14) were entirely accidental.

VI. Conclusion

The kinetic isotope effects are in accord with the polar mechanism illustrated in Scheme II but not with the SET process formulated in eq 1. This result is not unexpected in view of the fact that the ionization potentials of silanes (Ph-SiH_3 , 9.18 eV; Ph-SiMe_3 , 9.00 eV)²⁶ are considerably greater than the electron affinities of benzhydryl cations (Ph_2CH^+ , 7.32 eV),²⁷ which implies that SET is energetically unfavorable.

In the polar mechanism, hydride transfer takes place in the rate-determining step, and it may be assumed that the stabilization of the silicenium ions thus produced is reflected by the reaction rates. Most of the preceding discussions on substituent effects have, therefore, implied the proportionality between ΔG^\ddagger and ΔG° of the rate-determining step (Bell–Evans–Polanyi principle).²⁸

(26) Frey, J. E. *Appl. Spectrosc. Rev.* **1987**, *23*, 247.

(27) Franklin, J. L. In *Carbenium Ions*; Olah, G. A., Schleyer, P. v. R., Eds.; Wiley-Interscience: New York, 1968; Vol. 1, Chapter 2.

(28) For an excellent discussion see: Dewar, M. J. S.; Dougherty, R. C. *The PMO-Theory of Organic Chemistry*; Plenum Press: New York, 1975; p 218.

Table VI. Reaction of Bis(*p*-methoxyphenyl)carbenium Triflate (An₂CH⁺OTf⁻) with Dimethylphenylsilane in Various Solvents^a

solvent	ϵ^b	$E_T(30)^c$	temp range, °C	ΔH^\ddagger , kJ·mol ⁻¹	ΔS^\ddagger , J·mol ⁻¹ ·K ⁻¹	$\Delta G^\ddagger(-20\text{ °C})$, kJ·mol ⁻¹	$k_2(-20\text{ °C})$, L·mol ⁻¹ ·s ⁻¹
CH ₂ Cl ₂	8.9	40.7	-70 to +10	30.1	-92.5	53.5	47.7
(CH ₂ Cl) ₂	10.37	41.3	-20 to +30	27.9	-98.2	52.8	68.4
H ₃ C-CN	37.5 ^d	46.0	-20 to +30	27.6	-99.5	52.8	67.5
H ₃ C-NO ₂	35.94	46.3	-10 to +20	35.1	-69.4	52.7	71.4

^a [An₂CH⁺]₀ = 10⁻⁴ mol·L⁻¹; [PhMe₂SiH] = 2 × 10⁻⁴ to 1 × 10⁻³ mol·L⁻¹. ^b Dielectric constant at 25 °C; ref 24. ^c $E_T(30)$ in kcal·mol⁻¹; ref 24. ^d 20 °C.

Table VII. Rate Constants for the Reactions of Diarylcarbenium Ions [(*p*-XC₆H₄)(*p*-YC₆H₄)CH⁺] with Deuteriosilanes in CH₂Cl₂ at -70 °C

silane	X, Y	k_2 , L·mol ⁻¹ ·s ⁻¹	k_H/k_D
DSiEt ₃	OMe, H	73.6	1.68
DSiMe ₂ Ph	OMe, H	75.8	1.97
DSiPh ₃	OMe, H	4.38	1.89
DSi(^t Pr) ₃	OMe, H	17.3	2.09
DSiPh ₃	OMe, OMe	0.0305	2.14
DSiPh ₃	Me, Me	168	2.07
HDSiPh ₂	OMe, H	0.802	1.50
D ₂ SiPh ₂	OMe, H	0.516	2.33

While this relationship has been found to hold roughly for hydride abstractions from CH groups, i.e. hydride transfer is usually the faster the better stabilized the produced carbenium ion,²⁹ deviations from the proportionality between ΔpK_R^+ and $\log k$ (hydride transfer) have been reported.³⁰ Figure 5, for example, which compares hydride transfer rates from SiH and some CH groups, shows that triphenylmethane is a considerably less active hydride donor than bis(*p*-anisyl)methane though the resulting carbenium ions differ by less than one pK_R^+ unit. Caution is necessary, therefore, when thermodynamic properties are being derived from these kinetic data.

VII. Experimental Section

For the kinetic experiments the previously described apparatus and methods have been used.^{8h} The silanes were either commercially available or synthesized by LiAlH₄ (or LiAlD₄) reduction of the corresponding chlorosilanes.

The partial rate constants for the reaction of HDSiPh₂ with AnPhCH⁺ were obtained by mass spectroscopic determination of the product ratio AnPhCH₂/AnPhCHD (EI, 70 eV). For this purpose the relative intensities of the peaks with $m/e = 197, 198, 199$, and 200 were determined for AnPhCH₂ (39.0:100:14.9:1.17) and AnPhCHD (5.63:38.0:100:14.9). Three mixtures with molar ratio AnPhCH₂/AnPhCHD from 0.5 to 2.5 have then been prepared, and the product ratio H/D (H for AnPhCH₂; D for AnPhCHD) has been calculated by solving the following system of linear equations for the intensities I of the peaks with $m/e = 197-200$, using the method of least squares:

$$39H + 5.63D = I(197)$$

$$100H + 38.0D = I(198)$$

$$14.9H + 100D = I(199)$$

$$1.17H + 14.9D = I(200)$$

(29) Dauben, H. J.; McDonough, L. M. Unpublished results. McDonough, L. M. Ph.D. Thesis, University of Washington, 1960.

(30) Bethell, D.; Hare, G. J.; Kearney, P. A. *J. Chem. Soc., Perkin Trans. 2* 1981, 684.

The deviation between calculated and experimental product ratios was 2-3%.

Three competition experiments (AnPhCH⁺ TiCl₅⁻ was combined with 1:1 to 4:1 mixtures of D₂SiPh₂ and H₂SiPh₂) were then carried out to give relative reactivities $k(\text{H}_2\text{SiPh}_2)/k(\text{D}_2\text{SiPh}_2) = 2.30, 2.25$, and 2.10, in good agreement with the reactivity ratio, determined by kinetic measurements (2.33, Table VII). With the method thus confirmed, the intramolecular competition experiment has been performed: AnPhCHCl (0.859 mmol) was dissolved in 90 mL of CH₂Cl₂ at -78 °C (N₂ atmosphere), and a solution of HDSiPh₂ (1.79 mmol) in CH₂Cl₂ (2 mL) was added. After the addition of TiCl₄ (0.2 mmol), the mixture was kept at -78 °C for 15 h and then poured into concentrated aqueous ammonia (100 mL). After extraction with CH₂Cl₂ (3 × 50 mL), the solution was dried over CaCl₂ and evaporated, and the residue was analyzed by GC/MS. Since the AnPhCH₂/AnPhCHD peak obtained by GC showed an inhomogeneous isotope distribution, the mass spectra summarized over the whole GC peak were evaluated (the same procedure has been used for confirming the method). From the peaks m/e (relative intensity) = 197 (35.4), 198 (100), 199 (58.0), and 200 (7.87) a reactivity ratio of 1.82 is calculated, which allows the measured rate constants to be split into the partial rate constants 0.284 (for D abstraction) and 0.518 (for H abstraction).

Acknowledgment. We thank Ms. Andrea Riemann for the mass spectroscopic analysis, Dr. E. Popowski (Rostock) for a generous gift of pentamethyldisiloxane and Mr. S. Seifert for improving the software to control the work station for the kinetic experiments. Financial support by the Deutsche Forschungsgemeinschaft and the Fonds der Chemischen Industrie is gratefully acknowledged.

Registry No. HSiMe₃, 993-07-7; HSiMe₂Et, 758-21-4; HSiMeEt₂, 760-32-7; HSiEt₃, 617-86-7; HSi(^tPr)₃, 998-29-8; HSi(ⁱPr)₃, 6485-79-6; HSi(ⁿBu)₃, 998-41-4; H₃Si(ⁿHex), 1072-14-6; H₂Si(ⁿHex)₂, 1002-78-4; HSi(ⁿHex)₃, 2929-52-4; H₃SiPh, 694-53-1; H₂SiPh₂, 775-12-2; HSiPh₃, 789-25-3; H₂SiMePh, 766-08-5; HSiMe₂Ph, 766-77-8; HSiMe₂Tol, 1432-39-9; HSiMe₂An, 1432-38-8; HSiMe₂(*p*-Cl-C₆H₄), 1432-31-1; HSiMePh₂, 776-76-1; HSiMe₂CH₂Ph, 1631-70-5; HSiMe₂CH₂Cl, 3144-74-9; HSiMe₂Cl, 1066-35-9; HSiMeCl₂, 75-54-7; HSi(SiMe₃)₃, 1873-77-4; HSiMe₂(OSiMe₃), 1438-82-0; DSiEt₃, 1631-33-0; DSiMe₂Ph, 22034-19-1; DSiPh₃, 18536-60-2; DSi(^tPr)₃, 139408-37-0; HDSiPh₂, 126971-98-0; D₂SiPh₂, 17950-94-6; AnPhCH⁺, 13270-92-3; An₂CH⁺, 13948-07-7; An(*p*-PhOC₆H₄)CH⁺, 99796-93-7; AnTolCH⁺, 42289-58-7; Ph(*p*-PhOC₆H₄)CH⁺, 104910-81-8; Tol₂CH⁺, 58493-75-7; PhTolCH⁺, 42289-57-6; MeSiH₂⁺, 27516-00-3; (H₂N)SiH₂⁺, 66639-71-2; (HO)SiH₂⁺, 66639-72-3; FSiH₂⁺, 68992-43-8; (H₃Si)SiH₂⁺, 86860-82-4; (H₂P)SiH₂⁺, 80401-42-9; (HS)SiH₂⁺, 80401-43-0; ClSiH₂⁺, 80401-44-1; MeCH₂⁺, 14936-94-8; (H₂N)CH₂⁺, 54088-53-8; (HO)CH₂⁺, 17691-31-5; FCH₂⁺, 35310-31-7; (H₃Si)CH₂⁺, 53696-43-8; (H₂P)CH₂⁺, 111129-73-8; (HS)CH₂⁺, 20879-50-9; ClCH₂⁺, 59000-00-9.

Supplementary Material Available: Tables with concentrations and rate constants of the kinetic experiments described in Tables I, II, V, VI, and VII (19 pages). Ordering information is given on any current masthead page.