

Chemische Berichte

Inhalt

Contents

A

Teil A: Anorganische und elementorganische Chemie / Part A: Inorganic and Organometallic Chemistry

Kreiter, C. G., Lehr, K.,
Leyendecker, M., Sheldrick, W. S.,
Exner, R.

Westermann, H., Nieger, M.,
Niecke, E.

Herberich, G. E., Englert, U.,
Hostalek, M., Laven, R.

Herberich, G. E., Negele, M.,
Ohst, H.

Erhart, M., Mews, R., Pauer, F.,
Stalke, D., Sheldrick, G. M.

Maggiulli, R., Mews, R.,
Noltemeyer, M., Offermann, W.,
Paape, R., Stohrer, W.-D.

Habben, C. D., Heine, A.,
Sheldrick, G. M., Stalke, D.,
Bühl, M., von R. Schleyer, P.

Boese, R., Haas, A.,
Spehr, M.

Piana, H., Wagner, H.,
Schubert, U.

Baudler, M., Arndt, V.

Jutzi, P., Kroos, R.,
Müller, A., Bögge, H.,
Penk, M.

Notizen / Notes

Seth, N., Gupta, V. D.,
Linti, G., Nöth, H.

Bittner, J., Seppelt, K.

- 3 ► Photochemische Reaktionen von Übergangsmetall-Organyl-Komplexen mit Olefinen, 3. – [4 + 5]-Cycloaddition konjugierter Diene an Tricarbonyl(η^5 -2,4-dimethyl-2,4-pentadienyl)mangan
Photochemical Reactions of Transition Metal Organyl Complexes with Olefins, 3. – [4 + 5] Cycloaddition of Conjugated Dienes to Tricarbonyl(η^5 -2,4-dimethyl-2,4-pentadienyl)manganese
- 13 ► Synthese und Struktur von Bis[bis(diisopropylamino)phosphino]chalkogeniden
Synthesis and Structure of Bis[bis(diisopropylamino)phosphino]chalcogenides
- 17 ► Derivate des Borols, XVI. – Bis(borol)-nickel-Komplexe
Derivatives of Borole, XVI. – Bis(borol)-nickel Complexes
- 25 ► Derivate des Borols, XVII. – { η^5 -[1-(Diisopropylamino)borol]}metall-Komplexe: Synthesen, Protonierung, interne Rotation
Derivatives of Borole, XVII. – { η^5 -[1-Diisopropylamino]borole}metal Complexes: Syntheses, Protonation, Internal Rotation
- 31 ► (Perfluoroalkyl)(dimethylamino)sulfonium-hexafluoroarsenate
(Perfluoroalkyl)(dimethylamino)sulfonium Hexafluoroarsenates
- 39 ► Addition von Alkenen an 5-(Trifluormethyl)-1,3,2,4,6-dithiatriazin
Addition of Alkenes to 5-(Trifluoromethyl)-1,3,2,4,6-dithiatriazine
- 47 ► Synthese und Eigenschaften von 1,2,4,3-Thiadiazaboretidinen. Kristallstruktur des 2,4-Di-*tert*-butyl-3-phenyl-1,2,4,3-thiadiazaboretidins
Synthesis and Properties of 1,2,4,3-Thiadiazaboretidines. Crystal Structure of 2,4-Di-*tert*-butyl-3-phenyl-1,2,4,3-thiadiazaboretidine
- 51 ► Darstellung von (Perfluoroalkyl)halogeno-1,3-diselenetanen aus entsprechenden Selenocarbonylfluoriden und Umsetzung mit Bortrichlorid sowie Arsenpentaffluorid
Preparation of (Perfluoroalkyl)halogeno-1,3-diselenetanes from the Corresponding Selenocarbonyl Fluorides and Reactions with Boron Trichloride or Arsenic Pentaffluoride
- 63 ► Übergangsmetall-Silyl-Komplexe, 36. – Zur Frage von Gold-Gold-Wechselwirkungen bei $\text{Ph}_2\text{P}(\text{CH}_2)_{12}\text{PPh}_2$ -verbrückten zweikernigen Gold-Silyl-Komplexen
Transition-Metal Silyl Complexes, 36. – On the Question of Gold-Gold Interactions in $\text{Ph}_2\text{P}(\text{CH}_2)_{12}\text{PPh}_2$ -Bridged Dinuclear Gold Silyl Complexes
- 69 ► Beiträge zur Chemie des Phosphors, 210. – Pentaorganyltricyclonaphosphane mit Brexan-Struktur, $b\text{-P}_9\text{R}_5$ ($\text{R} = \text{Me}, \text{Et}, \text{iPr}$)
Contributions to the Chemistry of Phosphorus, 210. – Pentaorganyltricyclonaphosphanes with a Structure Analogous to Brexane, $b\text{-P}_9\text{R}_5$ ($\text{R} = \text{Me}, \text{Et}, \text{iPr}$)
- 75 ► Synthese und Struktur der Polycyclohexaphosphane $\text{P}_6(\text{C}_5\text{Me}_5)_4$ und $\text{P}_6(\text{C}_5\text{Me}_5)_2$
Synthesis and Structure of the Polycyclohexaphosphines $\text{P}_6(\text{C}_5\text{Me}_5)_4$ and $\text{P}_6(\text{C}_5\text{Me}_5)_2$
- 83 Untersuchungen über das Produkt aus Monobutylzinn(IV)-Verbindungen und Benzoyl(thiobenzoyl)methan: ein $1,3,2\lambda^5,4\lambda^5$ -Dithiadistannetan
► Investigation on the Product from Monobutyltin(IV) Compounds and Benzoyl-thiobenzoyl)methane: a $1,3,2\lambda^5,4\lambda^5$ -Di-thiadistannetane
- 87 Molekülstruktur von $\text{Hg}[\text{C}(\text{COF})=\text{SF}_2=\text{O}]_2$
► Molecular Structure of $\text{Hg}[\text{C}(\text{COF})=\text{SF}_2=\text{O}]_2$

► Publikationssprache

► Language of Publication

Inhalt (Fortsetzung)

- Kuhn, N., Jendral, K.,
Boese, R., Bläser, D.
- Wiberg, N., Schuster, H.
- 93 ► Zur Kenntnis des „Silanphosphimins“
(Phosphasilaethens) $t\text{Bu}_2\text{Si} = \text{PSi}t\text{Bu}_3$

Kurzmitteilung / Short Communication

- Sommovigo, M., Pasquali, M.,
Leoni, P., Braga, D.,
Sabatino, P.
- 97 ► P – C-Bindungsaktivierung und η^4 -Koordinierung von Aren: Röntgen-Kristallstruktur eines zweikernigen μ -Phosphido- μ - η^2 : η^2 -Phenoxy-Zwitterionenkomplexes von Palladium, der ein Aggregat von drei Wasserstoff-verbrückten Phenol-Molekülen abfängt

Querverweise / Cross References

- Bickelhaupt, F., Maier, G. et al. 185 Massenspektrometrie und Matrix-Isolierung einiger 9-Silaanthracene
- Eilbracht, P. et al. 191 ► Umwandlungen des (+)-2- und (+)-3-Carens in optisch aktive Siebenring-Systeme
- Aumann, R. et al. 213 ► Organische Synthesen mit Übergangsmetall-Komplexen, 50

Contents (Continued)

- Heterocycles as Ligands, X. – 2,2',5,5'-Tetra-*tert*-butyl-1,1'-diazaferrrocene – Stabilization of the Diheterometallocene Structure by Steric Shielding
- On the Knowledge of the “Silanephosphimine” (Phosphasilaethene) $t\text{Bu}_2\text{Si} = \text{PSi}t\text{Bu}_3$

- P – C Bond Activation and η^4 -Coordination of Arene: X-ray Crystal Structure of a Dinuclear μ -Phosphido μ - η^2 : η^2 -Phenoxy Zwitterionic Complex of Palladium Trapping an Aggregate of Three Hydrogen-Bonded Phenol Molecules

B

Teil B: Organische Chemie / Part B: Organic Chemistry

- Hoffmann, V. T., Musso †, H. 103 ► Asterane, XXII. – Synthese eines doppelten Tetraasterans: Nonacyclo[10.8.0.0^{2,11}.0^{4,9}.0^{6,17}.0^{7,16}.0^{9,14}.0^{14,19}]eicosan
- Korbonits, D., Simon, K., Kolonits, P. 111 Synthese von 4-Aminopyrimidinen aus 1,2,4-Oxadiazolen, III. – Eine neuartige Bildung von Olefinen aus Aminen
- Reißig, H.-U., Hippeli, C. 115 Lithiierte 5,6-Dihydro-4H-1,2-oxazine: Synthese, hoch diastereoselektive Reaktionen mit Elektrophilen und nachfolgende Umwandlungen
- Schulz, A., Kaim, W. 129 ► Elektrochemische und spektroskopische Charakterisierung von *N,N'*-Dialkylchinonoxalium-Redoxsystemen
- Prüssé, T., Czekay, G., Schwarz, H. 141 Wird die Fe⁺-induzierte C – H/C – C-Bindungsaktivierung von α,ω -Alkanedinitrilen durch bifunktionelle Wechselwirkungen beeinflusst?
- Godt, A., Schlüter, A.-D. 149 ► Cyclobutene-Ringöffnung: Eine nützliche Reaktion zur Synthese doppelsträngiger Moleküle
- Effenberger, F., Streicher, W. 157 ► Darstellung aromatischer Fluor-Verbindungen durch nucleophilen Austausch von Nitro-Gruppen gegen Fluorid
- Effenberger, F., Koch, M., Streicher, W. 163 ► Nucleophile Substitution von Nitrit in Nitrobenzolen, Nitrobiphenylen und Nitronaphthalinen
- Öhler, E., Zbiral, E. 175 ► Zur Reaktion von Dialkylphosphiten mit α -Enonen, I. – Synthese und Allyl-Umlagerung von (1-Hydroxy-2-alkenyl)- und (1-Hydroxy-2-cycloalkenyl)phosphonsäuredimethylestern
- van den Winkel, Y., van Baar, B. L. M., Bickelhaupt, F., Kulik, W., Sierakowski, C., Maier, G. 185 Neutralisations-Reionisations-Massenspektrometrie und Matrix-Isolierung einiger 9-Silaanthracene
- Publikationssprache
- Mass Spectrometry and Matrix Isolation of Some 9-Silaanthracenes
- Conversions of (+)-2- and (+)-3-Carene into Optically Active Seven-Membered Ring Systems
- Organic Syntheses via Transition Metal Complexes, 50
- Asteranes, XXII. – Synthesis of a Double Tetraasterane: Nonacyclo[10.8.0.0^{2,11}.0^{4,9}.0^{6,17}.0^{7,16}.0^{9,14}.0^{14,19}]eicosane
- Synthesis of 4-Aminopyrimidines from 1,2,4-Oxadiazoles, III. – A Novel Type of Formation of Olefins from Amines
- Lithiated 5,6-Dihydro-4H-1,2-oxazines: Synthesis, Highly Diastereoselective Reactions with Electrophiles, and Subsequent Transformations
- Electrochemical and Spectroscopic Characterization of *N,N'*-Dialkylquinoxalinium Redox Systems
- Is the Fe⁺-Induced C – H/C – C Bond Activation of α,ω -Alkanedinitriles Affected by Bifunctional Interactions?
- Cyclobutene Ring Opening: A Useful Reaction for the Synthesis of Double-Stranded Molecules
- Synthesis of Aromatic Fluoro Compounds by Nucleophilic Exchange of Nitro Groups by Fluoride
- Nucleophilic Substitution of Nitrite in Nitrobenzenes, Nitrobiphenyls, and Nitronaphthalenes
- Reaction of Dialkyl Phosphites with α -Enones, I. – Synthesis and Allylic Rearrangement of Dimethyl (1-Hydroxy-2-alkenyl)- and (1-Hydroxy-2-cycloalkenyl)-phosphonates
- Neutralization-Reionization Mass Spectrometry and Matrix Isolation of Some 9-Silaanthracenes
- Language of Publication

Inhalt (Fortsetzung)

- Eilbracht, P., Winkels, I.
- Bestmann, H. J., Röder, T., Bremer, M., Löw, D.
- Mayr, H., Heilmann, W., Bäuml, E., Vorbrüggen, H.
- Victory, P., Alvarez-Larena, A., Beti, C., Borrell, J. I., Batllori, X., Córdoba, C.
- Aumann, R., Hinterding, P.
- Enders, D., Nakai, S.
- Adam, W., Hadjiarapoglou, L., Smerz, A.
- Gast, A., Breitmaier, E.
- Kirmse, W., Mönch, D.
- Kirmse, W., Mrotzek, U., Siegfried, R.
- Schinzer, D., Ruppelt, M.
- 3
- 39
- 191 ► Stereospezifische Umwandlungen des (+)-2- und (+)-3-Carens in optisch aktive Siebenring-Systeme
- 199 ► Bororganyle, II. — Darstellung, Struktur und Reaktionen von Lithium-[cyan-(triphenylphosphoranylidien)methyl]trihydroborat(-1)
- 203 ► Synthese von 2,2,5,5-Tetramethylcyclopentacarbonsäure — Ein Baustein eines Süßstoffes auf Aminosäure-Basis
- 207 ► Trioxypentafulvene, 2. — Synthese 1-substituierter 2,3,6-Trioxypentafulvene
- 213 ► Organische Synthesen mit Übergangsmetall-Komplexen, 50. — 2H-Pyrrol-Komplexe durch stufenweise Insertion von Alkinen und Isocyaniden in $M=C$ -Bindungen von Chrom- und Wolfram-Carbonylkomplexen
- 219 ► Regio-, diastereo- und enantioselektive Synthese von *vic*-Diolen über α -Silylketone nach der SAMP/RAMP-Hydrazone-Methode
- 227 ► Dioxiran-Epoxidierung von α,β -ungesättigten Ketonen
- 233 ► Neue O₆-Makrocyclen mit 2,2-Bis(2-furyl)-propan-Einheiten
- 237 ► Umlagerungen von 1,4,4- und 2,2,5-Trimethylbicyclo[3.2.1]oct-6-yl-Kationen
- 241 ► Umlagerungen von 5,5,6,6-Tetraalkyl-2-norbornyl-Kationen
- 247 ► Säure-katalysierte und Lewis-Säure-be schleunigte Synthese von 6-Hydroxybicyclo[4.3.0]nonan-3-onen

Contents (Continued)

- Stereospecific Conversions of (+)-2- and (+)-3-Carens into Optically Active Seven-Membered Ring Systems
- Organoboron Compounds, II. — Preparation, Structure, and Reactions of Lithium [Cyanotriphenyl]phosphoranylidene)methyl]trihydroborate(-1)
- Synthesis of 2,2,5,5-Tetramethylcyclopentanecarboxylic Acid — A Building Block of an Amino Acid Based Sweetener
- Trioxypentafulvenes, 2. — Synthesis of 1-Substituted 2,3,6-Trioxypentafulvenes
- Organic Syntheses via Transition Metal Complexes, 50. — 2H-Pyrrole Complexes by a Stepwise Insertion of Alkynes and Isocyanides into $M=C$ Bonds of Carbene Chromium or Tungsten Complexes
- Regio-, Distereco-, and Enantioselective Synthesis of *vic*-Diols via α -Silyl Ketones According to the SAMP/RAMP Hydrazone Method
- Dioxirane Epoxidation of α,β -Unsaturated Ketones
- New O₆ Macrocycles with 2,2-Bis(2-furyl)-propane Units
- Rearrangements of 1,4,4- and 2,2,5-Trimethylbicyclo[3.2.1]oct-6-yl Cations
- Rearrangements of 5,5,6,6-Tetraalkyl-2-norbornyl Cations
- Acid-Catalyzed and Lewis Acid-Promoted Synthesis of 6-Hydroxybicyclo[4.3.0]nonan-3-ones

Kurzmitteilung / Short Communication

- Schinzer, D., Ruppelt, M.
- 247 ► Säure-katalysierte und Lewis-Säure-be schleunigte Synthese von 6-Hydroxybicyclo[4.3.0]nonan-3-onen

Querverweise / Cross References

- Kreiter, C. G. et al.
- Mews, R. et al.
- 3 ► Photochemische Reaktionen von Übergangsmetall-Organyl-Komplexen mit Olefinen, 3
- 39 ► Addition von Alkenen an 5-(Trifluormethyl)-1,3,2,4,6-dithiatriazin

A I Richtlinien für Autoren

- Publikationssprache

Photochemical Reactions of Transition Metal Organyl Complexes with Olefines, 3

Addition of Alkenes to 5-(Trifluoromethyl)-1,3,2,4,6-dithiatriazine

B I Instructions for Authors

- Language of Publication

Synthesis of 2,2,5,5-Tetramethylcyclopentanecarboxylic Acid – A Building Block of an Amino Acid Based Sweetener

Herbert Mayr^a, Werner Heilmann^a, Englbert Bäuml^a, and Helmut Vorbrüggen^b

Institut für Chemie, Medizinische Universität zu Lübeck^a,
Ratzeburger Allee 160, D-2400 Lübeck 1

Schering AG^b,
Postfach 650311, D-1000 Berlin 65


Received May 29, 1990

Key Words: Sweetener / Electrophilic addition / Cycloalkanecarboxylic acid

The title compound **2** is synthesized in three steps from readily available 2,5-dichloro-2,5-dimethylhexane (**8**) and vinylidene


chloride (**9**). The key steps are carbocationic addition and cyclization reactions.

N-L-Aspartyl-*N'*-(alkylcarbonyl)-1,1-diaminoethanes **1** have been reported to be thermally stable, water-soluble sweeteners without bitter after-taste¹⁾. The sweetness of **1** strongly depends on the nature of the acyl group R, and among 14 tested compounds (**1**), the maximal taste intensity has been found for R = 2,2,5,5-tetramethylcyclopentylcarbonyl (1000-fold sweetness relative to sucrose).




The incorporated carboxylic acid **2** has previously been synthesized by permethylation of the α positions of cyclopentanone, Wittig reaction, hydroboration, and Cr(VI) oxidation as shown in Scheme 1^{1a)}.

Scheme 1




Since compounds with quaternary carbons are often readily accessible via carbocationic addition reactions²⁾, we have attempted to improve the synthesis of **2** by employing the carbocationic cyclization of **3** as the key step³⁾.



For the synthesis of precursors of **3**, two different approaches, indicated by the disconnections *a* and *b* in formula **3**, have been considered.

Disconnection a: The aldehyde **6** has previously been prepared in 88% yield from the anion of isobutyraldehyde (**5**) and prenyl bromide⁴⁾. Since in this procedure, the enolate anion is generated with potassium hydride in THF, we have preferred to combine compounds **4** and **5** under phase-transfer conditions as shown in Scheme 2, making use of a procedure previously described for the synthesis of similar compounds⁵⁾. Treatment of **6** with triphenylphosphine/tetrachloromethane⁶⁾ gives the diene **7** in good yield.

Scheme 2



Disconnection b yields a C₈ fragment, which is technically produced from acetylene and acetone. The commercially available 2,5-dimethyl-2,5-hexanediol is quantitatively converted into the dichloride **8**⁷⁾ by treatment with hydrochloric acid. Attempts to produce the five-membered ring in a one-pot reaction from **8** have not been successful. While reactions of **8** with vinylsilanes have not been considered for economic reasons, the reaction of **8** with propyne gives complex mixtures of products under a variety of conditions^{3b)}. When **8** and **9** are combined in sulfuric acid (Bott-Hellmann conditions⁸⁾), a diacid is formed from **8** and two equivalents of vinylidene chloride **9**.

Since **8** incorporates two isolated equivalent functional groups, we have not succeeded in selectively engaging a single C–Cl bond in the reaction with vinylidene chloride. Mild Lewis acids (ZnCl₂) do not initiate the reaction of **8** with **9**, while AlCl₃, EtAlCl₂, or FeCl₃ produce mixtures of

the 1:1 adduct **10** and the 2:1 adduct **11**. Among these Lewis acids, AlCl_3 gives rise to the highest yields of **10**.





Figure 1. Product composition during the AlCl_3 -promoted reaction of **8** (50.0 mmol) with **9** (325 mmol) at 0°C. Since the reaction takes place only within the first few minutes after the addition of the catalyst, this figure refers to a single experiment, in which the AlCl_3 portions have been added at 30-min intervals.


Though vinylidene chloride is usually employed as the solvent, the AlCl_3 -initiated reactions terminate after a certain degree of conversion due to deactivation of the Lewis acid. In accord with this rationalization, a progress of the reactions can be achieved by adding more aluminum chloride. The degree of conversion thus depends on the percentage of Lewis acid, and Figure 1 shows that optimal yields of **10** are obtained with 8–10% of AlCl_3 . Larger amounts of the Lewis acid give rise to the formation of a higher percentage of **11** and of unidentified nonvolatile compounds. Since the ratio of **10/11** decreases with increasing degree of conversion, one can improve the yield of the process by using a smaller amount of Lewis acid and recovering the nonconverted reactants **8** and **9**.



Treatment of **10** with a base under phase-transfer conditions ($\text{NaOH}/\text{CH}_2\text{Cl}_2$) gives mixtures of **12** and **13**; on

heating, pure **13** is formed in 79% yield. Since the cyclization of **13** appears to be easier than the cyclization of **12**, no efforts have been made to selectively produce **12**.


Cyclizations: In analogy to investigations by Bott⁹ and Gallo¹⁰, we tried to achieve the conversion of **13** into **2** in sulfuric acid. Compound **2** is obtained under these conditions, but generally only in low yield (<30%). Therefore, the cyclization has been carried out in boiling formic acid, conditions previously employed by Lansbury for analogous conversions¹¹. In this medium, like in 70–95% aqueous acetic acid, **7** as well as **13** are cyclized to give 50–60% of **2**, but we have not found conditions which avoid the concomitant formation of the six-membered ring compounds **14** and/or **15**.



Compound **15** is formed in 95% yield, however, when **7** is treated with FeCl_3 in CH_2Cl_2 . Possibly, this conversion is due to proton catalysis (traces of moisture) since the selective rearrangement of **7** to **15** is also catalyzed by FSO_3H .

The mechanism of these reactions has not been elucidated, and Scheme 3, which is self-explanatory, only represents a rationalization of the isolated products. Probably, thermodynamic product control is responsible for the exclusive formation of **15** from **7** in the absence of water: Under these

Scheme 3



conditions, irreversible trapping of the cyclopentyl-substituted carbenium ion **18** is not possible, so that the less strained six-membered ring compound **15** is produced selectively. It is remarkable that compound **15** is also generated from **13** in formic acid; thus, one has to assume the presence of an ion pair [**20** Cl⁻] which collapses to give the dichloride **15**.

The dependence of the product composition on the reaction conditions indicates the occurrence of mutual rearrangements of the intermediate carbenium ions, so that the product ratio **2**/**(14 + 15)** does not provide information on the relative rates of the cyclization steps (*6-endo* or *5-exo*¹²).

Conclusion: The sequence **8** → **10** → **13** → **2** leads to the title compound **2** in a three-step synthesis in 32% overall yield. Since only inexpensive reagents are needed, which are combined under convenient reaction conditions, the carbocationic approach to **2** is superior to the carbanionic route described in Scheme 1. Alternative routes¹³, which have been published after the submission of our first report^{3a}, also employ 2,2,5,5-tetramethylcyclopentanone as starting compound and, in our opinion, are less straightforward than the procedures reported in this article.

Experimental

NMR: XL 200 (Varian), internal standard TMS. — Mass spectra: 70–250 E (VG-Instruments). — IR: IR-435 (Shimadzu).

2,2,5-Trimethyl-4-hexenal (6): A solution of **4** (13.6 g, 130 mmol) and **5** (7.21 g, 100 mmol) in toluene (10 ml) is added dropwise (10 min) to a vigorously stirred mixture of 50% aqueous NaOH (30 g), toluene (70 ml), and tetrabutylammonium iodide (1.4 g, 3.8 mmol) at 80°C. The mixture is stirred for 4.5 h (80°C), cooled and poured onto water. The product is extracted with ether, and the ethereal extract dried with Na₂SO₄. Distillation over a Vigreux column yields **6** (11.9 g, 85%) as a colorless liquid with b.p. 89–92°C/71 mbar (ref.¹⁴: 65–66°C/20 Torr; purity 94% by GC). Spectral data are given in ref.¹⁴.

1,1-Dichloro-3,3,6-trimethyl-1,5-heptadiene (7): A solution of **6** (5.00 g, 35.7 mmol) in CCl₄ (5 ml) is added dropwise (5 min) with stirring to a mixture of triphenylphosphine (21.0 g, 80.0 mmol) in CCl₄ (40 ml) at 80°C (N₂ atmosphere). After 3.5 h, the mixture is cooled and poured onto 300 ml of hexane to precipitate triphenylphosphine and triphenylphosphine oxide. The suspension is filtered over silica gel and the resulting solution evaporated to give **7** (6.12 g, 83%), as a colorless oil, which is used without further purification. Bulb-to-bulb distillation [34–37°C (bath)/0.15 mbar] yields an analytically pure product. — ¹H NMR (CDCl₃): δ = 1.16 (s, 6 H, 3-CH₃), 1.62, 1.73 (2 br. s, 6 H, 6-CH₃, 7-H), 2.14 (br. d, *J* = 7.6 Hz, 2 H, 4-H), 5.13 (br. t, *J* = 7.6 Hz, 1 H, 5-H), 5.87 (s, 1 H, 2-H). — ¹³C NMR (CDCl₃): δ = 17.95, 26.04 (2 q, 6-CH₃, C-7), 26.70 (q, 3-CH₃), 37.66 (s, C-3), 40.52 (t, C-4), 118.83 (s, C-1), 120.21 (d, C-5), 134.10 (s, C-6), 137.93 (d, C-2). — IR (neat): ν = 2956, 2917, 1606, 1466, 1450, 1383, 1376, 879 cm⁻¹. — MS (70 eV): *m/z* (%) = 208, 206 (0.1, 0.15) [M⁺], 141, 139, 137 (3, 17, 26), 118, 116 (6, 8), 69 (100), 41 (47).

C₁₀H₁₆Cl₂ (207.1) Calcd. C 57.98 H 7.79
Found C 57.97 H 7.84

2,5-Dichloro-2,5-dimethylhexane (8): 2,5-Dimethyl-2,5-hexanediol (73.1 g, 0.500 mol) is stirred with 37% aqueous HCl (250 ml) for 1 h. The initially biphasic mixture first becomes homogeneous, and

later on the precipitation of crystalline **8** takes place. The product is extracted with 600 ml of petroleum ether (b.p. 40–60°C) and dried with CaCl₂. Evaporation of the solvent yields 81.9 g (89%) of an NMR-spectroscopically pure solid, which is recrystallized from petroleum ether: m.p. 68–68.5°C (ref.⁷: 64°C). — ¹H NMR (CCl₄): δ = 1.55 (s, 12 H, CH₃), 1.86 (s, 4 H, CH₃). — ¹³C NMR (CDCl₃): δ = 32.51 (q), 41.11 (t), 70.22 (s).

Reaction of 8 with Vinylidene Chloride (9): Aluminum chloride (3.75 g, 28.0 mmol) is added in portions during 45 min to a well-stirred suspension of **8** (54.9 g, 300 mmol) in **9** (135 ml, 1.70 mol) at 0°C. After stirring for another 100 min at 0°C, 10 ml of water and 20 g of CaCl₂ are added successively, and the mixture is filtered. Evaporation of **9** leaves 88.4 g of a dark viscous residue from which 17.5 g (95.6 mmol) of **8** are recovered by distillation. Distillation of the residue yields 34.3 g of **10** (60% with respect to consumed **8**) and 26.2 g of **11** (34% with respect to consumed **8**) which solidifies at –20°C.

1,1,1,6-Tetrachloro-3,3,6-trimethylheptane (10): B.p. 79–80°C/0.5 mbar. — ¹H NMR (CDCl₃): δ = 1.20 (s, 6 H, CH₃), 1.59 (s, 6 H, CH₃), 1.59–1.83 (m, 4 H, CH₂-CH₂), 2.81 (s, 2 H, CH₂-CCl₃). — ¹³C NMR (CDCl₃): δ = 27.80 (q), 32.43 (q), 35.69 (s), 38.38 (t), 39.97 (t), 63.25 (t), 70.74 (s), 98.60 (s). — IR (neat): ν = 2961, 2931, 2870, 1470, 1452, 1387, 1370, 1100, 896, 790, 726, 703, 605 cm⁻¹. — MS (70 eV): *m/z* (%) = 229, 227 (0.9, 1.3) [M⁺ – HCl – CH₃], 171 (4), 147 (7), 137 (13), 123 (10), 111 (50), 77 (64), 69 (83), 31 (100).

C₁₀H₁₈Cl₄ (280.1) Calcd. C 42.89 H 6.48
Found C 43.06 H 6.54

1,1,1,8,8,8-Hexamchloro-3,3,6,6-tetramethyloctane (11): Colorless needles with m.p. 52.5–53.5°C (petroleum ether). — ¹H NMR (CDCl₃): δ = 1.18 (s, 12 H, CH₃), 1.46 (s, 4 H, CH₂-CH₂), 2.81 (s, 4 H, CH₂-CCl₃). — ¹³C NMR (CDCl₃): δ = 27.79 (q), 35.91 (s), 37.16 (t), 63.32 (t), 98.67 (s). — IR (KBr): ν = 2976, 2953, 2904, 2870, 1472, 1465, 1390, 1373, 1337, 1255, 1188, 1154, 1123, 1042, 903, 784, 717, 608 cm⁻¹. — MS (70 eV): *m/z* (%) = 344, 342, 340, 338, (0.07, 0.18, 0.29, 0.18) [M⁺ – HCl], 229, (1.3), 227 (1.4), 193 (1.0), 191 (1.2), 173 (3), 171 (7), 149 (23), 147 (70), 137 (19), 111 (81), 77 (100), 69 (36). C₁₂H₂₀Cl₆ (377.0) Calcd. C 38.23 H 5.35
Found C 38.38 H 5.37

1,6-Dichloro-3,3,6-trimethyl-1-heptyne (13): Compound **10** (25.2 g, 90.0 mmol) and 0.60 g of tricaprylmethylammonium chloride (Aliquat 336) are stirred vigorously with 40% aqueous NaOH (50 ml) at 80°C for 22 h. After cooling, 60 ml of CH₂Cl₂ is added, and the organic layer is dried with CaCl₂. The solvent is evaporated, and the residue is distilled to give 14.7 g (79%) of **13** with b.p. 39–43°C/1 mbar. — ¹H NMR (CDCl₃): δ = 1.21 (s, 6 H, CH₃), 1.59 (s, CH₃), 1.53–1.93 (m, 10H including the s at δ 1.59, CH₂-CH₂). — ¹³C NMR (CDCl₃): δ = 28.98 (q), 31.26 (s), 32.49 (q), 38.25 (t), 41.48 (t), 57.90 (s), 70.66 (s), 75.80 (s). — IR (neat): ν = 2967, 2920, 2864, 2220, 1465, 1450, 1385, 1370, 1267, 1098, 989, 788 cm⁻¹. — MS (70 eV): *m/z* (%) = 208, 206 (0.4, 0.6) [M⁺], 155 (28), 135 (20), 115 (15), 103 (33), 101 (100), 77 (16), 69 (74), 65 (28).

C₁₀H₁₆Cl₂ (207.1) Calcd. C 57.98 H 7.79
Found C 58.29 H 7.67

Cyclization of 13: Compound **13** (8.29 g, 40.0 mmol) is added dropwise with stirring to 80 ml of refluxing 90% aqueous formic acid (30 min) and heated under reflux for 6 h. Water (300 ml) is added after cooling, and the mixture is extracted with two 50-ml portions of petroleum ether. The combined organic layers are extracted with five 30-ml portions of 10% aqueous Na₂CO₃ solution. The aqueous layers are combined and acidified with conc. HCl to precipitate 4.63 g (68%) of spectroscopically pure **2**.

The remaining petroleum ether solution is dried with CaCl_2 , and the solvent is evaporated to give 2.07 g of a brown liquid residue which is separated by chromatography (silica gel, petroleum ether/ CH_2Cl_2) to give 574 mg (7%) of **15** and 1.02 g (17%) of **14**.

2,2,5,5-Tetramethylcyclopentanecarboxylic Acid (2): M.p. 128–129°C (petroleum ether). — ^1H NMR (CDCl_3): δ = 1.15 (s, 6H, 2,5- CH_3), 1.16 (s, 6H, 2,5- CH_3), 1.51–1.63 (m, 4H, 3,4-H), 2.26 (s, 1H, 1-H). — ^{13}C NMR (CDCl_3): δ = 25.42, 31.41, (2 q, CH_3), 40.80 (t, C-3,4), 42.56 (s, C-2,5), 64.20 (d, C-1), 179.26 (s, CO_2H). — IR (KBr): $\tilde{\nu}$ = 2938, 2860, 2724, 2325, 1691, 1459, 1409, 1379, 1362, 1274, 1253, 1230, 1204, 935, 730 cm^{-1} . — MS (70 eV): m/z (%) = 170 (0.21) [M^+], 155 (1.8), 137 (1.5), 114 (2.4), 110 (4), 109 (10), 102 (5), 101 (100), 83 (8), 70 (32), 55 (17).

$\text{C}_{10}\text{H}_{18}\text{O}_2$ (170.3) Calcd. C 70.55 H 10.66
Found C 70.32 H 10.51

2,3,6,6-Tetramethyl-2-cyclohexen-1-one (14): B.p. 30–40°C (bath)/0.6 mbar. — ^1H NMR (CDCl_3): δ = 1.04 (s, 6H, 6- CH_3), 1.69 (t, J = 7.4 Hz, 2H, 5-H), 1.85 (br. s, 3H, 2- CH_3), 2.24 (t, J = 2 Hz, 3H, 3- CH_3), 2.50 (m, 2H, 4-H). — ^{13}C NMR (CDCl_3): δ = 20.41 (q), 24.07 (q, 6- CH_3), 24.43 (q), 25.27, 34.77 (2 t, C-4,5), 46.48 (s, C-6), 130.12, 148.28 (2s, C-2,3), 211.22 (s, C-1). — IR (neat): $\tilde{\nu}$ = 2951, 2850, 1706 (s, $\text{C}=\text{O}$), 1631 (s, $\text{C}=\text{C}$), 1450, 1378, 1364, 1270, 1192, 1114, 989, 849, 786, 762 cm^{-1} . — MS (70 eV): m/z (%) = 152 (78) [M^+], 137 (93), 109 (25), 96 (31), 81 (12), 68 (100), 67 (36).

$\text{C}_{10}\text{H}_{16}\text{O}$ (152.2) Calcd. C 78.90 H 10.59
Found C 79.50 H 10.75

Cyclization of 7 in Formic Acid: Compound 7 (1.0 g, 4.8 mmol) is added dropwise within 10 min to a refluxing mixture of 98% formic acid (10 ml) and BF_3 –ether (1 ml). The mixture is heated under reflux for 6 h and worked up as described above to yield 0.40 g (49%) of **2** and 0.20 g (20%) of **15**.

1,2-Dichloro-3,3,6,6-tetramethyl-1-cyclohexene (15): A solution of 7 (210 mg, 1.01 mmol) in 3 ml of anhydrous CH_2Cl_2 is added to a suspension of FeCl_3 (40 mg, 0.25 mmol) in 5 ml of CH_2Cl_2 . During stirring, the mixture adopts a deep red color after approximately 5 min. Pentane (20 ml) is added after 2.5 h, and the mixture is passed through 2.0 g of silica gel. After elution with pentane (ca. 50 ml), the pentane solutions are concentrated to give 200 mg (95%) of spectroscopically (^1H NMR) pure **15**; b.p. 20–30°C (bath)/0.5 mbar. — ^1H NMR (CDCl_3): δ = 1.27 (s, 12H, CH_3), 1.63 (s, 4H, CH_2). — ^{13}C NMR (CDCl_3): δ = 26.13 (q, 3,6- CH_3), 41.00 (t, C-

4,5), 47.00 (s, C-3,6), 155.06 (s, C-1,2). — IR (neat): $\tilde{\nu}$ = 2987, 2950, 2864, 1604, 1588, 1458, 1363, 1216, 1145, 1007, 907, 866, 847, 778 cm^{-1} . — MS (70 eV): m/z (%) = 208, 206 (9, 14) [M^+], 193 (8), 191 (13), 173 (17), 171 (52), 150 (58), 135 (18), 115 (59), 91 (12), 83 (100), 77 (21), 55 (18).

$\text{C}_{10}\text{H}_{16}\text{Cl}_2$ (207.1) Calcd. C 57.98 H 7.79
Found C 58.14 H 7.78

CAS Registry Numbers

2: 96188-82-8 / **4:** 503-60-6 / **5:** 78-84-2 / **6:** 1000-30-2 / **7:** 128600-84-0 / **8:** 6223-78-5 / **9:** 75-35-4 / **10:** 109749-69-1 / **11:** 128600-85-1 / **13:** 109749-70-4 / **14:** 53343-33-2 / **15:** 128600-86-2 / 2,5-dimethyl-2,5-hexanediol: 110-03-2

- ¹⁾ ^{1a)} Cumberland Packing Corporation (M. S. Verlander, W. D. Fuller, M. Goodman, Inv.) European Pat. Appl. 0128654 (1984) [Chem. Abstr. **102** (1985) 185492 c]. — ^{1b)} W. D. Fuller, M. Goodman, M. S. Verlander, J. Am. Chem. Soc. **107** (1985) 5821.
- ²⁾ Reviews: ^{2a)} M. T. Reetz, Angew. Chem. **94** (1982) 97; Angew. Chem. Int. Ed. Engl. **21** (1982) 96. — ^{2b)} H. Mayr in *Selectivities in Lewis Acid-Promoted Reactions*, NATO ASI Series C, (D. Schinzer, Ed.), vol. 289, p. 21, Kluwer Academic Publishers, Dordrecht 1989.
- ³⁾ ^{3a)} Schering AG (H. Vorbrüggen, H. Mayr, W. Heilmann, Inv.) DOS 3534613.2 (24.09.86) [Chem. Abstr. **107** (1987) 115282 p]. — ^{3b)} W. Heilmann, H. Mayr, H. Vorbrüggen FOCUS MHL **4** (1987) 94 [Chem. Abstr. **111** (1989) 57078 w].
- ⁴⁾ P. Groenewegen, H. Kallenberg, A. van der Gen, Tetrahedron Lett. **1978**, 491.
- ⁵⁾ H. K. Dietl, K. C. Brannock, Tetrahedron Lett. **1973**, 1273.
- ⁶⁾ R. Appel, Angew. Chem. **87** (1975) 863; Angew. Chem. Int. Ed. Engl. **14** (1975) 801.
- ⁷⁾ L. R. C. Barclay, E. E. Betts, J. Am. Chem. Soc. **77** (1955) 5735.
- ⁸⁾ K. Bott, Angew. Chem. **92** (1980) 169; Angew. Chem. Int. Ed. Engl. **19** (1980) 171.
- ⁹⁾ K. Bott, H. Hellmann, Angew. Chem. **78** (1966) 932; Angew. Chem. Int. Ed. Engl. **5** (1966) 870.
- ¹⁰⁾ S. Randriamahefa, P. Deschamps, R. Gallo, H. Grangette, Synthesis **1985**, 493.
- ¹¹⁾ P. T. Lansbury, R. C. Stewart, Tetrahedron Lett. **1973**, 1569.
- ¹²⁾ J. E. Baldwin, J. Chem. Soc. Chem. Commun. **1976**, 734.
- ¹³⁾ Kyowa Hakko Kogyo Co. (T. Yamauchi, K. Hattori, S. Ikeda, K. Tamaki, Invs.) European Pat. Appl. 0196461 (1986) [Chem. Abstr. **110** (1989) 172746 d].
- ¹⁴⁾ P. Cresson, Bull. Soc. Chim. Fr. **1964**, 2618.

[184/90]