

ORIGINAL ARTICLE

Prevalence of *BRCA1/2* germline mutations in 21 401 families with breast and ovarian cancer

Karin Kast,^{1,2} Kerstin Rhiem,³ Barbara Wappenschmidt,³ Eric Hahnen,³ Jan Hauke,³ Britta Bluemcke,³ Verena Zarghooni,³ Natalie Herold,³ Nina Ditsch,⁴ Marion Kiechle,⁵ Michael Braun,⁶ Christine Fischer,⁷ Nicola Dikow,⁷ Sarah Schott,^{8,9} Nils Rahner,¹⁰ Dieter Niederacher,¹¹ Tanja Fehm,¹¹ Andrea Gehrig,¹² Clemens Mueller-Reible,¹² Norbert Arnold,¹³ Nicolai Maass,¹³ Guntram Borck,¹⁴ Nikolaus de Gregorio,¹⁵ Caroline Scholz,¹⁶ Bernd Auber,¹⁶ Raymonda Varon-Manteeva,¹⁷ Dorothee Speiser,¹⁸ Judit Horvath,¹⁹ Nadine Lichey,¹⁹ Pauline Wimberger,^{1,2} Sylvia Stark,²⁰ Ulrike Faust,²¹ Bernhard H F Weber,²² Gunter Emons,²³ Silke Zachariae,²⁴ Alfons Meindl,⁵ Rita K Schmutzler,³ Christoph Engel,²⁴ on behalf of the German Consortium for Hereditary Breast and Ovarian Cancer (GC-HBOC)

For numbered affiliations see end of article.

Correspondence to

Karin Kast, Department of Gynecology and Obstetrics, University Hospital Carl Gustav Carus, TU Dresden, Fetscherstraße 74, Dresden 01069, Germany; karin.kast@uniklinikum-dresden.de

Received 28 November 2015

Revised 21 January 2016

Accepted 2 February 2016

Published Online First

29 February 2016

ABSTRACT

Purpose To characterise the prevalence of pathogenic germline mutations in *BRCA1* and *BRCA2* in families with breast cancer (BC) and ovarian cancer (OC) history.

Patients and methods Data from 21 401 families were gathered between 1996 and 2014 in a clinical setting in the German Consortium for Hereditary Breast and Ovarian Cancer, comprising full pedigrees with cancer status of all individual members at the time of first counselling, and *BRCA1/2* mutation status of the index patient.

Results The overall *BRCA1/2* mutation prevalence was 24.0% (95% CI 23.4% to 24.6%). Highest mutation frequencies were observed in families with at least two OCs (41.9%, 95% CI 36.1% to 48.0%) and families with at least one breast and one OC (41.6%, 95% CI 40.3% to 43.0%), followed by male BC with at least one female BC or OC (35.8%; 95% CI 32.2% to 39.6%). In families with a single case of early BC (<36 years), mutations were found in 13.7% (95% CI 11.9% to 15.7%). Postmenopausal unilateral or bilateral BC did not increase the probability of mutation detection. Occurrence of premenopausal BC and OC in the same woman led to higher mutation frequencies compared with the occurrence of these two cancers in different individuals (49.0%; 95% CI 41.0% to 57.0% vs 31.5%; 95% CI 28.0% to 35.2%).

Conclusions Our data provide guidance for healthcare professionals and decision-makers to identify individuals who should undergo genetic testing for hereditary breast and ovarian cancer. Moreover, it supports informed decision-making of counselees on the uptake of genetic testing.

probability of finding a deleterious germline mutation in a woman affected with BC or OC depends on her familial cancer history in terms of type, number and ages of onset of these cancers.^{2–5} To date, the decision to perform mutation testing is mainly guided by the presence of a family cancer history, which is indicative for a *BRCA* mutation with a certain probability.^{6–8} Consequently, to define appropriate clinical selection criteria, precise knowledge on expected mutation frequencies for individual family histories is required. Since the discovery of the *BRCA1* and *BRCA2* genes, several studies have analysed the relationship between family history and *BRCA* mutation prevalence.^{9–14} In the largest study so far, Frank *et al*¹¹ correlated familial disease histories of 10 000 consecutively enrolled individuals with mutation status, resulting in detailed tabulations of empiric mutation prevalences.

In 1996, the German Consortium for Hereditary Breast and Ovarian Cancer (GC-HBOC) had established a panel of clinical criteria for genetic testing of individuals in a clinical setting, based on familial BC and OC history.⁷ By the end of 2014, a total of 21 401 families suspected of having a deleterious *BRCA* mutation according to this panel of clinical criteria were enrolled into the central registry of GC-HBOC. Based on this data set, we aimed to comprehensively analyse the correlation of family history of BC and OC with *BRCA* mutation frequencies. We were particularly interested in the predictive value of the number of premenopausal versus postmenopausal BCs in the family, and the presence of premenopausal or postmenopausal bilateral breast cancer (bBC) versus unilateral cancer. With this analysis, we aim to provide guidance for counsellors, counselees and decision-makers on the offer and uptake of genetic testing.

MATERIALS AND METHODS

The GC-HBOC comprises 15 university centres. Using standardised clinical criteria (table 2),

INTRODUCTION

Women with pathogenic germline mutations in the *BRCA1* and *BRCA2* genes are at an increased lifetime risk for breast cancer (BC) and ovarian cancer (OC) compared with the general population.¹ Identification of mutation carriers is an important prerequisite for targeted clinical management. The



CrossMark

To cite: Kast K, Rhiem K, Wappenschmidt B, *et al*. *J Med Genet* 2016;**53**: 465–471.

Table 1 Family characteristics

	Total	BRCA1	BRCA2	Negative
Families, no	21 401	3398	1766	16 265
Members				
Total number	617 578	99 708	52 861	466 013
Median per family	25	26	26	25
Members with cancer, no				
uBC ₅₀₋	26 236	4996	2519	18 770
uBC ₅₁₊	24 452	2789	2033	19 655
bBC ₅₀₋	4132	1304	427	2414
bBC ₅₁₊	1896	239	168	1492
OC	7250	2650	730	3897
BC and OC	1917	805	202	916
Male BC	671	62	193	420
Age of onset (mean)*				
BC	49.0	44.4	47.5	50.2
OC	51.5	49.8	55.6	51.9
Male BC	58.0	56.9	60.4	57.0

*Mean ages of onset for BC and OC were significantly different in all pairwise group comparisons ($p < 0.001$). For male BC, mean age of onset was significantly different between BRCA2-positive and BRCA1/2-negative families ($p = 0.006$). bBC, bilateral breast cancer; BC, breast cancer; OC, ovarian cancer.

families with clustering or early onset of BC or OC are registered and tested for the presence of deleterious germline mutations in *BRCA1* and *BRCA2*. Comprehensive data on familial cancer history, including a detailed pedigree, pathology reports and results of molecular testing, are documented in a central database using standardised electronic case report forms.

A total number of 21 401 families, who were registered from 1996 until 2014, were included in the present analysis. All families fulfilled the clinical inclusion criteria shown in table 2. Families who were ascertained through a known pathogenic mutation rather than by clinical criteria were not included. If possible, the family member with the most severe phenotype (defined as bBC, BC and OC, or earliest age of onset) was

chosen as the index patient and was searched for *BRCA1* and *BRCA2* mutations. In case that no DNA from an affected family member could be obtained, mutation analysis was performed in unaffected individuals. BC included ductal carcinoma in situ. OC included cancers of the fallopian tube and primary peritoneal cancers irrespective of histopathological subtype and grading. BC cases with an age of onset of 50 years or earlier are hereafter referred to as premenopausal BC (BC₅₀₋). BC at the age from 51 years onwards are referred to as postmenopausal BC (BC₅₁₊). For bBC, the age of onset of the first cancer was considered. The present study did not include families with single cases of OC or single cases of male breast cancer (mBC) since these families were not part of the clinical inclusion criteria.

Mutation analysis

Mutation analysis was performed using direct sequencing or a pre-screening step followed by direct sequencing of suspect fragments. Pre-screening methods comprised mainly denaturing high-performance liquid chromatography and high-resolution melting.^{15 16} Before the year 1999, single-strand conformation polymorphism and protein truncation test were used. If no deleterious sequence alterations were found in these steps, an additional screening for large genomic alterations was performed using multiplex ligation-dependent probe amplification. In the present study, 88.4% of all *BRCA1/2*-negative index patients were searched for large genomic alterations. Mutations were classified according to the International Agency for Research on Cancer (IARC) system and considered pathogenic or likely pathogenic (class 4 or 5) based on literature evidence, multifactorial likelihood and functional analyses of the ENIGMA consortium that comprises genetic data of the GC-HBOC database.¹⁷⁻¹⁹

Statistical analysis

IBM SPSS 22 was used for statistical analysis. The 95% CIs for mutation frequencies were calculated using Wilson's score

Table 2 Familial breast and ovarian cancer histories used as inclusion criteria for *BRCA1/2* mutation testing in German Consortium for Hereditary Breast and Ovarian Cancer and observed mutation prevalences

Distinct groups of familial BC and OC history (including proband)	Families with pathogenic mutation										
	Families		BRCA1/2			BRCA1			BRCA2		
	N	% of total	n	Prev (%)	95% CI (%)	n	Prev (%)	95% CI (%)	n	Prev (%)	95% CI (%)
Total	21 401	100.0	5136	24.0	23.4 to 24.6	3398	15.9	15.4 to 16.4	1766	8.3	7.9 to 8.6
≥3 females with BC ₅₁₊ (no BC<51, no OC, no mBC)	684	3.2	25	3.7	2.5 to 5.3	9	1.3	0.7 to 2.5	17	2.5	1.6 to 3.9
≥2 females with BC, of these ≥1 with BC ₅₀₋ (no OC, no mBC)	12 996	60.7	2379	18.3	17.7 to 19.0	1439	11.1	10.5 to 11.6	949	7.3	6.9 to 7.8
Single female with unilateral BC ₃₅₋ (no further female BC, no OC, no mBC)	1267	5.9	173	13.7	11.9 to 15.7	116	9.2	7.7 to 10.9	57	4.5	3.5 to 5.8
Single female with bBC ₅₀₋ (no further female BC, no OC, no mBC)	480	2.2	109	22.7	19.2 to 26.7	73	15.2	12.3 to 18.7	36	7.5	5.5 to 10.2
≥1 females with BC and ≥1 female with OC (no mBC)	5072	23.7	2111	41.6	40.3 to 43.0	1624	32.0	30.7 to 33.3	500	9.9	9.1 to 10.7
≥2 females with OC (no female BC, no mBC)	260	1.2	109	41.9	36.1 to 48.0	77	29.6	24.4 to 35.4	34	13.1	9.5 to 17.7
≥1 male with BC and ≥1 females with BC or OC	642	3.0	230	35.8	32.2 to 39.6	60	9.3	7.3 to 11.8	173	26.9	23.7 to 30.5

bBC, bilateral breast cancer; BC, breast cancer; mBC, male breast cancer; OC, ovarian cancer; Prev, prevalence.

method. For comparison of mutation frequencies between groups, the χ^2 test was used. *p* Values <0.05 were considered significant.

RESULTS

Basic characteristics of the study population are summarised in [table 1](#). Of 21 401 families, 3398 (15.9%) had a pathogenic mutation in *BRCA1* and 1766 (8.3%) in *BRCA2*. Of these, 28 families had mutations in both genes, *BRCA1* and *BRCA2*. The median number of family members documented in each pedigree was 25, regardless of *BRCA* mutation status. The mean age at cancer diagnosis was 44.4 years for BC and 49.8 years for OC in *BRCA1*-positive families compared with 47.5 and 55.6 years, in *BRCA2*-positive families, and 50.2 and 51.9 years in *BRCA*-negative families. The mean age at diagnosis of mBC in *BRCA1*-positive families was 56.9 years, which was younger than in *BRCA2* mutation carriers (60.4 years) and equal to *BRCA1/2*-negative families (57.0 years).

The families were classified into seven mutually exclusive groups of aggregated familial cancer histories ([table 2](#)). These groups cover the inclusion criteria of the GC-HBOC established

in 1996. The overall *BRCA1/2* mutation prevalence was 24.0% (95% CI 23.4% to 24.6%). The highest mutation frequencies were seen in families with at least two OC (41.9%, 95% CI 36.1% to 48.0%) and families with at least one BC and one OC (41.6%, 95% CI 40.3% to 43.0%) followed by families with mBC and at least one additional female with BC or OC (35.8%, 95% CI 32.2% to 39.6%). *BRCA1* mutations were more frequent in families with OC, whereas *BRCA2* mutations were more frequent in families with mBC. The largest group comprises families with at least two cases of BC, at least one that was diagnosed before the age of 51 (*n*=12 996, 60.7% of all families). The lowest mutation frequencies (3.7%, 95% CI 2.5% to 5.3%) were observed in families with three or more cases of postmenopausal BC, but no occurrence of premenopausal BC, OC or male BC.

To characterise mutation frequencies in more detail, the familial cancer histories were further refined. [Table 3](#) shows the mutation prevalences in families, in which exclusively female BC was present (72.1% of all families). Group 1a comprises families with exclusive occurrence of unilateral cases of BC (53.1% of all families), whereas groups 1b (14.6%) and 1c

Table 3 *BRCA* mutation prevalence in families with female breast cancer only

Families with pathogenic mutation													
Familial cancer history (including proband)		Families			BRCA1/2			BRCA1			BRCA2		
Group	BC ₅₀₋	BC ₅₁₊	N	% of total	n	Prev (%)	95% CI (%)	n	Prev (%)	95% CI (%)	n	Prev (%)	95% CI (%)
Group 1a:	0	≥3	522	2.4	20	3.8	2.5 to 5.8	6	1.1	0.5 to 2.5	15	2.9	1.7 to 4.7
female unilateral BC	1*	0	1267	5.9	173	13.7	11.9 to 15.7	116	9.2	7.7 to 10.9	57	4.5	3.5 to 5.8
(no bBC, OC, mBC)	1	1	2577	12.0	227	8.8	7.8 to 10.0	118	4.6	3.8 to 5.5	109	4.2	3.5 to 5.1
	1	2	1239	5.8	102	8.2	6.8 to 9.9	54	4.4	3.4 to 5.6	48	3.9	2.9 to 5.1
	1	≥3	579	2.7	43	7.4	5.6 to 9.9	14	2.4	1.4 to 4.0	29	5.0	3.5 to 7.1
	2	0	1725	8.1	302	17.5	15.8 to 19.4	187	10.8	9.5 to 12.4	116	6.7	5.6 to 8.0
	2	1	1256	5.9	204	16.2	14.3 to 18.4	99	7.9	6.5 to 9.5	105	8.4	7.0 to 10.0
	2	2	477	2.2	76	15.9	12.9 to 19.5	33	6.9	5.0 to 9.6	43	9.0	6.8 to 11.9
	2	≥3	239	1.1	41	17.2	12.9 to 22.4	15	6.3	3.8 to 10.1	26	10.9	7.5 to 15.5
	≥3	0	739	3.5	225	30.4	27.2 to 33.9	143	19.4	16.7 to 22.4	82	11.1	9.0 to 13.6
	≥3	1	462	2.2	127	27.5	23.6 to 31.7	83	18.0	14.7 to 21.7	45	9.7	7.4 to 12.8
	≥3	2	177	0.8	50	28.2	22.1 to 35.3	33	18.6	13.6 to 25.0	17	9.6	6.1 to 14.8
	≥3	≥3	103	0.5	25	24.3	17.0 to 33.4	13	12.6	7.5 to 20.4	12	11.7	6.8 to 19.3
	Total		11 362	53.1	1615	14.2	13.6 to 14.9	914	8.0	7.6 to 8.6	704	6.2	5.8 to 6.7
Group 1b:	1	0	480	2.2	109	22.7	19.2 to 26.7	73	15.2	12.3 to 18.7	36	7.5	5.5 to 10.2
female BC,	1	1	482	2.3	101	21.0	17.6 to 24.8	69	14.3	11.5 to 17.7	32	6.6	4.7 to 9.2
of these ≥1bBC ₅₀₋	1	2	204	1.0	41	20.1	15.2 to 26.1	28	13.7	9.7 to 19.1	13	6.4	3.8 to 10.6
(no OC, mBC)	1	≥3	99	0.5	14	14.1	8.6 to 22.3	10	10.1	5.6 to 17.6	4	4.0	1.6 to 9.9
	2	0	588	2.7	190	32.3	28.7 to 36.2	140	23.8	20.5 to 27.4	52	8.8	6.8 to 11.4
	2	1	348	1.6	109	31.3	26.7 to 36.4	76	21.8	17.8 to 26.5	34	9.8	7.1 to 13.3
	2	2	131	0.6	40	30.5	23.3 to 38.9	26	19.8	13.9 to 27.5	14	10.7	6.5 to 17.1
	2	≥3	55	0.3	14	25.5	15.8 to 38.3	9	16.4	8.9 to 28.3	5	9.1	3.9 to 19.6
	≥3	0	361	1.7	183	50.7	45.6 to 55.8	135	37.4	32.6 to 42.5	49	13.6	10.4 to 17.5
	≥3	1	222	1.0	96	43.2	36.9 to 49.8	72	32.4	26.6 to 38.8	27	12.2	8.5 to 17.1
	≥3	2	95	0.4	44	46.3	36.6 to 56.3	26	27.4	19.4 to 37.1	18	18.9	12.3 to 28.0
	≥3	≥3	64	0.3	22	34.4	23.9 to 46.6	13	20.3	12.3 to 31.7	9	14.1	7.6 to 24.6
	Total		3129	14.6	963	30.8	29.2 to 32.4	677	21.6	20.2 to 23.1	293	9.4	8.4 to 10.4
Group 1c:	0	≥3	162	0.8	5	3.1	1.3 to 7.0	3	1.9	0.6 to 5.3	2	1.2	0.3 to 4.4
female BC,	1	1	197	0.9	19	9.6	6.3 to 14.6	3	1.5	0.5 to 4.4	16	8.1	5.1 to 12.8
of these ≥1bBC ₅₁₊	1	2	175	0.8	14	8.0	4.8 to 13.0	4	2.3	0.9 to 5.7	10	5.7	3.1 to 10.2
(no bBC ₅₀₋ , OC, mBC)	1	≥3	132	0.6	14	10.6	6.4 to 17.0	4	3.0	1.2 to 7.5	10	7.6	4.2 to 13.4
	2	1	75	0.4	13	17.3	10.4 to 27.4	7	9.3	4.6 to 18.0	6	8.0	3.7 to 16.4
	2	2	69	0.3	9	13.0	7.0 to 23.0	7	10.1	5.0 to 19.5	2	2.9	0.8 to 10.0
	2	≥3	62	0.3	11	17.7	10.2 to 29.0	3	4.8	1.7 to 13.3	8	12.9	6.7 to 23.4
	≥3	1	24	0.1	10	41.7	24.5 to 61.2	7	29.2	14.9 to 49.2	3	12.5	4.3 to 31.0
	≥3	2	12	0.1	6	50.0	25.4 to 74.6	2	16.7	4.7 to 44.8	4	33.3	13.8 to 60.9
	≥3	≥3	28	0.1	7	25.0	12.7 to 43.4	6	21.4	10.2 to 39.5	1	3.6	0.6 to 17.7
	Total		936	4.4	108	11.5	9.6 to 13.7	46	4.9	3.7 to 6.5	62	6.6	5.2 to 8.4

*≤35 years.

bBC, bilateral breast cancer; BC, breast cancer; mBC, male breast cancer; OC, ovarian cancer; Prev, prevalence.

(4.4%) include also cases of premenopausal and postmenopausal bilateral BC, respectively. These groups were further stratified by the number of women with premenopausal and postmenopausal BC. The highest mutation frequency was seen in families with at least three females with premenopausal BC, at least one of which was bilateral (50.7%, 95% CI 45.6% to 55.8%). Mutations were detected significantly more frequent in families with a single case of premenopausal bilateral BC than in families with two different women with premenopausal BC (22.7%, 95% CI 19.2% to 26.7% vs 17.5%, 95% CI 15.8% to 19.4%, $p=0.012$). In families with a single case of very early BC before the age of 36, mutations were found in 13.7% (95% CI 11.9% to 15.7%). In all subgroups, the occurrence of additional cases of postmenopausal BC did not considerably change mutation frequencies. In contrast, additional cases of premenopausal BC increased the mutation frequencies considerably.

Table 4 lists families with OC only (group 2a, 1.2% of all families), families with BC and OC (group 2b, 23.7%) and families with occurrence of mBC (group 3, 3.0%). In families with one case of BC and one case of OC, mutation frequencies were considerably higher when the BC case was premenopausal (31.5% 95% CI 28.0% to 35.2% vs 19.2%, 95% CI 15.3% to 23.8%). Double primary premenopausal BC and OC in one individual was associated with a much higher mutation prevalence than the occurrence of premenopausal BC and OC in two different women (49.0%, 95% CI 41.0% to 57.0% vs 31.5%, 95% CI 28.0% to 35.2, $p<0.001$). However, there was no significant difference in mutation prevalence between double primary BC and OC (BCOC) and occurrence of BC and OC in two different women (BC/OC) when the BC was postmenopausal instead of premenopausal (20.3%, 95% CI 14.1% to 28.5% vs 19.2%, 95% CI 15.3% to 23.8, $p=0.788$).

In families with mBC (table 4, group 3) and an additional case of female BC, no significant difference in mutation frequencies could be detected depending on menopausal status of the BC (16.5%, 95% CI 10.4% to 25.1% vs 23.2%, 95% CI 16.0% to 32.5%, $p=0.284$). In families with mBC and additional cases of OC, mutation frequencies for *BRCA1* and *BRCA2* mutations were similar in contrast to families without additional OC cases.

DISCUSSION

Based on a large sample of 21 401 families, the present study provides a detailed characterisation of *BRCA1/2* mutation prevalences for defined patterns of familial BC and OC. The underlying data were collected over a period of almost 20 years (1996–2015) in a standardised way within a German multicentre consortium of interdisciplinary university centres specialised in providing healthcare for families with HBOC. All families suspected of having HBOC were selected for genetic testing according to a set of defined clinical criteria that were compulsory for all participating centres.

The German HBOC consortium has defined inclusion criteria for *BRCA1/2* testing based on at least three generation pedigree analysis.⁶ Currently, GC-HBOC offers genetic testing to index patients if the expected *BRCA* mutation probability is $\geq 10\%$ based on the individual family cancer history. In our study, the overall mutation prevalence was 24.0%. The decision threshold of 10% is exceeded in almost all subgroups of HBOC families including mBC. However, families with exclusive occurrence of three or more postmenopausal BC cases (3.2% of all families) were below the 10% threshold with a mutation of prevalence of only 3.7%. The latter finding is in line with the study of Frank *et al.*,¹¹ who reported a prevalence of 3.9%. Interestingly, the

low prevalence observed in our study was independent of whether bilateral cases were present among the postmenopausal BCs in the family (3.1%) or not (3.8%). Moreover, mutation prevalences did not increase with the number of postmenopausal BC cases in the family. On the contrary, in a previous study we showed that an increasing number of females with BC diagnosed at an age of 60 or later was associated with a decreasing *BRCA1/2* mutation frequency.²⁰

In contrast to postmenopausal BC, mutation prevalence increased considerably with each additional case of premenopausal BC both in families with exclusive occurrence of BC and in families with BC and OC. This agrees well with the results of the study of Frank *et al.*¹¹ In our study, mutation prevalence was even higher if at least one case of premenopausal BC was bilateral. In contrast, the presence of postmenopausal bilateral BC did not increase the chance to detect a deleterious mutation. This observation is in line with a previous study of Gershoni-Baruch *et al.*,²¹ who suggested that bBC per se is not reflective of genetic predisposition, unless associated with early age of onset. However, other studies came to the conclusion that mutation frequencies are similar if two cancers (bBC or BCOC) occurred in one individual compared with two separate individuals.^{10 22–24} In our study, mutations were detected significantly more often in families with a single case of premenopausal bilateral BC than in families with two independent premenopausal BC (22.7% vs 17.5%, $p=0.012$).

In accordance with other studies, the highest mutation rates were observed in families with BC and OC.^{4 9 11 12} As observed in families with exclusive occurrence of BC, mutation prevalence was considerably higher if BC cases were premenopausal. Moreover, mutation prevalence was significantly higher in individuals with double primaries of ovarian and premenopausal BC than if these two cancers occurred in different women.

Our study comprised 642 families with mBC. Due to the inclusion criteria, these families had at least one additional case of female BC or OC. As described in previous studies, *BRCA2* mutations were found more frequently than *BRCA1* mutations in families with mBC but without occurrence of OC. In cases where additional OC cases were present in these families, a similar prevalence of mutations was found in *BRCA1* and *BRCA2*. The additional presence of a premenopausal versus postmenopausal BC increased the mutation prevalence from 16.5% to 23.2%, although this difference was not significant due to low sample sizes in these groups.

The present study revealed a mutation prevalence of 13.7% (9.2%, 95% CI 7.7% to 10.9% for *BRCA1* and 4.5%, 95% CI 3.5% to 5.8% for *BRCA2*) for individuals with early unilateral BC before the age of 36 years without further cancer cases in their family. This is in line with results from two other studies.^{25 26} However, there is evidence in the literature that the prevalence of *BRCA1* and *BRCA2* mutations is high in women with triple-negative breast cancer (TNBC) and that *BRCA1/2* mutations are not restricted to young women or patients with a positive family history.²⁷ Thus, single cases of TNBC might be considered in *BRCA1/2* genetic testing guidelines with an extended age of onset that requires further exploration to confirm a potential cut-off at the age of 60 years as suggested by the data of Couch *et al.*²⁷

Comparison of our data to empiric mutation frequencies of other groups is difficult due to varying inclusion criteria. For example, in high-risk HBOC families of Czech ancestry an overall detection rate of 29% was reported compared with 24.0% by us.¹² That work was restricted to first-degree and second-degree relatives in the maternal line and to third-degree

Table 4 BRCA mutation prevalence in families with ovarian cancer (OC) only (group 2a), both breast cancer (BC) and OC (group 2b), and male breast cancer (mBC) (group 3)

		Families with pathogenic mutation										
		Families		BRCA1/2			BRCA1			BRCA2		
Group	Familial cancer history (including proband)	N	% of total	n	Prev (%)	95% CI (%)	n	Prev (%)	95% CI (%)	n	Prev (%)	95% CI (%)
Group 2a: OC only	2 OC (no BC, mBC)	194	0.9	79	40.7	34.1 to 47.8	50	25.8	20.1 to 32.4	29	14.9	10.6 to 20.6
	≥3 OC (no BC, mBC)	66	0.3	30	45.5	34.0 to 57.4	27	40.9	29.9 to 53.0	5	7.6	3.3 to 16.5
	Total	260	1.2	109	41.9	36.1 to 48.0	77	29.6	24.4 to 35.4	34	13.1	9.5 to 17.7
Group 2b: OC and female BC, but no mBC	1 BC ₅₁₊ OC (no mBC)	118	0.6	24	20.3	14.1 to 28.5	17	14.4	9.2 to 21.9	7	5.9	2.9 to 11.7
	1 BC ₅₀ –OC (no mBC)	145	0.7	71	49.0	41.0 to 57.0	56	38.6	31.1 to 46.7	15	10.3	6.4 to 16.4
	≥2 BCOC (no mBC)	11	0.1	7	63.6	35.4 to 84.8	7	63.6	35.4 to 84.8	0	0.0	0.0 to 25.9
	≥2 BCOC+≥1 BC/OC (no mBC)	1482	6.9	796	53.7	51.2 to 56.2	641	43.3	40.8 to 45.8	160	10.8	9.3 to 12.5
	1 BC ₅₁₊ +1 OC (no mBC, BCOC)	333	1.6	64	19.2	15.3 to 23.8	39	11.7	8.7 to 15.6	26	7.8	5.4 to 11.2
	1 BC ₅₀ –+1 OC (no mBC, BCOC)	645	3.0	203	31.5	28.0 to 35.2	155	24.0	20.9 to 27.5	48	7.4	5.7 to 9.7
	1 BC+≥2 OC (no mBC, BCOC)	248	1.2	113	45.6	39.5 to 51.8	100	40.3	34.4 to 46.5	13	5.2	3.1 to 8.8
	≥2 BC+≥1 OC (no mBC, BCOC)	2090	9.8	833	39.9	37.8 to 42.0	609	29.1	27.2 to 31.1	231	11.1	9.8 to 12.5
	Total	5072	23.7	2111	41.6	40.3 to 43.0	1624	32.0	30.7 to 33.3	500	9.9	9.1 to 10.7
Group 3: mBC	1 mBC+1 BC ₅₁₊ (no OC)	97	0.5	16	16.5	10.4 to 25.1	3	3.1	1.1 to 8.7	13	13.4	8.0 to 21.6
	1 mBC+1 BC ₅₀ – (no OC)	99	0.5	23	23.2	16.0 to 32.5	5	5.1	2.2 to 11.3	18	18.2	11.8 to 26.9
	1 mBC+≥2 BC (no OC)	331	1.5	128	38.7	33.6 to 44.0	29	8.8	6.2 to 12.3	100	30.2	25.5 to 35.4
	≥2 mBC+≥1 BC (no OC)	23	0.1	15	65.2	44.9 to 81.2	1	4.3	0.8 to 21.0	15	65.2	44.9 to 81.2
	1 mBC+≥1 OC (no BC)	15	0.1	8	53.3	30.1 to 75.2	4	26.7	10.9 to 52.0	4	26.7	10.9 to 52.0
	≥1 mBC+≥1 BC +≥1 OC	77	0.4	40	51.9	41.0 to 62.7	18	23.4	15.3 to 34.0	23	29.9	20.8 to 40.8
	Total	642	3.0	230	35.8	32.2 to 39.6	60	9.3	7.3 to 11.8	173	26.9	23.7 to 30.5

Prev, prevalence.

relatives in the case of paternal transmission. In the work of Frank *et al.*¹¹ the overall detection frequency was 15.7% in non-Ashkenazi individuals, but postmenopausal BC was not taken into account.

Some limitations have to be mentioned. First, in our study only aggregated familial cancer histories were correlated with mutation prevalence, that is, the pedigree size and family structure was not considered. Mutation prediction algorithms such as BOADICEA, BRCAPRO or IBIS, which use the pedigree structure and an underlying genetic model of inheritance, might have a better predictive performance.^{28–31} Second, it is known that BRCA1/2-associated OCs are characterised by a high grade (G2/3) serous histology.^{32–34} In our study, we did not restrict the analysis to OC with specific pathological features since pathology information was not available for all OC. Thus, we cannot exclude that a restriction to this specific type of OC would have led to larger mutation frequencies.

In summary, we provide a detailed overview on empiric BRCA1/2 mutation frequencies for at-risk individuals with different familial cancer histories. The number of postmenopausal BC failed to show a systematic correlation with mutation frequency. Bilateral BC or premenopausal BC and OC in one affected family member conferred a higher mutation prevalence

than two primaries in separate individuals. Our analysis provides a simple means to get an overview about expected mutation probabilities during clinical and genetic counselling before mutation testing is considered for both counsellors and counselees. Moreover, our results provide guidance for healthcare professionals and decision-makers to define consistent clinical criteria for decision-making to undergo genetic testing for individuals with suspected HBOC.

Author affiliations

¹Department of Gynecology and Obstetrics, University Hospital Carl Gustav Carus, TU Dresden, Dresden, Germany

²German Cancer Consortium (DKTK) Dresden and German Cancer Research Center (DKFZ), Heidelberg, Germany

³Center for Hereditary Breast and Ovarian Cancer, Center for Integrated Oncology (CIO) and Center for Molecular Medicine Cologne (CMMC), Medical Faculty, University of Cologne and University Hospital Cologne, Cologne, Germany

⁴Department for Gynecology and Obstetrics, LMU Munich, Munich, Germany

⁵Department for Gynecology and Obstetrics, Technical University of Munich, Munich, Germany

⁶Breast Center, Department of Gynecology, Red Cross Hospital, Munich, Germany

⁷Institute of Human Genetics, Heidelberg University, Heidelberg, Germany

⁸Department of Gynecology and Obstetrics, Heidelberg University Hospital, Heidelberg, Germany

⁹German Cancer Consortium (DKTK), NCT Heidelberg and German Cancer Research Center (DKFZ), Heidelberg, Germany

¹⁰Medical Faculty, Institute of Human Genetics and Anthropology, Heinrich-Heine University, Düsseldorf, Germany

¹¹Department of Gynecology and Obstetrics, Medical Faculty, Heinrich-Heine University, Düsseldorf, Germany

¹²Department of Human Genetics, University of Würzburg, Würzburg, Germany

¹³Department of Gynaecology and Obstetrics, University Hospital of Schleswig-Holstein, Campus Kiel, Christian-Albrechts University Kiel, Kiel, Germany

¹⁴Institute of Human Genetics, University of Ulm, Ulm, Germany

¹⁵Department of Gynecology and Obstetrics, University Hospital, Universität Ulm, Ulm, Germany

¹⁶Institute of Human Genetics, Hannover Medical School, Hannover, Germany

¹⁷Institute of Medical and Human Genetics, Charité-University Medical Center, Berlin, Germany

¹⁸Department of Gynecology, Charité University Medicine Berlin, Berlin, Germany

¹⁹Institute for Human Genetics, University of Münster, Münster, Germany

²⁰Department of Gynecology and Obstetrics, University of Leipzig, Leipzig, Germany

²¹Institute of Medical Genetics and Applied Genomics, University of Tübingen, Tübingen, Germany

²²Institute of Human Genetics, University of Regensburg, Regensburg, Germany

²³Department of Obstetrics and Gynecology, University of Göttingen, Göttingen, Germany

²⁴Institute for Medical Informatics, Statistics and Epidemiology, University of Leipzig, Leipzig, Germany

Acknowledgements The authors thank all the participating women and the German Cancer Aid for supporting GC-HBOC.

Contributors KK: conception of the work, acquisition and interpretation of data, writing of the manuscript, final approval of the version to be published. AM and RKS: conception of the work, acquisition and interpretation of data, critical revision of the manuscript for important intellectual content, final approval of the version to be published. CE: conception of the work, analysis and interpretation of data, writing of the manuscript, final approval of the version to be published. All other authors: acquisition of data, critical revision of the manuscript for important intellectual content, final approval of publication. All authors agree to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Funding Deutsche Krebshilfe (German Cancer Aid), grant no. 110837.

Competing interests None declared.

Patient consent Obtained.

Ethics approval The study has been approved by the Institutional Review Boards of each participating centre.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- Chen S, Parmigiani G. Meta-analysis of *BRCA1* and *BRCA2* penetrance. *J Clin Oncol* 2007;25:1329–33.
- Claus EB, Risch N, Thompson WD. Autosomal dominant inheritance of early-onset breast cancer. Implications for risk prediction. *Cancer* 1994;73:643–51.
- Fishman A, Dekel E, Chetrit A, Lerner-Geva L, Bar-Am A, Beck D, Beller U, Ben-Baruch G, Piura B, Friedman E, Struewing JP, Modan B. Patients with double primary tumors in the breast and ovary—clinical characteristics and *BRCA1-2* mutations status. *Gynecol Oncol* 2000;79:74–8.
- Ford D, Easton DF, Stratton M, Narod S, Goldgar D, Devilee P, Bishop DT, Weber B, Lenoir G, Chang-Claude J, Sobol H, Teare MD, Struewing J, Arason A, Scherneck S, Peto J, Rebbeck TR, Tonin P, Neuhausen S, Barkardottir R, Eyfjord J, Lynch H, Ponder BA, Gayther SA, Zelada-Hedman M, and the Breast Cancer Linkage Consortium. Genetic heterogeneity and penetrance analysis of the *BRCA1* and *BRCA2* genes in breast cancer families. The Breast Cancer Linkage Consortium. *Am J Hum Genet* 1998;62:676–89.
- Hall JM, Lee MK, Newman B, Morrow JE, Anderson LA, Huey B, King MC. Linkage of early-onset familial breast cancer to chromosome 17q21. *Science (New York, NY)* 1990;250:1684–9.
- Meindl A, Ditsch N, Kast K, Rhiem K, Schmutzler RK. Hereditary breast and ovarian cancer: new genes, new treatments, new concepts. *Dtsch Arztebl Int* 2011;108:323–30.
- Kreienberg R, Albert U, Follmann M, Kopp I, Kühn T, Wöckel A. Interdisziplinäre S3-Leitlinie für die Diagnostik, Therapie und Nachsorge des Mammakarzinoms. *Senologie-Zeitschrift für Mammadiagnostik und-therapie* 2013;10:164–92.
- Familial Breast Cancer. *Classification and care of people at risk of familial breast cancer and management of breast cancer and related risks in people with a family history of breast cancer*. Cardiff, UK: National Collaborating Centre for Cancer, 2013.
- Capalbo C, Ricevuto E, Vestri A, Ristori E, Sidoni T, Buffone O, Adamo B, Cortesi E, Marchetti P, Scambia G, Tomao S, Rinaldi C, Zani M, Ferraro S, Frati L, Screpanti I, Gulino A, Giannini G. *BRCA1* and *BRCA2* genetic testing in Italian breast and/or ovarian cancer families: mutation spectrum and prevalence and analysis of mutation prediction models. *Ann Oncol* 2006;17(Suppl 7):vi34–40.
- Claes K, Poppe B, Coene I, Papee AD, Messiaen L. *BRCA1* and *BRCA2* germline mutation spectrum and frequencies in Belgian breast/ovarian cancer families. *Br J Cancer* 2004;90:1244–51.
- Frank TS, Deffenbaugh AM, Reid JE, Hulick M, Ward BE, Lingenfelter B, Gumpfer KL, Scholl T, Tavtigian SV, Pruss DR, Critchfield GC. Clinical characteristics of individuals with germline mutations in *BRCA1* and *BRCA2*: analysis of 10,000 individuals. *J Clin Oncol* 2002;20:1480–90.
- Machackova E, Foretova L, Lukesova M, Vasicikova P, Navratilova M, Coene I, Pavlu H, Kosinova V, Kuklova J, Claes K. Spectrum and characterisation of *BRCA1* and *BRCA2* deleterious mutations in high-risk Czech patients with breast and/or ovarian cancer. *BMC Cancer* 2008;8:140.
- Menkiszak J, Gronwald J, Gorski B, Jakubowska A, Huzarski T, Byrski T, Fosczyńska-Kłoda M, Haus O, Janiszewska H, Perkowska M, Brozek I, Grzybowska E, Zientek H, Gozdz S, Kozak-Klonowska B, Urbanski K, Miturski R, Kowalczyk J, Pluzanska A, Niepsuj S, Koc J, Swiec M, Drosik K, Mackiewicz A, Lamperska K, Strozzyk E, Godlewski D, Stawicka M, Wasiko B, Bebenek M, Rozmiarek A, Rzepka-Gorska I, Narod SA, Lubinski J. Hereditary ovarian cancer in Poland. *Int J Cancer* 2003;106:942–5.
- Risch HA, McLaughlin JR, Cole DE, Rosen B, Bradley L, Fan I, Tang J, Li S, Zhang S, Shaw PA, Narod SA. Population *BRCA1* and *BRCA2* mutation frequencies and cancer penetrances: a kin-cohort study in Ontario, Canada. *J Natl Cancer Inst* 2006;98:1694–706.
- Arnold N, Gross E, Schwarz-Boeger U, Pfisterer J, Jonat W, Kiechle M. A highly sensitive, fast, and economical technique for mutation analysis in hereditary breast and ovarian cancers. *Hum Mutat* 1999;14:333–9.
- Gross E, Arnold N, Pfeifer K, Bandick K, Kiechle M. Identification of specific *BRCA1* and *BRCA2* variants by DHPLC. *Hum Mutat* 2000;16:345–53.
- Eccles DM, Mitchell G, Monteiro AN, Schmutzler R, Couch FJ, Spurdle AB, Gómez-García EB; ENIGMA Clinical Working Group. *BRCA1* and *BRCA2* genetic testing—pitfalls and recommendations for managing variants of uncertain clinical significance. *Ann Oncol* 2015;26:2057–65.
- Plon SE, Eccles DM, Easton D, Foulkes WD, Genuardi M, Greenblatt MS, Hogervorst FB, Hoogerbrugge N, Spurdle AB, Tavtigian SV; IARC Unclassified Genetic Variants Working Group. Sequence variant classification and reporting: recommendations for improving the interpretation of cancer susceptibility genetic test results. *Hum Mutat* 2008;29:1282–91.
- Spurdle AB, Healey S, Devereau A, Hogervorst FB, Monteiro AN, Nathanson KL, Radice P, Stoppa-Lyonnet D, Tavtigian S, Wappenschmidt B, Couch FJ, Goldgar DE. ENIGMA—evidence-based network for the interpretation of germline mutant alleles: an international initiative to evaluate risk and clinical significance associated with sequence variation in *BRCA1* and *BRCA2* genes. *Hum Mutat* 2012;33:2–7.
- Kast K, Schmutzler RK, Rhiem K, Kiechle M, Fischer C, Niederacher D, Arnold N, Grimm T, Speiser D, Schlegelberger B, Varga D, Horvath J, Beer M, Briest S, Meindl A, Engel C. Validation of the Manchester scoring system for predicting *BRCA1/2* mutations in 9,390 families suspected of having hereditary breast and ovarian cancer. *Int J Cancer* 2014;135:2352–61.
- Gershoni-Baruch R, Dagan E, Fried G, Kepten I, Robinson E. *BRCA1* and *BRCA2* founder mutations in patients with bilateral breast cancer. *Eur J Hum Genet* 1999;7:833–6.
- Couch FJ, DeShano ML, Blackwood MA, Calzone K, Stopfer J, Campeau L, Ganguly A, Rebbeck T, Weber BL. *BRCA1* mutations in women attending clinics that evaluate the risk of breast cancer. *N Engl J Med* 1997;336:1409–15.
- Evans DG, Ahmed M, Bayliss S, Howard E, Lalloo F, Wallace A. *BRCA1*, *BRCA2* and CHEK2 c.1100 delC mutations in patients with double primaries of the breasts and/or ovaries. *J Med Genet* 2010;47:561–6.
- Steinmann D, Bremer M, Rades D, Skawran B, Siebrands C, Karstens JH, Dork T. Mutations of the *BRCA1* and *BRCA2* genes in patients with bilateral breast cancer. *Br J Cancer* 2001;85:850–8.
- FitzGerald MG, MacDonald DJ, Krainer M, Hoover I, O’Neil E, Silva-Arrieto S, Finkelstein DM, Beer-Romero P, Englert C, Sgroi DC, Smith BL, Younger JW, Garber JE, Duda RB, Mayzel KA, Isselbacher KJ, Friend SH, Haber DA. Germ-line *BRCA1* mutations in Jewish and non-Jewish women with early-onset breast cancer. *N Engl J Med* 1996;334:143–9.
- Langston AA, Malone KE, Thompson JD, Daling JR, Ostrander EA. *BRCA1* mutations in a population-based sample of young women with breast cancer. *N Engl J Med* 1996;334:137–42.
- Couch FJ, Hart SN, Sharma P, Toland AE, Wang X, Miron P, Olson JE, Godwin AK, Pankratz VS, Olswold C, Slettedahl S, Hallberg E, Guidugli L, Davila JJ, Beckmann MW, Janni W, Rack B, Ekici AB, Slamon DJ, Konstantopoulou I, Fostira F, Vratimos A, Fountzilas G, Peltari LM, Tapper WJ, Durcan L, Cross SS, Pilarski R, Shapiro CL, Klemp J, Yao S, Garber J, Cox A, Brauch H, Ambrosone C, Nevanlinna H, Yannoukakis D, Slager SL, Vachon CM, Eccles DM, Fasching PA. Inherited

- mutations in 17 breast cancer susceptibility genes among a large triple-negative breast cancer cohort unselected for family history of breast cancer. *J Clin Oncol* 2015;33:304–11.
- 28 Chen S, Wang W, Broman KW, Katki HA, Parmigiani G. BayesMendel: an R environment for Mendelian risk prediction. *Stat Appl Genet Mol Biol* 2004;3:Article21.
 - 29 Lee AJ, Cunningham AP, Kuchenbaecker KB, Mavaddat N, Easton DF, Antoniou AC. BOADICEA breast cancer risk prediction model: updates to cancer incidences, tumour pathology and web interface. *Br J Cancer* 2014;110:535–45.
 - 30 Tyrer J, Duffy SW, Cuzick J. A breast cancer prediction model incorporating familial and personal risk factors. *Stat Med* 2004;23:1111–30.
 - 31 Fischer C, Kuchenbaecker K, Engel C, Zachariae S, Rhiem K, Meindl A, Rahner N, Dikow N, Plendl H, Debatin I, Grimm T, Gadzicki D, Flottmann R, Horvath J, Schrock E, Stock F, Schafer D, Schwaab I, Kartsonaki C, Mavaddat N, Schlegelberger B, Antoniou AC, Schmutzler R, German Consortium for Hereditary Breast and Ovarian Cancer. Evaluating the performance of the breast cancer genetic risk models BOADICEA, IBIS, BRCAPRO and Claus for predicting *BRCA1/2* mutation carrier probabilities: a study based on 7352 families from the German Hereditary Breast and Ovarian Cancer Consortium. *J Med Genet* 2013;50:360–7.
 - 32 Lakhani SR, Manek S, Penault-Llorca F, Flanagan A, Arnout L, Merrett S, McGuffog L, Steele D, Devilee P, Klijn JG, Meijers-Heijboer H, Radice P, Pilotti S, Nevanlinna H, Butzow R, Sobol H, Jacquemier J, Lyonet DS, Neuhausen SL, Weber B, Wagner T, Winqvist R, Bignon YJ, Monti F, Schmitt F, Lenoir G, Seitz S, Hamman U, Pharoah P, Lane G, Ponder B, Bishop DT, Easton DF. Pathology of ovarian cancers in *BRCA1* and *BRCA2* carriers. *Clin Cancer Res* 2004;10:2473–81.
 - 33 Gilks CB, Prat J. Ovarian carcinoma pathology and genetics: recent advances. *Hum Pathol* 2009;40:1213–23.
 - 34 Prat J, Ribe A, Gallardo A. Hereditary ovarian cancer. *Hum Pathol* 2005;36: 861–70.