

Editor-in-Chief

HERBERT D. KAESZ

*Department of Chemistry and Biochemistry
University of California, Los Angeles*

.....
**INORGANIC
SYNTHESES**

Volume 26



A Wiley-Interscience Publication
JOHN WILEY & SONS

New York Chichester Brisbane Toronto Singapore

CONTENTS

Chapter One COMPOUNDS OF THE MAIN GROUP ELEMENTS AND THE LANTHANIDES

1. Potassium Tetradeca hydronaborate(1 $-$)	1
2. Bromotrimethylsilane	4
3. Trimethylphosphine	7
4. Phosphorus Trifluoride	12
5. (η^5 -Cyclopentadienyl)lanthanide Complexes from the Metallic Elements	17
A. Tris(η^5 -cyclopentadienyl)(tetrahydrofuran)neodymium(III).	20
B. Tris(η^5 -cyclopentadienyl)(tetrahydrofuran)samarium(III)	21
C. Bis(η^5 -cyclopentadienyl)(1, 2-dimethoxyethane)-ytterbium(II)	22

Chapter Two MONONUCLEAR TRANSITION METAL COMPLEXES Part I

6. Ammonium [(1 <i>R</i>)-(endo, anti)]-3-bromo-1, 7-dimethyl-2-oxobicyclo[2.2.1]heptane-7-methanesulfonate, and Synthesis and Resolution of <i>cis</i> -Dichlorobis(1, 2-ethanediamine)-chromium(III) Chloride	24
A. Ammonium [(1 <i>R</i>)-(endo, anti)]-3-bromo-1, 7-dimethyl-2-oxobicyclo[2.2.1]heptane-7-methanesulfonate [ammonium (+) ₅₈₉ - α -bromocamphor- π -sulfonate].	25
B. <i>cis</i> -Dichlorobis(1, 2-ethanediamine)chromium(III) Chloride Monohydrate	27
C. (+) ₅₈₉ - Λ - <i>cis</i> -Dichlorobis(1, 2-ethanediamine)-chromium(III) Chloride Monohydrate	28
7. Organometallic Complexes of Isomeric Acyl Isocyanides Chromium Carbonyl (Acyl Isocyanide) and (Acyl Cyanide) Complexes	31

A.	(Benzoyl isocyanide)dicarbonyl(η^6 -methylbenzoate)chromium(0).....	32
B.	(Benzoyl isocyanide)pentacarbonylchromium(0).....	34
C.	(Benzoyl cyanide)pentacarbonylchromium(0).....	35
8.	Diammonium pentachlorooxomolybdate(V)	36
9.	Carbyne Complexes of Tungsten.....	40
A.	Pentacarbonyl[(diethylamino)methylidyne]tungsten Tetrafluoroborate.....	40
B.	<i>trans</i> -Tetracarbonyl[(diethylamino)methylidyne]- (isocyanato)tungsten.....	42
10.	(2, 2-Dimethylpropylidyne)tungsten(VI) Complexes and Precursors for Their Syntheses.....	44
A.	Methoxytrimethylsilane.....	44
B.	Trichlorotrimethoxytungsten	45
C.	(2, 2-Dimethylpropyl)magnesium Chloride.....	46
D.	Tris(2, 2-dimethylpropyl)(2, 2-dimethylpropylidyne)tungsten(VI)	47
E.	Trichloro(1, 2-dimethoxyethane)(2, 2-dimethylpropylidyne)tungsten(VI).....	50
11.	Zerovalent Iron Isocyanide Complexes.....	52
A.	Tetracarbonyl(2-isocyano-1, 3-dimethylbenzene)iron(0)	52
B.	Tricarbonylbis(2-isocyano-1, 3-dimethylbenzene)iron(0)	54
C.	Dicarbonyltris(2-isocyano-1, 3-dimethylbenzene)iron(0)....	56
D.	Carbonyltetrakis(2-isocyano-1, 3-dimethylbenzene)iron(0) ..	57
E.	Pentakis(2-isocyano-1, 3-dimethylbenzene)iron(0)	57
12.	Tetracarbonyliron(0) Complexes Containing Group V Donor Ligands	59
A.	Tetracarbonyl(triphenylphosphine)iron(0).....	61
B.	Other Tetracarbonyl(Group V Donor Ligand) Iron(0) Complexes	61
13.	<i>cis</i> -Tetraamminedihaloruthenium(III) Halide Complexes.....	65
A.	<i>cis</i> -Tetraamminedichlororuthenium(III) Chloride	66
B.	<i>cis</i> -Tetraamminedibromoruthenium(III) Bromide	67
14.	(η^4 -1, 5-Cyclooctadiene)ruthenium(II) Complexes	68
A.	Di- μ -chloro(η^4 -1, 5-cyclooctadiene)ruthenium(II).....	69

B.	Bis(acetonitrile)dichloro(η^4 -1, 5-cyclooctadiene)ruthenium(II).....	69
C.	Bis(benzonitrile)dichloro(η^4 -1, 5-cyclooctadiene)ruthenium(II).....	70
D.	Bis(acetonitrile)dibromo(η^4 -1, 5-cyclooctadiene)ruthenium(II).....	70
E.	Bis(benzonitrile)dibromo(η^4 -1, 5-cyclooctadiene)ruthenium(II).....	71
F.	Tris(acetonitrile)chloro(η^4 -1, 5-cyclooctadiene)ruthenium(II) Hexafluorophosphate(1-) ...	71
G.	Tris(acetonitrile)bromo(η^4 -1, 5-cyclooctadiene)ruthenium(II) Hexafluorophosphate(1-) ...	72
H.	Tetrakis(acetonitrile)(η^4 -1, 5-cyclooctadiene)ruthenium(II) bis[hexafluorophosphate(1-)]	72
I.	(η^4 -1, 5-Cyclooctadiene)tetrakis(hydrazine)ruthenium(II) bis[tetraphenylborate(1-)].....	73
J.	(η^4 -1, 5-Cyclooctadiene)tetrakis(methylhydrazine)ruthenium(II)bis[hexafluorophosphate(1-)]	74
K.	(η^4 -1, 5-Cyclooctadiene)tetrakis(methylhydrazine)ruthenium(II)bis[tetraphenylborate(1-)]	74
15.	Pentacarbonylhydridorhenium	77
16.	Organometallic Fluoro Complexes	81
A.	Tetrakis[tricarbonylfluororhenium(I)]tetrahydrate.....	82
B.	(2, 2'-Bipyridine)tricarbonylfluororhenium(I).....	82
C.	(2, 2'-Bipyridine)tricarbonyl(phosphorodifluoridato)rhenium(I).....	83
D.	Dicarbonylbis[1, 2-ethanediylbis(diphenylphosphine)]fluromolybdenum(II) hexafluorophosphate(1-)	84
17.	(Tetrahydrothiophene)gold(I) or Gold(III) Complexes.....	85
A.	Chloro(tetrahydrothiophene)gold(I)	86
B.	(Pentafluorophenyl)(tetrahydrothiophene)gold(I)	86
C.	Tris(pentafluorophenyl)(tetrahydrothiophene)gold(III).....	87
D.	Benzyltriphenylphosphonium Chloro(pentafluorophenyl)aurate(I).....	88
E.	Bis[1, 2-phenylenebis(dimethylarsine)]gold(I) bis(pentafluorophenyl)aurate(I)	89
F.	(Pentafluorophenyl)- μ -thiocyanato(triphenylphosphene)digold(I).....	90

Chapter Three MONONUCLEAR TRANSITION METAL COMPLEXES Part II. COMPLEXES WITH WEAKLY BONDED ANIONS

Preface W. Beck	92
18. Carbonyl(η^5 -cyclopentadienyl)(tetrafluoroborato)molybdenum and -tungsten Complexes	96
A. Tricarbonyl(η^5 -cyclopentadienyl)(tetrafluoroborato)molybdenum and -tungsten	96
B. Dicarbonyl(η^5 -cyclopentadienyl)(tetrafluoroborato)(triphenylphosphine)-molybdenum and -tungsten, $MCp(CO)_2(PPh_3)(FBF_3)$ ($M = Mo, W$)	98
C. Tricarbonyl(η^5 -cyclopentadienyl)(η^2 -ethene)molybdenum-(1+)tetrafluoroborate(1-)	102
D. Carbonyl(η^5 -cyclopentadienyl)bis(diphenylacetylene)molybdenum(1+)tetrafluoroborate(1-)	102
E. Carbonyl(η^5 -cyclopentadienyl)(diphenylacetylene)(triphenylphosphine)molybdenum(1+)tetrafluoroborate-(1-)	104
F. (Acetone)(tricarbonyl)(η^5 -cyclopentadienyl)molybdenum(1+) and -tungsten(1+)tetrafluoroborate(1-)	105
19. Pentacarbonyl(tetrafluoroborato)rhenium and -manganese and Reactions Thereof	106
A. Pentacarbonylmethylrhenium	107
B. Pentacarbonyl(tetrafluoroborato)rhenium(I)	108
C. Pentacarbonyl(η^2 -ethene)rhenium(1+)tetrafluoroborate(1-)	110
D. Di- μ_3 -(carbon dioxide)octadecacarbonyltetrarhenium	111
20. Manganese(I) and Rhenium(I) Pentacarbonyl(trifluoromethanesulfonato) Complexes	113
A. Pentacarbonyl(trifluoromethanesulfonato)manganese(I), $Mn(CO)_5(O_3SCF_3)$	114
B. Pentacarbonyl(trifluoromethanesulfonato)rhenium(I), $Re(CO)_5(O_3SCF_3)$	115
21. Iridium(III) Complexes with the Weakly Bonded Anions $[BF_4]^-$, and $[OSO_2CF_3]^-$	117

A.	Carbonylchlorohydrido(tetrafluoroborato)bis-(triphenylphosphine)iridium(III)	117
B.	Carbonylchloromethyl(tetrafluoroborato)bis-(triphenylphosphine)iridium(III)	118
C.	Chloro(dinitrogen)hydrido(tetrafluoroborato)bis-(triphenylphosphine)iridium(III)	119
D.	Carbonylhydridobis(trifluoromethanesulfonato)bis-(triphenylphosphine)iridium(III)	120
22.	Dihydridobis(solvent)bis(triphenylphosphine)iridium(III) Tetrafluoroborates	122
A.	(η^4 -1, 5-Cyclooctadiene)bis(triphenylphosphine)iridium(I) Tetrafluoroborate(1 -)	122
B.	Bis(acetone)dihydridobis(triphenylphosphine)iridium(III) Tetrafluoroborate(1 -)	123
C.	Diaquadihydridobis(triphenylphosphine)iridium(III) Tetrafluoroborate(1 -)	124
D.	(1, 2-Diodobenzene)dihydridobis(triphenylphosphine)- iridium(III) Tetrafluoroborate(1 -)	125
23.	<i>cis</i> -Chlorobis(triethylphosphine)- (trifluoromethanesulfonato)platinum(II).....	126
24.	Acetonitrile Complexes of Selected Transition Metal Cations ..	128
A.	Tetrakis(acetonitrile)palladium(II) bis(tetrafluoroborate)- (1 -)	128
B.	<i>cis</i> -Tetrakis(acetonitrile)dinitrosylmolybdenum(II) bis(tetrafluoroborate)(1 -).....	132
C.	<i>cis</i> -Tetrakis(acetonitrile)dinitrosytungsten(II) bis(tetrafluoroborate)(1 -).....	133
25.	The Formation of the Hydrido(methanol)bis(triethyl- phosphine)platinum(II) Cation and Its Reactions with Unsaturated Hydrocarbons.....	134
A.	<i>trans</i> -Hydrido(methanol)bis(triethylphosphine)platinum(II) bis(trifluoromethanesulfonate)(1 -).....	135
B.	μ -Hydrido-hydridophenyltetrakis(triethylphosphine)di- platinum(II) tetraphenylborate(1 -)	136
C.	<i>cis</i> (3-Methoxy-3-oxo- κO -propyl- κC^1)bis(triethylphos- phine)platinum(II) tetraphenylborate(1 -)	138
D.	(η^3 -Cyclooctenyl)bis(triethylphosphine)platinum(II) tetra- phenylborate(1 -).....	139

E. <i>trans</i> -Chloro(<i>cis</i> -1, 2-diphenylethenyl)bis(triethylphosphine)platinum(II)	140
Chapter Four MONONUCLEAR TRANSITION METAL COMPLEXES Part III. METALLOCYCLIC COMPLEXES	
Preface E. Lindner	142
26. Main Group Metal (Li or Mg) <i>o</i> -Phenylenedimethyl[<i>o</i> -C ₆ H ₄ (CH ₂) ₂ or <i>o</i> -C ₆ H ₄ {CH(SiMe ₃) ₂ } Transfer Reagents	144
A. The di-Grignard Reagent of <i>o</i> -bis(Chloromethyl)-benzene, 1	146
B. <i>o</i> -Bis[(trimethylsilyl)methyl]benzene	148
C. μ -[(α , α' , 1, 2- <i>h</i> : α , α' , 1, 2- <i>h</i>)- <i>O</i> -bis-[trimethylsilyl)methyl]-benzene]bis(<i>N</i> , <i>N</i> '-tetramethyl-1, 2-ethanediamine)-dilithium.....	148
27. Cyclometallated Organolithium Compounds	150
A. [2-[(Dimethylamino)methyl]phenyl]lithium	152
B. [2-[(Dimethylamino)methyl]-5-methylphenyl]lithium	152
C. [2-[(Dimethylamino)phenyl]-methyl]lithium	153
D. (Diethyl ether)[8-[(dimethylamino)-1-naphthyl]lithium....	154
28. Metallation of Aromatic Ketones with Pentacarbonylmethylmanganese(I)	155
A. Pentacarbonylmethylmanganese(I) and (2-Acetylphenyl- <i>C</i> , <i>O</i>)tetracarbonylmanganese	156
B. Octacarbonyl-1- κ^4 <i>C</i> , 2 κ^4 <i>C</i> - μ -[carbonyl-1 κ <i>C</i> :2 κ [<i>O</i> -(3-diphenylphosphino-1 κ <i>P</i>)- <i>o</i> -phenylene-2 κ <i>C</i> ¹]-dimanganese, 5	158
29. Cyclo-Cotrimorization of Phosphinothioyls with Alkynes. Phosphinothioyl-containing Manganese Analogs of Cyclopentadiene and Bicycloheptadienes	161
A. Dimethylphosphine sulfide	162
B. Octacarbonylbis- μ -(dimethylphosphinothioyl- <i>P</i> : <i>S</i>)-dimanganese	162
C. Tetracarbonyl[2-(dimethylphosphinothioyl)-1, 2-bis-(methoxycarbonyl)ethenyl- <i>C</i> ': <i>S</i>]manganese	163
D. Tricarbonyl[η^4 -3, 6-dihydro-3, 4, 5, 6-tetrakis(methoxycarbonyl)-2, 2-dimethyl-2 <i>H</i> -1, 2-thiaphosphorin-2-iun-3, 6-diyl]manganese.....	165

E. Tetramethyl thiophenetetracarboxylate	166
F. Tricarbonyl[η^4 -2, 5-dihydro-2, 3, 4, 5-tetrakis(methoxycarbonyl)-1, 1-dimethyl-2 <i>H</i> -phosphol-1-ium-2, 5-diyl]manganese	167
30. Tetracarbonyl{[2-(diphenylphosphino)phenyl]-hydroxymethyl- <i>C</i> , <i>P</i> }manganese	169
31. Cyclometallation Reactions.....	171
A. Benzylpentacarbonylmanganese	172
B. Tetracarbonyl[2-(phenylazo)phenyl- <i>C</i> ¹ , <i>N</i> ²]manganese and μ -(azodi-1, 2-phenylene- <i>C</i> ¹ , <i>N</i> ² : <i>C</i> ^{1'} , <i>N</i> ¹)-octacarbonyldimanganese	173
C. Bis[μ -chloro-[2-(phenylazo)phenyl- <i>C</i> ¹ , <i>N</i> ²]palladium].....	175
D. Tricarbonyl[2-(phenylazo)phenyl]- <i>C</i> ¹ , <i>N</i> ²]cobalt	176
E. <i>cis</i> -Bis(benzo[<i>h</i>]quinolin-10-yl- <i>C</i> ¹⁰ , <i>N</i> ¹)dicarbonylruthenium(II).....	177
F. (η^5 -Cyclopentadienyl)[2-[diphenoxyphosphino]-oxy]phenyl- <i>C</i> , <i>P</i>]triphenylphosphite- <i>P</i>)ruthenium(II)	178
32. (η^6 -Hexamethylbenzene)ruthenium Complexes Containing Hydride and <i>ortho</i> -Metallated Triphenylphosphine	180
A. Chloro(η^6 -hexamethylbenzene)(hydrido)(triphenylphosphine)ruthenium(II)	181
B. [6-Diphenylphosphino)phenyl- <i>C</i> ¹ , <i>P</i>](η^6 -hexamethylbenzene)(hydrido)ruthenium(II).....	182
33. An Osmium Containing Benzene Analog, Os(CSCHCHCHCH)(CO)(PPh ₃) ₂ , Carbonyl (5-thioxo-1, 3-pentadiene-1, 5-diyl- <i>C</i> ¹ , <i>C</i> ⁵ , <i>S</i>)bis(triphenylphosphine)osmium and Its Precursors	184
A. Dichlorotris(triphenylphosphine)osmium(II)	184
B. Dichloro(thiocarbonyl)tris(triphenylphosphine)osmium(II) ..	185
C. Dihydrido(thiocarbonyl)tris(triphenylphosphine)-osmium(II)	186
D. Carbonyl(thiocarbonyl)tris(triphenylphosphine)osmium(0)..	187
E. Carbonyl(5-thioxo-1, 3-pentadiene-1, 5-diyl- <i>C</i> ¹ , <i>C</i> ⁵ , <i>S</i>)-bis(triphenylphosphine)osmium, Os(CSCHCHCHCH)(CO)(PPh ₃) ₂	188
34. (1, 3-butadiene-1, 4-diyl)(η^5 -Cyclopentadienyl)-(triphenylphosphine)cobalt with Various Substituents.....	189
A. Chlorotris(triphenylphosphine)cobalt	190

B.	(η^5 -Cyclopentadienyl)bis(triphenylphosphine)cobalt	191
C.	(η^5 -Cyclopentadienyl)[1, 1'-(η^2 -1, 2-ethynediyl)dibenzene]- (triphenylphosphine)cobalt and (η^5 -cyclopentadienyl)-(methyl-3-phenyl- η^2 -2-propynoate)(triphenylphosphine)- cobalt.....	192
D.	(η^5 -Cyclopentadienyl)(2, 3-dimethyl-1, 4-diphenyl-1, 3-butadiene-1, 4-diyl)(triphenylphosphine)cobalt.....	195
E.	[1, 4-Bis(methoxycarbonyl)-2methyl-3-phenyl-1, 3-butadiene-1, 4-diyl](η^5 -cyclopentadienyl)(triphenylphosphine)cobalt and [1,3-bis(methoxycarbonyl)-2-methyl-4-phenyl-1, 3-butadiene-1, 4-diyl](η^5 -cyclopentadienyl)-(triphenylphosphine)- cobalt.....	197
35.	An Iridium(III) Complex Containing Cyclometallated Tri-phenylphosphine Formed by Isomerization of an Iridium(I) Triphenylphosphine Complex	200
A.	Chlorotris(triphenylphosphine)iridium(I).....	201
B.	(OC-6-53)Chloro[(2-diphenylphosphino)phenyl- C^1 , <i>P</i>]- hydridobis(triphenylphosphine)iridium(III)	202
36.	Nickel-Containing Cyclic Amide Complexes	204
A.	[2-Methylpropanamidato(2-)- C^3 , <i>N</i>](tricyclohexylphosphine)nickel(II), (tcyp) $\text{NiCH}_2\text{CH}(\text{CH}_3)\text{CONH}$	205
B.	[Butanamidato(2-) C^4 , <i>N</i>]- (tricyclohexylphosphine)nickel(II), $\text{NiCH}_2\text{CH}_2\text{CH}_2\text{CONH}$ (tcyp).....	206
37.	Six-Membered Cyclopalladated Complexes of 2-Benzylpyridine.	208
A.	Bis(μ -acetato)bis[2-(2-pyridinylmethyl)phenyl- C^1 , <i>N</i>]- dipalladium(II).....	208
B.	Di- μ -chloro-bis[2-(2-pyridinylmethyl)phenyl- C^1 , <i>N</i>]- dipalladium(II).....	209
C.	Chloro(3, 5-dimethylpyridine)[2-(2-pyridinylmethyl)- phenyl- C^1 , <i>N</i>] palladium(II)	210
38.	Cyclopalladated Compounds.....	211
A.	Di- μ -chloro-bis[2-[(dimethylamino)methyl]- phenyl- <i>C</i> , <i>N</i>]dipalladium(II).....	212
B.	Di- μ -chloro-bis(8-quinolylmethyl- <i>C</i> , <i>N</i>)- dipalladium(II).....	213

Chapter Five POLYNUCLEAR TRANSITION METAL COMPLEXES

Preface J. R. Shapley	215
39. Ditungsten Tetracarboxylates	219
A. Tungsten Tetrachloride $[WCl_4]_x$	221
B. Tetrakis(trifluoroacetato)ditungsten(II). $W_2(O_2CCF_3)_4(W^{\frac{4}{2}}W)$	222
C. Tetrakis(2, 2-dimethylpropanoato)ditungsten(II). $W_2(O_2C-t-Bu)_4(W^{\frac{4}{2}}W)$	223
D. Tetrakis(acetato)ditungsten(II). $W_2(O_2CCH_3)_4(W^{\frac{4}{2}}W)$	224
40. Diphenylphosphino-Bridged Dimanganese Complexes	225
A. (μ -Diphenylphosphino)- μ -hydrido-bis(tetracarbonyl-manganese)(Mn—Mn)	226
B. μ -Nitrido-bis(triphenylphosphorus)(1+)(μ -diphenyl-phosphino)bis(tetracarbonylmanganate)(Mn—Mn)(1-) ...	228
C. Heterometallic Trinuclear Clusters	
1. Octacarbonyl- $1\kappa^4C, 2\kappa^4C(\mu$ -diphenylphosphino- $1:2\kappa P$)-(triphenylphosphine- $3\kappa P$)-triangulo- dimanganesegold	229
2. (μ -Chloromercurio)(μ -diphenylphosphino)-bis(tetra- carbonylmanganese)(Mn—Mn)	230
41. Preparation of Some Bimetallic μ (η^1 -C, O) Acetyl Complexes ..	231
A. Dicarbonyl(η^5 -cyclopentadienyl)(tetrahydrofuran)iron Hexafluorophosphate	232
B. (μ -Acetyl-C:O)-Bis(dicarbonyl- η^5 -cyclopentadienyliron)- (1+) Hexafluorophosphate(1-)	235
C. (μ -Acetyl- $2\kappa C^1:1\kappa O$ -tricarbonyl- $1\kappa^2C, 2\kappa C$ -bis[$1:2(\eta^5$ - cyclopentadienyl)] (triphenylphosphine- $2\kappa P$)diiron(1+) Hexafluorophosphate(1-)	237
D. (μ -Acetyl- $2\kappa C^1:1\kappa O$)-pentacarbonyl- $\kappa^3C, 2\kappa^2C$ -bis[$1:2(\eta^5$ - cyclopentadienyl)]molybdenumiron(1+) Hexa- fluorophosphate(1-)	239
E. (μ -Acetyl- $2\kappa C^1:1\kappa O$)-tetracarbonyl- $1\kappa^3C,$ $2\kappa^2C$ -bis[$1:2(\eta^5$ -cyclopentadienyl)](triphenyl- (η^5 -cyclopentadienyl)phosphine- $1\kappa P$)- molybdenumiron(1+) Hexafluorophosphate(1-)	241
42. Trinuclear Metal Complexes	243
A. Nonacarbonyl dihydrido- μ_3 -thio-triiron	244
B. Nonacarbonyl- μ_3 -thio-dicobaltiron	245

43.	Bis[μ -nitrido-bis(triphenylphosphorus)(1+)[μ_4 -carbido-dodecacarbonyltetraferrate(2-)]	246
44.	Dinuclear Ruthenium(II) Carboxylate Complexes	249
A.	Di- μ -chloro(η^4 -bicyclo[2.2.1]hepta-2, 5-diene)ruthenium(II) Polymer	250
B.	(η^4 -Bicyclo[2.2.1]hepta-2, 5-diene)bis(η^3 -2-propenyl)-ruthenium(II).	251
C.	Di- μ -chloro-(η^4 -cycloocta-1, 5-diene)ruthenium(II) Polymer	253
D.	(η^4 -Cycloocta-1, 5-diene)bis(η^3 -2-propenyl)ruthenium(II)....	254
E.	μ -Aqua-bis-(μ -trifluoroacetato)-bis[η^4 -cycloocta-1, 5-diene)(trifluoroacetato)ruthenium(II)]	254
F.	μ -Aqua-bis-(μ -chloroacetato)-bis[(chloroacetato)(η^4 -cycloocta-1, 5-diene)ruthenium(II)].....	256
G.	μ -Aqua-bis-(μ -trichloroacetato)-bis[(η^4 -bicyclo[2.2.1]-hepta-2, 5-diene)(trichloroacetato)ruthenium(II)].....	256
45.	Polynuclear Ruthenium Complexes.....	259
A.	Dodecacarbonyltriruthenium, Ru ₃ (CO) ₁₂	259
B.	Dodecacarbonyltetra- μ -hydrido-tetraruthenium, Ru ₄ (μ -H) ₄ (CO) ₁₂	262
46.	A Phosphino Bridged Ruthenium Cluster: Nonacarbonyl- μ -hydrido-(μ -diphenylphosphino)triruthenium(0)	264
47.	[μ -Nitrido-bis(triphenylphosphorus)(1+)] [deca-carbonyl-1 κ^3 C, 2 κ^3 C, 3 κ^4 C- μ -hydrido-1:2 κ^2 H-bis-(triethylsilyl)-1 κ Si, 1 κ Si)triangulo-triruthenate(1-)]	269
48.	Tri- and Tetranuclear Carbonyl-Ruthenium Cluster Complexes Containing Isocyanide, Tertiary Phosphine, and Phosphite Ligands. Radical Ion-Initiated Substitution of Metal Cluster Carbonyl Complexes Under Mild Conditions.....	271
A.	Sodium Benzophenone Ketyl Solution.....	273
B.	Undecacarbonyl(dimethylphenylphosphine)triruthenium, Ru ₃ (CO) ₁₁ (PMe ₂ Ph)	273
C.	Decacarbonyl(dimethylphenylphosphine)(2-isocyano-2-methylpropane)triruthenium, Ru ₃ (CO) ₁₀ (CN- <i>t</i> -Bu)-(PMe ₂ Ph)	275
D.	Decacarbonyl[methylenebis(diphenylphosphine)]-triruthenium, Ru ₃ (CO) ₁₀ [(Ph ₂ P) ₂ CH ₂]	276

E. [μ -Ethynediylbis(diphenylphosphine)]bis[undeca-carbonyltriruthenium], $\{(Ru_3(CO))_1)_2[\mu-C_2(PPh_2)_2]\}$	277
F. Undecacarbonyltetrahydrido[tris(4-methylphenyl) phosphite]tetraruthenium, $Ru_4H_4(CO)_{11}[P(OC_6H_4Me-p)_3]$	277
G. Decacarbonyl(dimethylphenylphosphine)tetra-hydrido[tris(4-methylphenyl) phosphite]tetraruthenium, $Ru_4H_4(CO)_{10}(PMe_2Ph)[P(OC_6H_4Me-p)_3]$	278
 49. Carbido-carbonyl Ruthenium Cluster Complexes	280
A. μ_6 -Carbido-heptadecacarbonylhexaruthenium.....	281
B. μ_5 -Carbido-pentadecacarbonylpentaruthenium.....	283
C. Disodium or Bis[μ -nitrido-bis-(triphenylphosphorus)(1+)] Salts of [μ_5 -Carbidotetradecacarbonylpentaruthenate(2-)]	284
50. Nitrido-Ruthenium Cluster Complexes.....	286
A. μ -Nitrido-bis(triphenylphosphorus)(1+)azide, $PPN[N_3]$...	286
B. μ -Nitrido-bis(triphenylphosphorus)(1+)hexadeca-carbonylnitridohexaruthenate(1-), $[PPN][Ru_6N(CO)_{16}]$.	287
C. μ -Nitrido-bis(triphenylphosphorus)(1+)tetraadeca-carbonylnitridopentaruthenate(1-), $[PPN][Ru_5N(CO)_{14}]$	288
 51. Some Useful Derivatives of Dodecacarbonyltriosmium	289
A. (Acetonitrile)undecacarbonyltriosmium	290
B. Undecacarbonyl(pyridine)triosmium	291
C. Bis(acetonitrile)decacarbonyltriosmium	292
 52. Dodecacarbonyltetra- μ -hydrido-tetrahedro-tetraosmium	293
 53. High Nuclearity Carbonyl Cluster Complexes of Osmium.....	295
A. Octadecacarbonylhexaosmium	295
B. Bis[μ -nitrido-bis(triphenylphosphorus)(1+)] pentadeca-carbonylpentaosmate(2-).	299
C. Bis[μ -nitrido-bis(triphenylphosphorus)(1+)] octadeca-carbonylhexaosmate(2-).	300
D. Octadecacarbonyldihydridohexaosmium.....	301
 54. Thioosmium Clusters	303
A. (μ -Benzenthiolato)decacarbonyl- μ -hydrido-triosmium, 1 ..	304
B. μ_3 -Carbonyl-nonacarbonyl- μ_3 -thio-triosmium, 2	305
C. Nonacarbonyldi- μ_3 -thio-triosmium, 3	306
D. Tridecacarbonyldi- μ_3 -thio-tetraosmium, 4	307
E. Dodecacarbonyldi- μ_3 -thio-tetraosmium, 5	307

F. Analogous Selenido Clusters.....	308
55. Tris(η^5 -cyclopentadienyl)bis(μ_3 -phenyl-methylidyne)-tricobalt.....	309
56. Bis(tetramethylammonium) Hexa- μ -carbonyl-hexacarbonylhexanickelate(2-), $[\text{N}(\text{CH}_3)_4]_2[\text{Ni}_6(\text{CO})_6(\mu\text{-CO})_6]$	312
57. Bis(tetrabutylammonium) Hexa- μ -carbonyl-hexa-carbonylhexaplatinate(2-), $[\text{N}(\text{C}_4\text{H}_9)_4]_2[\text{Pt}_6(\text{CO})_6(\mu\text{-CO})_6]$	316
58. Preparation of the Carbonyl Platinum Anions, $[\text{Pt}_3(\text{CO})_6]_n^{2-}$ ($n = 3-5$).....	319
A. Bis(tetraethylammonium) Pentakis[tri- μ -carbonyl-tricarbonyltriplatinate](2-), $[\text{N}(\text{C}_2\text{H}_5)_4]_2[\text{Pt}_3(\text{CO})_6]_5$	320
B. Bis(tetraethylammonium) Tetrakis[tri- μ -carbonyl-tri-carbonyltriplatinate](2-), $[\text{N}(\text{C}_2\text{H}_5)_4]_2[\text{Pt}_3(\text{CO})_6]_4$	321
C. Bis(tetraethylammonium) Tris[tri- μ -carbonyl-tricarbonyl-triplatinate](2-), $[\text{N}(\text{C}_2\text{H}_5)_4]_2[\text{Pt}_3(\text{CO})_6]_3$	322
59. Synthesis of Gold-Containing Mixed-Metal Cluster Complexes.	324
A. Chloro(triphenylphosphine)gold, $\text{AuCl}(\text{PPh}_3)$	325
B. μ_3 -Oxo-[tris(triphenylphosphinegold)](1+)tetra-fluoroborate(1-), $[\text{O}\{\text{Au}(\text{PPh}_3)\}_3][\text{BF}_4]$	326
C. Dodecacarbonyltris(triphenylphosphine)cobalttrigoldtriruthenium, $\text{Au}_3\text{CoRu}_3(\text{CO})_{12}(\text{PPh}_3)_3$	327
60. Mercury-Bridged Transition Metal Cluster Derivatives and Their Precursor.....	328
A. Nonacarbonyl- μ_3 -($\kappa:\eta^2:\eta^2$ -3,3-dimethyl-1-butynyl)- μ -hydrido- <i>triangulo</i> -triruthenium	329
B. 1. Nonacarbonyl- μ_3 -($\kappa:\eta^2:\eta^2$ -3,3-dimethyl-1-butynyl)- μ -iodomercurio- <i>triangulo</i> -triruthenium	330
2. μ -(Bromomercurio)-nonacarbonyl- μ_3 -($\kappa:\eta^2:\eta^2$ -3,3-dimethyl-1-butynyl)- <i>triangulo</i> -triruthenium	331
C. μ_4 -Mercurio-bis-[Nonacarbonyl- μ_3 -($\kappa:\eta^2:\eta^2$ -3,3-dimethyl-1-butynyl)- <i>triangulo</i> -triruthenium]	333
D. Nonacarbonyl- μ_3 -($\kappa:\eta^2:\eta^2$ -3,3-dimethyl-1-butynyl)- μ -{[tricarbonyl(η^5 -cyclopentadienyl)molybdio]-mercurio}- <i>triangulo</i> -triruthenium	333
61. Heterobinuclear Nonacarbonyl Complexes and Hydride Complexes of Iron-Chromium, Iron-Molybdenum, and Iron-Tungsten	335

A.	μ -Nitrido-bis(triphenylphosphorus)(1+)tetra-carbonyl hydridoferrate(1-)	336
B.	μ -Nitrido-bis(triphenylphosphorus)(1+)-nonacarbonylhydridoferrate tungstate(1-)	336
C.	Bis[μ -nitrido-bis(triphenylphosphorus)(1+)]nonacarbonylchromateferrate(1-)	339
62.	Cyclopentadienylsodium and some Mono-, Tri-, and Tetrานuclear Metal Carbonyl Derivatives and Cluster Complexes	341
A.	Cyclopentadienylsodium—1,2-dimethoxyethane	342
B.	Sodium tricarbonyl(η^5 -cyclopentadienyl)metallates, DME solvates, $\text{Na}[\text{M}(\text{CO})_3(\eta^5\text{-C}_5\text{H}_5)] \cdot 2\text{DME}$ ($\text{M} = \text{Cr, Mo, or W}$)	343
C.	A Linear Dimetalliplatinum Complex	
1.	Bis(benzonitrile)-dichloroplatinum	345
2.	A Linear Dimetallic Platinum Complex	346
D.	Planar Tetrานuclear Clusters with Mixed-Metal Cores $\text{M}'_2\text{Mo}_2(\text{C}_2\text{H}_5)_2(\text{CO})_6(\text{PPh}_3)_2$, ($\text{M}' = \text{Pd or Pt}$)	
1.	Hexacarbonylbis(η^5 -cyclopentadienyl)bis(triphenylphosphine)dimolybdenumdiplatinum, $[\text{Pt}_2\text{Mo}_2(\eta^5\text{-C}_5\text{H}_5)_2(\text{CO})_6(\text{PPh}_3)_2]$	347
2.	Hexacarbonylbis(η^5 -cyclopentadienyl)bis(triphenylphosphine)dimolybdenumdipalladium, $[\text{Pd}_2\text{Mo}_2(\eta^5\text{-C}_5\text{H}_5)_2(\text{CO})_6(\text{PPh}_3)_2]$	348
63.	Mixed Trinuclear Dicobaltiron and Dicobaltruthenium Cluster Complexes	351
A.	Nonacarbonyl- μ_3 -thio-dicobaltiron, $\text{FeCo}_2(\text{CO})_9\text{S}$	352
B.	Nonacarbonyl- μ_3 -thio-dicobaltruthenium, $\text{RuCo}_2(\text{CO})_9\text{S}$	352
C.	Nonacarbonyl- μ_3 -(phenylphosphinidene)dicobaltiron, $\text{FeCo}_2(\text{CO})_9(\text{PC}_6\text{H}_5)\text{S}$	353
D.	Undecacarbonyldicobaltruthenium, $\text{RuCo}_2(\text{CO})_{11}$	354
64.	Stepwise Syntheses of Ruthenium Mixed-Metal Cluster Complexes	356
A.	Tetraethylammonium Bis(acetonitrile)tetrachlororuthenate(1-), $(\text{Et}_4\text{N})[\text{RuCl}_4(\text{CH}_3\text{CN})_2]$	356
B.	Tetraethylammonium Dodecacarbonyltricobaltruthenate(1-), $(\text{Et}_4\text{N})[\text{Co}_3\text{Ru}(\text{CO})_{12}]$	358
C.	(Acetonitrile) dodecacarbonyltricobaltrutheniumcopper, $\text{Co}_3\text{RuCu}(\text{CH}_3\text{CN})(\text{CO})_{12}]$	359
65.	Mixed-Metal Cluster Complexes of Nickel with Ruthenium or with Osmium	360

A.	Nonacarbonyl(η^5 -cyclopentadienyl)tri- μ -hydrido-nickel triosmium, Os ₃ Ni(μ -H) ₃ (η^5 -C ₅ H ₅)(CO) ₉	362
B.	Nonacarbonyl(η^5 -cyclopentadienyl)tri- μ -hydrido-nickel-triruthenium, Ru ₃ Ni(μ -H) ₃ (η^5 -C ₅ H ₅)(CO) ₉	363
66.	Triosmium and Trinickeltriosmium Cluster Complexes.....	365
A.	Nonacarbonyltris(η^5 -cyclopentadienyl)-trinickeltriosmium .	365
B.	Decacarbonyldihydridotriosmium.....	367
67.	A. Dicobaltplatinum Complex and Its Platinum Precursor.....	369
A.	Dichloro[1, 2-ethanediylbis(diphenylphosphine)platinum ...	370
B.	Heptacarbonyl[1, 2-ethanediylbis(diphenylphosphine)]-dicobaltplatinum.....	370
68.	Salts of Mixed Platinum-Tetrarhodium Cluster Complexes.....	372
A.	Bis[μ -nitrido-bis(triphenylphosphorus)(1 +)][tetra-decacarbonylplatinumtetrarhodate(2 -), [(Ph ₃ P) ₂ N] ₂ [PtRh ₄ (CO) ₁₄]].....	373
B.	Bis[μ -nitrido-bis(triphenylphosphorus)(1 +)][dodeca-carbonylplatinumtetrarhodate(2 -), [(Ph ₃ P) ₂ N] ₂ [PtRh ₄ (CO) ₁₂]].....	375

Chapter Six SOLID STATE

69.	Synthesis and Crystal Growth of A ₃ [M ₂ X ₉](A = Cs, Rb; M = Ti, V, Cr; X = Cl, Br).....	377
A.	Single Crystals of A[Cr ₂ X ₉].....	381
B.	Synthesis of TiBr ₃	382
C.	Polycrystalline A ₃ [Ti ₂ X ₉] and A ₃ [V ₂ X ₉].....	383
D.	Single Crystals of A ₃ [Ti ₂ X ₉] and A ₃ [V ₁ X ₉]	384
70.	Highly Conducting and Superconducting Synthetic Metals.....	386
A.	Synthesis of 5, 5', 6, 6'-tetrahydro-2, 2'-Bi-1, 3-dithiolo[4, 5-b][1, 4]-dithiinylidene, BEDT-TTF 1. 1, 3-Dithiolo[4, 5-b]dithiin-2-thione, EDT-DTT, II	389
	2. 5, 5', 6, 6'-tetrahydro-2, 2'-Bi-1, 3-dithiolo[4, 5-b]-dithiinylidene, BEDT-TTF, III	390
B.	Tetrabutylammonium Perrhenate and Superconducting Bis(5, 5', 6, 6'-tetrahydrdo-2, 2'-bi-1, 3-dithiolo-[4, 5-b][1, 4]dithiinylidene)perrhenate.....	391

C. Tetrabutylammonium fluorosulfate and 2,2'-bi-1,3-dithiolo[4,5- <i>b</i>][1,4]dithiinylidene fluorosulfate	393
<i>Index of Contributors</i>	395
<i>Subject Index</i>	399
<i>Formula Index</i>	411


18. CARBONYL(η^5 -CYCLOPENTADIENYL)-(TETRAFLUOROBORATO)MOLYBDENUM AND TUNGSTEN COMPLEXES

Submitted by WOLFGANG BECK, KLAUS SCHLOTER, KARLHEINZ
SÜNKEL, and GÜNTHER URBAN*

Checked by THOMAS FORSCHNER, ALICIA TODARO, and ALAN CUTLER†

A. TRICARBONYL(η^5 -CYCLOPENTADIENYL)(TETRAFLUOROBORATO)MOLYBDENUM AND -TUNGSTEN^{1a}

An efficient method for the preparation of tetrafluoroborato complexes is hydride abstraction from metal hydrides using triphenylmethylium tetrafluoroborate.^{1a} This method has been first reported by Sanders for hydridoruthenium complexes.²



In a similar way, the hexafluoroarsenate and hexafluoroantimonato complexes $\text{MCp}(\text{CO})_3\text{EF}_5$ ($\text{M} = \text{Mo, W}; \text{E} = \text{As, Sb}$) have been prepared from $\text{MCp}(\text{CO})_3\text{H}$ and $[\text{Ph}_3\text{C}][\text{EF}_6]$.^{1b}

An alternative method for the preparation of $\text{MCp}(\text{CO})_3(\text{FBF}_3)$ ($\text{M} = \text{Mo, W}$) is protonation of $\text{MCP}(\text{CO})_3\text{CH}_3$ by $\text{H}[\text{BF}_4]\cdot\text{Et}_2\text{O}$.^{1c}

Procedure

Tritiylium tetrafluoroborate is commercially available (Fluka AG) and should be freshly recrystallized from dichloromethane or dichloromethane–ethyl acetate prior to use. The hydrido complexes, $\text{MCp}(\text{CO})_3\text{H}$ ³ should be purified by sublimation or by chromatography (neutral alumina, activity 3, pentane eluant) prior to use. All solvents must be rigorously dried and handled under an inert atmosphere, see the preceding general comments.

A quantity of $[\text{Ph}_3\text{C}][\text{BF}_4]$ (0.33 g, 1.0 mmol) is dissolved in 10 mL of CH_2Cl_2 in a 50-mL Schlenk tube, under an inert atmosphere. The solution is cooled to -40°C (using Dry Ice–acetone and a low-temperature thermometer). To this is added $\text{MoCp}(\text{CO})_3\text{H}$ (Ref. 3) (0.22 g, 0.89 mmol) or

*Institut für Anorganische Chemie der Universität München, Meiserstr. 1, 8000 München 2, Federal Republic of Germany.

†Department of Chemistry, Rensselaer Polytechnic Institute, Troy, 12180-3590.

$\text{WCp}(\text{CO})_3\text{H}$ (Ref. 3) (0.30 g, 0.90 mmol). An immediate color change from yellow to purple-red is observed. After stirring for 10 min, 0.2 mL of the solution is syringed into an infrared (IR) solution cell and a spectrum is taken. If a more or less intense band is observed at $\sim 1355 \text{ cm}^{-1}$, indicating the presence of unreacted tritylium salt, small amounts of the corresponding hydrides are then added via a spatula. After stirring for 5 min, the IR spectrum is recorded for another solution aliquot. The addition of hydride is repeated until the IR spectrum of the solution shows no band at 1355 cm^{-1} . As soon as this equivalence point is reached, a sudden color change from dark red to lilac or violet is observed (see the solution in Section A). If this color change does not occur, the presence of moisture can be suspected. In this case the solution may be used for a reaction with stronger ligands than water, otherwise the preparation has to be tried again.

Two procedures are given for the treatment of the solution in Section A. In the first the solution is cooled down to -60°C , and 20 mL of hexane is added. Careful evaporation under vacuum to $\sim 20 \text{ mL}$ removes most of the CH_2Cl_2 . The lilac precipitate is isolated by centrifugation ($\sim 2 \text{ min}$ at 1500 rpm) and decanting off the solution. Hexane (20 mL) is added at -60°C and the suspension is stirred for 10 min. Centrifugation, decanting, and washing are repeated three times. Then the product is dried at -20°C for 8 h on a high-vacuum line (10^{-3} torr).

Alternate Procedure for Treatment of the Solution in Section A

The lilac colored reaction mixture is transferred into a second Schlenk flask (100 mL) using a double-ended stainless steel cannula. The second flask contains hexane previously cooled to -78°C (Dry Ice-acetone bath). A lilac colored solid precipitates. The solvent is siphoned off and the solid is washed three times with hexane (20 mL) previously cooled to -78°C and transferred into the flask using the double ended cannula technique. The wash solvent is siphoned off and remaining solid is dried under vacuum (10^{-3} torr, oil pump) at -40°C for 8 h. Yields: for $\text{MoCp}(\text{CO})_3(\text{FBF}_3)$: 282–319 mg (85–96%), for $\text{WCp}(\text{CO})_3(\text{FBF}_3)$: 357–378 mg (85–90%).

Anal. Calcd. for $\text{C}_8\text{H}_5\text{BF}_4\text{MoO}_3$: C, 28.95; H, 1.52. Found: C, 28.27; H, 1.64. Calcd. for $\text{C}_8\text{H}_5\text{BF}_4\text{O}_3\text{W}$: C, 22.28; H, 1.20. Found: C, 23.63; H, 1.39.


*Properties**

*See Section B.

B. DICARBONYL(η^5 -CYCLOPENTADIENYL)(TETRAFLUOROBORATO)(TRIPHENYLPHOSPHINE)MOLYBDENUM AND TUNGSTEN, $\text{MCp}(\text{CO})_2(\text{PPh}_3)(\text{FBF}_3)$ ($\text{M} = \text{Mo, W}$)⁴

Substitution of a CO group by a phosphine ligand makes the metal center electron richer and therefore less Lewis acidic. This weakens the coordination of the $[\text{BF}_4]^-$ ion. In addition, steric interactions with the phosphine ligands, the possibility of cis-trans isomerism in the complexes with “four legged piano stool” geometry,⁵ and the introduction of the ^{31}P nucleus as another sensitive NMR probe make this variation of the synthesis described in Section A, an interesting field of further investigation. The preparation described here for the PPh_3 compounds, can also be used with other PR_3 ligands such as PMe_3 , PEt_3 , P(OPh)_3 , or $\frac{1}{2}(\text{dppe})$ [$\text{dppe} = 1,2$ -ethanediyl-bis(diphenylphosphine)].[†]

Dicarbonyl(η^5 -cyclopentadienyl)hydrido(triphenylphosphine)molybdenum and -tungsten



Monophosphine substituted carbonylcyclopentadienylhydrido complexes of molybdenum and tungsten have been obtained by protonation of the anions $[\text{MCp}(\text{CO})_2(\text{PR}_3)]^-$,⁶ or by substitution of CO with phosphines in the hydrides $\text{MH}(\text{CO})_3\text{Cp}$.⁷ The straightforward synthesis of the hydrides $\text{MH}(\text{CO})_3\text{Cp}$ ($\text{M} = \text{Mo, W}$)³ makes the latter procedure preferable, at least for PPh_3 , P(OPh)_3 , PMe_3 , and PEt_3 , where fast reactions and good yields can always be obtained. For the analogous syntheses of PMe_3 or PEt_3 substituted hydrides, special precautions for handling these highly toxic, malodorous, and highly inflammable phosphines must be taken.⁸

Procedure

A quantity of freshly sublimed $\text{MCp}(\text{CO})_3\text{H}$ (0.49 g, $\text{M} = \text{Mo}$ or 0.66 g, $\text{M} = \text{W}$, each 2.0 mmol) is dissolved in 15 mL of hexane at room temperature in a 100-mL Schlenk flask. To this is added PPh_3 (0.58 g, 2.1 mmol) under vigorous stirring. The Schlenk tube is then connected to a mercury bubbler and a stream of argon (0.5 L min^{-1}) is passed over the solution for 2 h (it is

[†]Commonly known as 1,2-bis(diphenylphosphino)ethane.


not necessary to bubble the argon through the solution). Soon a white precipitate forms, which is isolated by filtration under argon and washed twice with 5 mL of hexane. The product is dried at room temperature for 6 h *in vacuo*. It may be recrystallized from CH₂Cl₂–hexane. Yields: MoCp(CO)₂(PPh₃)H, 625 mg, 65%; WCp(CO)₂(PPh₃)H, 636 mg (56%).

Properties

The hydrides MCp(CO)₂(PPh₃)H are yellowish-white powders. They are air stable for several minutes exposure as solids, however, for extended storage they should be kept under argon. IR spectra (in CH₂Cl₂): ν_{CO} = 1936, 1856 cm⁻¹ (Mo); 1923, 1835 cm⁻¹ (W); ¹H NMR [in CD₂Cl₂ (Mo), CDCl₃ (W)]: $\delta_{(\text{C}_5\text{H}_5)}$ = 5.08 (Mo); 5.10 ppm (W); $\delta_{(\text{M}-\text{H})}$ = -5.56 (Mo), -7.06 ppm (W), “doublets” $^2J_{(^3\text{P}^1\text{H})_{\text{av}}}$ = 47 Hz (Mo), 55 Hz (W); ³¹P NMR (in CD₂Cl₂): $\delta_{(\text{PPh}_3)}$ = 74.3 (Mo), 40.9 ppm (W) (relative to H₃PO₄). A fast equilibrium between the cis and trans isomers⁵ leads to averaging of the signals and coupling constants at room temperature. Both isomers can be distinguished by low-temperature ¹H NMR [$\delta_{(\text{Mo}-\text{H})}$: -5.33d and -6.14d; $^2J_{(^3\text{P}^1\text{H})}$ = 64 and 21.4 Hz; $\delta_{(\text{W}-\text{H})}$: -6.90d, -7.36d, $^2J_{(^3\text{P}^1\text{H})}$ = 65 and 22 Hz].

The analogous PMe₃ and PEt₃ containing hydrides tend to form oils and decompose quickly on contact with air; the tungsten compounds are more stable than the molybdenum analogs. Their spectral properties are similar to those of the PPh₃ compounds. The best yields are obtained with the P(OPh)₃ ligand, which leads exclusively to the stable cis isomers.⁵

Dicarbonyl(η^5 -cyclopentadienyl)(tetrafluoroborato)(triphenylphosphine)-molybdenum and -tungsten⁴



Procedure

Generally, the same guidelines as described in Section A have to be followed. A quantity of MoCp(CO)₂(PPh₃)H (0.45 g, 0.94 mmol) or WCp(CO)₂(PPh₃)H (0.55 g, 0.97 mmol) is added to a solution of [Ph₃C][BF₄] (0.33 g, 1.00 mmol) in 10 mL CH₂Cl₂ at -40 °C contained in a Schlenk flask (100 mL) equipped with a magnetic stirring bar. The mixture is stirred for 20 min, after which an IR spectrum is recorded of an aliquot (0.2 mL) to inspect the intensity of the

band at 1355 cm^{-1} . Small amounts of the hydride are added until the IR spectrum, recorded at 5-min intervals shows no band at 1355 cm^{-1} . Usually, a lilac precipitate forms before the equivalence point is reached. The equivalence point is again indicated by a lilac color of the solution. Complete precipitation of the product is obtained by addition of 20 mL of hexane at -60°C , or by transfer of the complete reaction mixture to another Schlenk flask containing the hexane cooled to -60°C (see the procedure in Section A). Isolation of the product is the same as described in Section A. Yields: $\text{MoCp}(\text{CO})_2(\text{PPh}_3)(\text{FBF}_3) \cdot 2\text{CH}_2\text{Cl}_2$ 632 mg. (86%); $\text{WCp}(\text{CO})_2(\text{PPh}_3)(\text{FBF}_3)$ 556 mg (85%).

Anal. Calcd. for $\text{C}_{25}\text{H}_{20}\text{BF}_4\text{MoO}_2\text{P} \cdot 2\text{CH}_2\text{Cl}_2$: C, 44.1; H, 3.29. Found: C, 45.0; H, 3.32.

Properties

All tetrafluoroborato complexes are very sensitive to moisture. Schlenk tubes used for storage therefore have to be heated to 400°C or more under vacuum for several hours; O-ring stopcocks or similar grease-free stopcocks are superior to the usual ground glass stopcocks. Although the phosphine containing BF_4^- complexes are thermally more stable than the unsubstituted compounds, storage at temperatures below -25°C under Ar is recommended for all these compounds. They dissolve in CH_2Cl_2 and CHCl_3 below -40°C without decomposition, while solvents with donor properties like acetone or acetonitrile dissolve these complexes under substitution of tetrafluoroborato ligands by the solvent to give ionic complexes, for example, $[\text{MoCp}(\text{CO})_3(\text{acetone})][\text{BF}_4^-]$. They can be characterized by their IR spectra in the region from 1200 to 700 cm^{-1} and by their low-temperature ^{19}F and, where appropriate, ^{31}P NMR spectra.⁹

TABLE I. Spectroscopic Data of $\text{MoCp}(\text{CO})_2\text{L}(\text{FBF}_3)$.

$\text{L} = \text{CO}$
IR: $\nu_{(\text{CO})} = 2071, 1988\text{ cm}^{-1}$ (in CH_2Cl_2)
$\nu_{(^{11}\text{BF})} = 1130, 884, 722\text{ cm}^{-1}$ (in Nujol)
^{19}F NMR: ^a – 155d (MoFBF_3), – 370q (MoFBF_3), 95 Hz [$^2J_{(\text{F}-\text{F})}$]
$\text{L} = \text{PPh}_3$
IR: $\nu_{(\text{CO})} = 1991, 1903\text{ cm}^{-1}$ (in CH_2Cl_2)
$\nu_{(^{11}\text{BF})} = 1119, 901, 732\text{ cm}^{-1}$ (in Nujol)
^{19}F NMR: ^a – 155d (MoFBF_3), – 344q, – 391q (MoFBF_3), 90 Hz [$^2J_{(\text{F}-\text{F})}$]

^a δ in ppm, relative to CFCl_3 , in CD_2Cl_2 , -80°C .

TABLE II. Spectroscopic Data of $\text{WCp}(\text{CO})_2\text{L}(\text{FBF}_3)$.

$\text{L} = \text{CO}$
IR: $\nu_{(\text{CO})} = 2067, 1975 \text{ cm}^{-1}$ (in CH_2Cl_2)
$\nu_{(\text{FBF})} = 1149, 874, 704 \text{ cm}^{-1}$ (in Nujol)
^{19}F NMR: ^a — 153d (WFBF ₃), — 394q (WFBF ₃), 99 Hz [$^2J_{(\text{F}-\text{F})}$]
$\text{L} = \text{PPh}_3$
IR: $\nu_{(\text{CO})} = 1988, 1963, 1877 \text{ cm}^{-1}$ (in Nujol)
$\nu_{(\text{FBF})} = 1148, 887, 720 \text{ cm}^{-1}$ (in Nujol)
^{19}F NMR: ^b — 156d (WFBF ₃), — 371q (WFBF ₃), 98 Hz [$^2J_{(\text{F}-\text{F})}$].

^a δ in ppm, relative to CFCl_3 , in CD_2Cl_2 , —52°C.

^b δ in ppm, relative to CFCl_3 , in CD_2Cl_2 , —80°C.

Coordination of the $[\text{BF}_4]^-$ ion lowers the T_d symmetry of $[\text{BF}_4]^-$ and makes the fluorine atoms nonequivalent. Therefore the IR spectra show three instead of one $\nu_{(^{11}\text{B}-\text{F})}$ absorptions^{1,4} (Tables I and II); the low-temperature ^{19}F NMR spectra⁹ show two distinct fluorine resonances, a high-field quartet (which may be split by coupling to the phosphorus in the PR₃ substituted compounds and a doublet at lower field, close to the resonance of free $[\text{BF}_4]^-$;¹³ the ^{31}P NMR spectrum shows at low temperature a pseudodoublet, produced by coupling with the coordinated fluorine (Tables I and II). Compounds MoCp(CO)₂(PR₃)(FBF₃) are obtained as cis and trans isomers. In WCp(CO)₂(P(OPh)₃)(FBF₃) total isomerization from the pure cis hydride to the pure trans-BF₄ compound could be followed via NMR.⁹


Reactions of tetrafluoroborato complexes with ethylene, diphenylacetylene and acetone

General Remarks

The tetrafluoroborate ligand of these highly reactive complexes can be easily substituted by a series of N, O, P, and S σ donors^{1a,4,10,11} and π donors (see Section C–F).

As described in Sections A and B, a lilac solution of the corresponding tetrafluoroborato complex is prepared at —30°C in 10 mL of CH_2Cl_2 . Complete reaction of the trityaborptionium salt is verified by checking for the disappearance of the 1355 cm^{-1} band in the IR spectrum of the solution. This solution is used for the following reactions without isolation of the tetrafluoroborato complex.

C. TRICARBONYL(η^5 -CYCLOPENTADIENYL)(η^2 -ETHENE)-MOLYBDENUM(1+) TETRAFLUOROBORATE(1-)^{1a}



The title compound can be obtained in three ways. One method starts from $\text{Mo}(\text{C}_5\text{H}_5)(\text{CO})_3\text{Cl}$, which is reacted with C_2H_4 at a pressure of 70 bar in the presence of AlCl_3 and consecutive precipitation with ammonium salt.¹² A second method involves β -hydride abstraction from the ethyl group in $\text{MoCp}(\text{CO})_3(\text{C}_2\text{H}_5)$ by $[\text{Ph}_3\text{C}] [\text{BF}_4]$.¹³ The third method, described here, has the advantage of mild reaction conditions and a good overall yield. Analogous complexes with other olefins have been prepared similarly.⁴


Procedure

Ethylene (1 bar), dried over P_2O_5 , is bubbled through a vigorously stirred lilac solution of $\text{MoCp}(\text{CO})_3(\text{FBF}_3)$ (1 mmol) in CH_2Cl_2 in a Schlenk flask (50 mL) cooled to -30°C . With continuous ethylene bubbling, the cooling bath is removed and the flask is permitted to warm up to $+20^\circ\text{C}$ over a 4-h period. The flow of ethylene is then stopped, and the reaction mixture stirred under argon for another 30 min. The yellow precipitate is isolated by centrifugation or filtration under Ar. After washing four times with 5 mL of CH_2Cl_2 , the product is dried 1 h at $+40^\circ\text{C}$ on a high-vacuum line. It may be recrystallized from acetone-diethyl ether. Yield: 282 mg (79%).

Properties

The compound decomposes on heating at 102 to 108°C . Infrared spectra (in Nujol): $\nu_{(\text{CO})} = 2104, 2053, 2001 \text{ cm}^{-1}$; ^1H NMR (acetone- d_6) $\delta = 6.35 \text{ ppm}$ (C_5H_5).

D. CARBONYL(η^5 -CYCLOPENTADIENYL)BIS(DIPHENYL-ACETYLENE)MOLYBDENUM(1+) TETRAFLUOROBORATE(1-)^{1a}



Other syntheses of cationic bis(alkyne)complexes of molybdenum and tungsten of the same type include AgBF_4 oxidation of the dimer

[MoCp(CO)₃]₂ in CH₂Cl₂ in the presence of diphenylacetylene or several other alkynes.¹⁴ Protonation of MoCp(CO)₃CH₃ with CF₃COOH and consecutive addition of 2-butyne in acetonitrile, followed by precipitation with a methanolic solution of [NH₄][PF₆] gives the corresponding 2-butyne complex.¹⁵ Refluxing a solution of MoCp(CO)₃Cl with (HOCH₂)CC(CH₂OH) leads to an analogous compound.¹⁶ The method described here uses very mild conditions and can be applied also for other alkynes, like 2-butyne or acetylene.¹⁷

Procedure


Diphenylacetylene (535 mg, 3.0 mmol) is added to a lilac solution of MoCp(CO)₃FBF₃ (1 mmol in 10 mL CH₂Cl₂, prepared as described above) at -30 °C under vigorous stirring in a Schlenk flask (50 mL) equipped with a magnetic stirring bar. The flask is connected to a mercury bubbler and flushed by a constant flow of argon or nitrogen gas. After 30 min, the gas flow is stopped and the cooling bath removed. Stirring is continued for 4 days at room temperature, over which time a yellow precipitate is formed. Diethyl ether (20 mL) is added and the yellowish red suspension is filtered under argon. The residue on the frit is extracted three times with 10 mL of CH₂Cl₂. The combined extracts are evaporated to 5 mL, to which is added diethyl ether (20 mL). The orange-yellow precipitate is isolated by centrifugation or filtration under argon, washed three times with 10-mL aliquots of diethyl ether, and then dried for 1 h *in vacuo* at 40 °C. The product may be recrystallized from CH₂Cl₂-pentane. Yield: 235 mg (37%).

Anal. Calcd. for C₃₄H₂₅BF₄MoO: C, 64.58; H, 3.99. Found: C, 63.98; H, 4.09.

Properties

The yellow compound is soluble in polar solvents such as CH₂Cl₂, acetone, or acetonitrile. Although prolonged exposure to air leads to decomposition, the compound can be handled in air for short periods of time. Its IR spectrum in Nujol shows one $\nu_{12\text{CO}}$ vibration at 2088 cm⁻¹ and a weak $\nu_{13\text{CO}}$ band at 2040 cm⁻¹. Also a weak absorption at 1741 cm⁻¹ occurs, which may be due to the $\nu_{(\text{C}=\text{C})}$ band of the coordinated alkyne. The ¹H NMR spectrum in CH₂Cl₂ has a sharp singlet for the C₅H₅ protons at δ = 6.20 ppm, besides the broad resonance of the phenyl protons of the diphenylacetylene. Interestingly, KBr pellets of the compound several hours after initially formed, show a bathochromic shift of the $\nu_{(\text{CO})}$ band, which is also observed with other cationic alkyne complexes.¹⁷

E. CARBONYL(η^5 -CYCLOPENTADIENYL)(DIPHENYL-ACETYLENE)(TRIPHENYLPHOSPHINE)MOLYBDENUM(1+) TETRAFLUOROBORATE(1-) ¹⁸



Green and coworkers¹⁹ prepared the title compound and other related monoalkyne complexes by reaction of the corresponding bis(alkyne) complex with triphenylphosphine (or other phosphines) in good yields. The method described here works for several alkynes, for example, 2-butyne or phenyl-acetylene, and also for phosphines, for example, PEt₃ or P(OPh)₃.

Procedure

Diphenylacetylene (1.78 g, 10.0 mmol) is added to a magnetically stirred lilac suspension of MoCp(CO)₂PPh₃FBF₃ (1.0 mmol in 10 mL of CH₂Cl₂, as described previously) in a Schlenk flask (50 mL) cooled to -30 °C. The flask is connected to a mercury bubbler and purged with argon for 15 min. The gas flow is stopped and the cooling bath is allowed to warm up to room temperature. Stirring is continued for 2 days during which time the flask is purged several times with argon to remove the carbon monoxide evolved in the reaction. Then hexane (20 mL) is added. Stirring is continued for another day at ambient temperature, after which the dark green suspension is filtered under argon. The residue on the filter is washed four times with 15-mL aliquots of hexane and then dried 3 h under vacuum at 25 °C. Yield: 408 mg (57%).

Anal. Calcd. for C₃₈H₃₀BF₄MoOP: C, 63.7; H, 4.22. Found: C, 62.8; H, 4.15.

Properties

The title compound is soluble in polar organic solvents, for example, acetone, acetonitrile, or dichloromethane. Although storing under inert gas is recommended, no decomposition can be observed when handled as a solid in air for short periods of time. IR (in CH₂Cl₂): $\nu_{(\text{C}\equiv\text{C})} = 1987 \text{ cm}^{-1}$; ¹H NMR (CD₂Cl₂): $\delta_{(\text{C}_5\text{H}_5)} = 5.77 \text{ ppm}$, $\delta_{[\text{C}-\text{C}_6\text{H}_5,\text{P}(\text{C}_6\text{H}_5)_3]} = 8-7 \text{ ppm}$; ³¹P NMR (in CD₂Cl₂) $\delta_{(\text{PPh}_3)} = 54.8 \text{ ppm}$.

The crystal structure of this compound shows a slightly elongated C≡C bond of the alkyne and the usual deviation from linearity at the two carbon atoms of the triple bond.^{18,19}

F. (ACETONE)(TRICARBONYL)(η^5 -CYCLOPENTADIENYL)-
MOLYBDENUM(1+) AND -TUNGSTEN(1+)
TETRAFLUOROBORATE(1-) ^{1a,10}



M = Mo, W

Procedure

A lilac solution of the tetrafluoroborato complex $\text{MCp}(\text{CO})_3(\text{FBF}_3)$, M = Mo or W (1 mmol in 10 mL of CH_2Cl_2) is prepared as indicated above in a Schlenk flask (50 mL) and cooled to -30°C . To this is added acetone (0.1 mL, 1.38 mmol). An immediate color change to red occurs, and stirring is continued for 3 h. Hexane (15 mL) is then added. The solution is cooled to -78°C (Dry Ice) and stored overnight giving a dark red precipitate. This is isolated by centrifugation and washed twice with 19 mL of hexane at 0°C . Alternatively, the supernatant solution may be removed by a stainless steel cannula fitted with a sintered glass frit. The solids are washed with two aliquots of cold (0°C) hexane (19 mL), each removed by use of the stainless steel cannula fitted with the glass frit. The product is then dried for 6 h at 0°C on a high vacuum line. Yield: $[\text{MoCp}(\text{CO})_3(\text{OC}(\text{CH}_3)_2)][\text{BF}_4]$ 350 mg (90%); $[\text{WCp}(\text{CO})_3(\text{OC}(\text{CH}_3)_2)][\text{BF}_4]$ 420 mg (88%).

Anal. Calcd. for $\text{C}_{11}\text{H}_{11}\text{BF}_4\text{MoO}_4$: C, 33.88; H, 2.84. Found: C, 33.87; H, 2.85. Calcd. for $\text{C}_{11}\text{H}_{11}\text{BF}_4\text{O}_4\text{W}$: C, 27.65; H, 2.32. Found: C, 26.90; H, 2.45.

Similar acetone complexes can also be prepared from the PPh_3 containing tetrafluoroborate complexes.

Properties

Solutions of the compounds in CH_2Cl_2 or acetone decompose at 20°C within a short time, especially, when traces of water are present. The solid compounds can be stored under argon at -30°C for several weeks without decomposition.

IR (CH_2Cl_2): $\nu_{(\text{CO})} = 2072, 1987\text{ cm}^{-1}$ (Mo) IR (in Nujol): $\nu_{(\text{CO})} = 2050, 1930\text{ cm}^{-1}$ (W); $\nu_{(\text{M}-\text{O}-\text{C})} = 1660\text{ cm}^{-1}$ (Mo), 1640 cm^{-1} (W). ^1H NMR (in CD_2Cl_2): $\delta_{(\text{C}_5\text{H}_5)} = 6.11$ (Mo), 6.19 ppm (W); $\delta_{(\text{CH}_3)} = 2.39$ (Mo), 2.43 ppm (W).

References

1. (a) W. Beck and K. Schlöter, *Z. Naturforsch. Teil B*, **33**, 1214 (1978); (b) K. Sünkel, U. Nagel, and W. Beck, *J. Organomet. Chem.*, **251**, 227 (1983); (c) M. Appel, K. Schlöter, J. Heidrich, and W. Beck, *J. Organomet. Chem.*, **322**, 77 (1987).

2. J. R. Sanders, *J. Chem. Soc. Dalton Trans.*, **1972**, 1333.
3. (a) R. B. King and F. G. A. Stone, *Inorg. Synth.*, **7**, 107 (1963). E. O. Fischer, *Inorg. Synth.*, **7**, 136 (1963); (b) R. B. King, *Organometallic Syntheses*, Vol. 1, Academic Press, New York, 1965, p. 156; (c) W. P. Fehlhammer, W. A. Herrmann, and G. K. Öfsele, in *Handbuch der Präparativen Anorganischen Chemie*, Vol. 2, 3rd ed., F. Enke Verlag, Stuttgart, 1981.
4. K. Sünkel, H. Ernst, and W. Beck, *Z. Naturforsch. Teil B*, **36**, 474 (1980).
5. J. W. Faller and A. S. Anderson, *J. Am. Chem. Soc.*, **92**, 5852 (1970).
6. (a) A. R. Manning, *J. Chem. Soc. A*, **1968**, 651; (b) M. J. Mays and S. M. Pearson, *J. Chem. Soc. A*, **1968**, 2291.
7. (a) A. Bainbridge, P. J. Craig, and M. Green, *J. Chem. Soc. A*, **1968**, 2715; (b) P. Kalck, R. Pince, R. Poilblanc, and J. Roussel, *J. Organomet. Chem.*, **24**, 445 (1970).
8. R. T. Markham, E. A. Dietz, Jr., and D. R. Martin, *Inorg. Synth.*, **16**, 153 (1976).
9. K. Sünkel, G. Urban, and W. Beck, *J. Organomet. Chem.*, **252**, 187 (1983); M. Appel and W. Beck, *J. Organomet. Chem.*, **319**, C1 (1987).
10. K. Sünkel, G. Urban, and W. Beck, *J. Organomet. Chem.*, **290**, 231 (1985).
11. K. Schloter and W. Beck, *Z. Naturforsch. Teil B*, **35**, 985 (1980); G. Urban, K. Sünkel, and W. Beck, *J. Organomet. Chem.*, **290**, 329 (1985).
12. E. O. Fischer and K. Fichtel, *Chem. Ber.*, **94**, 1200 (1961).
13. M. Cousins and M. L. H. Green, *J. Chem. Soc.*, **1963**, 889.
14. M. Bottrill and M. Green, *J. Chem. Soc. Dalton Trans.*, **1977**, 2365.
15. P. L. Watson and R. G. Bergman, *J. Am. Chem. Soc.*, **102**, 2698 (1980).
16. J. W. Faller and H. H. Murray, *J. Organomet. Chem.*, **172**, 171 (1979).
17. K. Schloter, K. Sünkel and W. Beck, unpublished results.
18. K. Sünkel, U. Nagel, and W. Beck, *J. Organomet. Chem.*, **222**, 251 (1981).
19. S. R. Allen, P. K. Baker, S. G. Barnes, M. Green, L. Trollope, L. Manojlovic-Muir, and K. W. Muir, *J. Chem. Soc. Dalton Trans.*, **1981**, 873.