

## ION ACTIVITIES AND POTASSIUM UPTAKE MECHANISMS OF GLIAL CELLS IN GUINEA-PIG OLFACTORY CORTEX SLICES

By K. BALLANYI, P. GRAFE AND G. TEN BRUGGENCATE

*From the Institut für Physiologie der Universität München, Pettenkoferstrasse 12, D-8000 München 2, F.R.G.*

(Received 20 November 1985)

### SUMMARY

1. Double-barrelled ion-sensitive micro-electrodes were used to measure changes in the intracellular activities of  $K^+$ ,  $Na^+$  and  $Cl^-$  ( $a_K^i$ ,  $a_{Na}^i$ ,  $a_{Cl}^i$ ) in glial cells of slices from guinea-pig olfactory cortex during repetitive stimulation of the lateral olfactory tract.

2. Base-line levels of  $a_K^i$ ,  $a_{Na}^i$  and  $a_{Cl}^i$  were about 66, 25 and 6 mM, respectively, for cells with resting potentials higher than  $-80$  mV. During stimulation, intragial  $a_K^i$  and  $a_{Cl}^i$  increased, whereas  $a_{Na}^i$  decreased. Within about 2 min after stimulation the ion activities returned to their base-line levels.

3. The  $Cl^-$  equilibrium potential was found to be close to the membrane potential ( $E_m$ ). There was also a strong correlation between changes of  $E_m$  and  $a_{Cl}^i$ . These observations indicate a high  $Cl^-$  conductance of the glial cell membrane.

4. In the presence of  $Ba^{2+}$ , the usual depolarizing response of the glial cells to a rise of the extracellular  $K^+$  activity ( $a_K^e$ ) reversed into a membrane hyperpolarization. Furthermore,  $Ba^{2+}$  strongly reduced the stimulus-related rise of intragial  $a_K^i$ . An additional application of ouabain blocked both the membrane hyperpolarization as well as the remaining rise of  $a_K^i$ .

5. In conclusion, our data show that glial cells in guinea-pig olfactory cortex slices possess at least two mechanisms of  $K^+$  accumulation. One mechanism is sensitive to the  $K^+$  channel blocker  $Ba^{2+}$  and might be a passive KCl influx. The other appears to be the electrogenic  $Na^+/K^+$  pump, which can be activated by excess extracellular  $K^+$ .

### INTRODUCTION

Enhanced neuronal activity in the central nervous system is accompanied by transient elevations in extracellular  $K^+$  ( $a_K^e$ ; Somjen, 1979; Nicholson, 1981). Alterations of  $a_K^e$ , however, can modify the activity of neurones (Sykova, 1983). Therefore, it is of interest to investigate the mechanisms of  $K^+$  homeostasis in the brain. Glial cells surrounding the neurones seem to be involved in the regulation of  $a_K^e$  (Walz & Hertz, 1983a). Intracellular recordings with ion-sensitive micro-electrodes (Coles & Tsacopoulos, 1979; Coles & Orkand, 1983; Kettenmann, Sonnhof & Schachner, 1983; Schlue & Wuttke, 1983) and radiotracer studies (Walz & Hertz, 1982) have revealed the ability of glial cells to accumulate  $K^+$  during an increase of

$a_{\text{K}}^{\text{c}}$ . Several mechanisms such as spatial buffer (Orkand, Nicholls & Kuffler, 1966; Dietzel, Heinemann, Hofmeier & Lux, 1980, 1982; Coles & Orkand, 1983; Gardner-Medwin, 1983),  $\text{Na}^+/\text{K}^+$  pump (Kukes, Elul & de Vellis, 1976; Walz & Hertz, 1982; Grisar, Franck & Delgado-Escueta, 1983), KCl co-transport (Walz & Hertz, 1984; Wolpaw & Martin, 1984) or  $\text{Ba}^{2+}$ -sensitive  $\text{K}^+$  channels (Walz, Shargool & Hertz, 1984a) have been proposed to contribute to the  $\text{K}^+$  uptake process. Most of the data was obtained from glial cell lines or from preparations of invertebrates. However, no recordings have been described concerning ion activities in glial cells of functionally intact mammalian cortex. Therefore, the first aim of the present study was to investigate changes in the intraglial activities of  $\text{K}^+$ ,  $\text{Na}^+$  and  $\text{Cl}^-$  ( $a_{\text{K}}^{\text{i}}$ ,  $a_{\text{Na}}^{\text{i}}$  and  $a_{\text{Cl}}^{\text{i}}$ ) using double-barrelled ion-sensitive micro-electrodes in guinea-pig olfactory cortex slices during stimulation of cortical afferents. The second aim was to characterize the ionic mechanisms responsible for the increase in  $a_{\text{K}}^{\text{i}}$  observed during stimulation. With the aid of  $\text{Ba}^{2+}$  and ouabain, it was possible to discriminate between two mechanisms of  $\text{K}^+$  accumulation. One component seems to operate through  $\text{K}^+$  channels which can be blocked by  $\text{Ba}^{2+}$ . The other component appears to be the ouabain-sensitive electrogenic  $\text{Na}^+/\text{K}^+$  pump. Parts of the results have been presented at a meeting of the German Physiological Society (Ballanyi, Grafe & ten Bruggencate, 1985; Grafe, 1985).

#### METHODS

##### *Preparation and solutions*

Experiments were performed on slices of the olfactory cortex of guinea-pigs. After decapitation, the brain was rapidly removed and surface slices (approximately  $500\ \mu\text{m}$  thick) were cut using a plastic guide and a bow cutter. Slices were placed in a Perspex chamber (volume 2 ml) and superfused at  $25\ ^\circ\text{C}$  with a solution of the following composition (in mM): NaCl, 118; KCl, 3;  $\text{NaHCO}_3$ , 25;  $\text{NaH}_2\text{PO}_4$ , 1.2;  $\text{MgCl}_2$ , 1.0;  $\text{CaCl}_2$ , 1.5, and glucose, 10 (gassed with 95%  $\text{O}_2$ , 5%  $\text{CO}_2$ ; pH 7.4). The  $\text{Cl}^-$ -free solution was composed of (in mM): Na gluconate, 118; K gluconate, 3;  $\text{NaHCO}_3$ , 25;  $\text{NaH}_2\text{PO}_4$ , 1.2;  $\text{MgSO}_4$ , 1; Ca gluconate, 6.5. Drugs were added to the superfusion fluid (all chemicals in this study were purchased from Sigma; Munich, F.R.G.). Cortical neurones were activated by electrical stimulation of the lateral olfactory tract (l.o.t.) via platinum wire electrodes.

##### *Ion-sensitive micro-electrodes*

The methods used for the construction and calibration of the double-barrelled ion-sensitive micro-electrodes are described in detail elsewhere (Ballanyi, Grafe, Reddy & ten Bruggencate, 1984; Grafe, Ballanyi & ten Bruggencate, 1985). In brief, theta-capillaries were pulled with a Brown-Flaming type micro-electrode puller (Sutter Instrument Co., San Francisco, CA, U.S.A.) resulting in tip diameters below  $0.3\ \mu\text{m}$ . Silanization of one barrel with hexamethyldisilazane (Sigma, Munich, F.R.G.) was performed at  $400\ ^\circ\text{C}$  for 20 min. During this period, nitrogen was applied under pressure to the other barrel in order to prevent the silane from entering. Later on, a drop of liquid ion exchanger was injected into the silanized barrel, which was then backfilled with an internal reference solution. The reference barrel was filled with electrolyte solution. The following combinations of ion exchanger, backfilling solution and reference barrel solution were used:  $\text{K}^+$ -sensitive micro-electrodes (Corning 477317, 200 mM-KCl, 1 M-Mg acetate);  $\text{Na}^+$ -sensitive micro-electrodes (Fluka 71176, 200 mM-NaCl; 1 M-Mg acetate);  $\text{Cl}^-$ -sensitive micro-electrodes (WP Instruments IE-170, 200 mM-KCl, 0.5 M-K sulphate). There was no significant difference in the average membrane potential of the glial cells as determined by electrodes containing 1 M-Mg acetate, 3 M-KCl (single-barrelled conventional electrodes) or 0.5 M-K sulphate in the reference barrel (see Results). This indicates that the interpretation of the data obtained with the  $\text{Cl}^-$ -sensitive micro-electrodes is not seriously hindered by junction potentials of the sulphate-containing electrodes.

K<sup>+</sup> electrodes were calibrated in solutions containing 3, 12 and 60 mM-KCl with a constant background of 150 mM-NaCl. The mean values for the slope and the selectivity coefficient *versus* Na<sup>+</sup> were  $53.3 \pm 4.6$  mV and  $0.019 \pm 0.007$  (mean  $\pm$  s.d.,  $n = 29$ ). The Cl<sup>-</sup> electrodes were calibrated either in pure KCl solutions (15, 40, 80 and 150 mM) or in solutions containing KCl (1, 10, 40 or 100 mM) and Na glucuronate in order to keep the ionic strength constant. The average electrode slope in both cases was  $50.2 \pm 3.1$  mV ( $n = 25$ ). Intracellular Na<sup>+</sup> was determined by comparing the voltage of the ion signal obtained upon withdrawal of the electrode from the cell into the bathing solution with the voltage obtained upon changing the normal Krebs solution to solutions containing different Na<sup>+</sup> concentrations (1, 3, 12, 30 and 60 mM) with a variable background of K<sup>+</sup> to keep the sum of Na<sup>+</sup> and K<sup>+</sup> at 150 mM. The Ca<sup>2+</sup> concentration in these solutions was buffered to  $10^{-7}$  M. All values of intracellular Na<sup>+</sup>, K<sup>+</sup> and Cl<sup>-</sup> are given in activities assuming an intracellular activity coefficient of 0.74 for these ions (Meier, Ammann, Morf & Simon, 1980). Intracellular impalements were achieved by means of a piezo-driven micromanipulator (built by M. Frankenberger, Munich, F.R.G.).

## RESULTS

### General observations

The *in vitro* slice preparation of guinea-pig olfactory cortex is well suited to investigation of neurone-glia interactions in the mammalian central nervous system. On the one hand, it is possible to maintain stable intragial recordings either with conventional single-barrelled (Grafe, Reddy, Emmert & ten Bruggencate, 1983) or double-barrelled ion-sensitive micro-electrodes (Ballanyi *et al.* 1985). On the other hand, neuronal K<sup>+</sup> release can be elicited by electrical stimulation of afferent nerve fibres in the lateral olfactory tract (l.o.t.).

Membrane resting potentials ( $E_m$ ) of presumed glial cells were between  $-75$  and  $-95$  mV (no difference between single-barrelled conventional or double-barrelled ion-sensitive micro-electrodes). Such cells were characterized by the absence of any 'injury discharge' after the impalement and by the lack of synaptic potentials, which are the typical characteristics of neurones. Furthermore, repetitive stimulation of the l.o.t. (10–50 Hz, 5–50 s) resulted in a rapid depolarization of up to 35 mV amplitude with kinetics similar to the stimulus-induced increase of extracellular K<sup>+</sup> concentration. Measurements from glial cells in this study were discarded if  $E_m$  was less than  $-75$  mV, or when  $E_m$  or the ion signal were not stable for at least 10 min.

### Intracellular K<sup>+</sup> activity

The intracellular K<sup>+</sup> activity ( $a_K^i$ ) as measured in twelve cells with a mean  $E_m$  of  $-84.4 \pm 3.3$  mV was  $65.9 \pm 6.6$  mM (mean  $\pm$  s.d.). Stimulation of the l.o.t. (20 Hz, 15 s) in these cells resulted in an average  $a_K^i$  increase of  $16.1 \pm 6.1$  mM. Fig. 1 gives an example of the measurement of  $a_K^i$  in a cortical glial cell. In this experiment, one K<sup>+</sup>-sensitive micro-electrode was used to record intracellularly from a glial cell (Fig. 1A), whereas another was positioned extracellularly, close to the impaled cell (Fig. 1B). At the beginning of the intracellular recording, the base-line level of intragial  $a_K^i$  increased, a typical observation in the 'sealing phase' of the membrane. Later on, the l.o.t. was stimulated every 3 min with pulse trains of constant frequency (30 Hz) and variable duration (5–50 s). Each of these stimuli resulted in a rise of both  $a_K^e$  as well as  $a_K^i$ . Longer-lasting stimuli induced larger increases in intragial  $a_K^i$ . After each period of repetitive stimulation,  $a_K^i$  returned to the base line within about 2 min. Fig. 1B shows the recording of  $a_K^e$  as measured with the extracellularly positioned

K<sup>+</sup>-sensitive micro-electrode. It is apparent that there is a close relationship in the kinetics of changes in  $a_K^e$  and the glial membrane potential. However, the longer-lasting stimuli (30 and 50 s) were followed by transient after-hyperpolarizations of the glial  $E_m$ , whereas this behaviour was not mirrored by an 'undershoot' of  $a_K^e$ . To analyse this discrepancy between glial  $E_m$  and  $a_K^e$ , which was observed in several preparations,  $a_K^i$ ,  $a_K^e$ , glial  $E_m$ , and the calculated K<sup>+</sup> equilibrium potential were compared during tetanic stimulation.




Fig. 1. Effects of repetitive stimulation on intragial ( $a_K^i$ ) and extracellular K<sup>+</sup> activity ( $a_K^e$ ). In this Figure, the recordings of an intracellularly (A) and extracellularly positioned (B) K<sup>+</sup>-sensitive micro-electrode are illustrated during repetitive stimulation of the lateral olfactory tract with stimulation trains of constant frequency (30 Hz) and different durations (indicated by bars).  $E_m$  is the membrane potential of the glial cell,  $E_{ref}$  the voltage reading of the reference channel of the extracellular double-barrelled electrode. The voltage deflexions on the  $E_{ref}$  trace represent extracellular field potentials. These potentials did not affect the  $a_K^e$  measurements as the  $a_K^e$  signal is the difference of the voltage reading of the reference and the K<sup>+</sup>-sensitive barrel. At the beginning of the record, a voltage pulse of 20 mV (20 mV d.c.) was added to the ground potential in order to test the gain adjustment of the differential amplifiers.

In Fig. 2A, the 30 s stimulation period of Fig. 1 is illustrated on an extended time scale. First, it can be seen more clearly that the glial membrane potential shows a post-tetanic hyperpolarization, although no  $a_K^e$  undershoot as a possible cause for this behaviour was detectable. However, a calculation of the K<sup>+</sup> equilibrium potential ( $E_K$ ) based on both changes in  $a_K^e$  as well as  $a_K^i$ , reveals a good correlation between the glial  $E_m$  and  $E_K$ . Therefore, post-tetanic glial hyperpolarization in this situation can be explained by the still elevated  $a_K^i$  in this period (see also Kettenmann *et al.* 1983). A contribution of the electrogenic Na<sup>+</sup>/K<sup>+</sup> pump to the post-tetanic hyperpolarization is not necessarily involved since the behaviour of the membrane can be adequately explained on the basis of a high K<sup>+</sup> conductance. Application of




Fig. 2. Changes of ion activities and ionic equilibrium potentials. *A*, the 30 s stimulation period of Fig. 1 is illustrated on an extended time scale. Measured values of  $a_{\text{K}}^{\text{i}}$  and  $a_{\text{K}}^{\text{e}}$  were used to calculate the  $\text{K}^+$  equilibrium potential ( $E_{\text{K}}$ ; open circles) and to compare it with  $E_{\text{m}}$ . *B*, in this illustration, the  $\text{Cl}^-$  equilibrium potential ( $E_{\text{Cl}}$ ; open circles) before, during, and after repetitive stimulation of the l.o.t. was calculated using the measured value of the intracellular  $\text{Cl}^-$  activity ( $a_{\text{Cl}}^{\text{i}}$ ) and the  $\text{Cl}^-$  activity in the bathing solution. For further explanations see text.

ouabain is not a good tool to differentiate between these possibilities since it would modify both the rise of  $a_{\text{K}}^{\text{i}}$  as well as the pump-related membrane current. The good correlation between glial  $E_{\text{m}}$  and  $E_{\text{K}}$  also excludes the possibility that some other substance may have contributed to the voltage reading of the Corning ion exchanger-based  $\text{K}^+$  electrodes. Such a situation has been shown to interfere with measurements of  $a_{\text{K}}^{\text{i}}$  in glial cells of leech ganglia (Schlue & Wuttke, 1983). Secondly, it can also be seen that  $a_{\text{K}}^{\text{i}}$  rises more slowly with respect to the increase of  $a_{\text{K}}^{\text{e}}$ . This difference might be due to the kinetics of the  $\text{K}^+$  uptake mechanisms or due to time elapsed during intracellular diffusion. It is certainly not the result of a slow electrode response as most of the intracellularly used  $\text{K}^+$ -sensitive micro-electrodes responded very quickly when used extracellularly.

#### Intracellular $\text{Cl}^-$ activity

The intracellular  $\text{Cl}^-$  activity ( $a_{\text{Cl}}^{\text{i}}$ ) was measured in twenty-three glial cells with a mean  $E_{\text{m}}$  of  $-86.9 \pm 3.8$  mV (mean  $\pm$  s.d.). On the assumption of a pure  $\text{Cl}^-$  selectivity of the ion-sensitive micro-electrodes, an apparent  $a_{\text{Cl}}^{\text{i}}$  of  $6.0 \pm 1.5$  mM (mean  $\pm$  s.d.) and a  $\text{Cl}^-$  equilibrium potential ( $E_{\text{Cl}}$ ) of  $-71 \pm 5.9$  mV was calculated. During repetitive stimulation of the l.o.t. (20–30 Hz, 15 s)  $a_{\text{Cl}}^{\text{i}}$  increased by  $3.6 \pm 1.2$  mM (mean  $\pm$  s.d.,  $n = 19$ ). As an example, Fig. 3 illustrates an experiment in which a  $\text{Cl}^-$ -sensitive micro-electrode was used to follow changes in intragial  $a_{\text{Cl}}^{\text{i}}$  when the l.o.t. was stimulated with pulse trains of constant frequency (30 Hz) and variable duration (5–50 s). Based on the known extracellular  $\text{Cl}^-$  activity and the measured  $a_{\text{Cl}}^{\text{i}}$ ,  $E_{\text{Cl}}$  was then calculated. This calculation revealed a discrepancy




Fig. 3. Effects of repetitive stimulation on intragial  $\text{Cl}^-$  activity ( $a_{\text{Cl}}^i$ ). In this experiment, an intracellular  $\text{Cl}^-$ -sensitive micro-electrode was used to measure changes in  $a_{\text{Cl}}^i$  during repetitive stimulation of the l.o.t. with stimulation trains of constant frequency (30 Hz) and different durations (indicated by bars). At the beginning of the record, a voltage pulse of 20 mV (20 mV d.c.) was added to the ground potential in order to test the gain adjustment of the differential amplifier.  $V_{\text{Cl}}$  is the difference signal of the  $\text{Cl}^-$ -sensitive micro-electrode.

between resting glial  $E_m$  and  $E_{\text{Cl}}$  of around 15 mV (see Fig. 2B) and would indicate that either  $a_{\text{Cl}}^i$  is about 3 mM above a passive distribution or that interfering anions contribute to the voltage reading of the  $\text{Cl}^-$ -sensitive ligand. In order to differentiate between these two possibilities, experiments in  $\text{Cl}^-$ -free bathing solution were performed. A typical example of such an experiment is shown in Fig. 4.

In this experiment, one  $\text{Cl}^-$ -sensitive micro-electrode was positioned extracellularly about  $100 \mu\text{m}$  below the pial surface, and a second was placed intracellularly in an adjacent glial cell to measure  $a_{\text{Cl}}^i$ . The apparent  $a_{\text{Cl}}^i$  at the resting potential was about 7.4 mM, indicating a  $\text{Cl}^-$  equilibrium potential of  $-64$  mV and therefore 16 mV above  $E_m$  ( $-80$  mV). However, 8 min after the replacement of the extracellular  $\text{Cl}^-$  with gluconate,  $a_{\text{Cl}}^i$  was still at 4.2 mM in spite of a fall of  $a_{\text{Cl}}^e$  to about the same value. This observation leads us to conclude that some other anions, to which the  $\text{Cl}^-$  ligand is sensitive, result in an apparent intracellular  $\text{Cl}^-$  equivalent of about 3–4 mM. If one subtracts about 3 mM from the average apparent  $a_{\text{Cl}}^i$  of 6 mM, then an  $a_{\text{Cl}}^i$  of 3.2 mM would result in an equal  $E_{\text{Cl}}$  and  $E_m$  of  $-86$  mV each. Direct proof of the contribution of interfering anions to the voltage reading of  $\text{Cl}^-$ -sensitive micro-electrodes (liquid exchanger-based) was recently obtained in skeletal muscle by means of comparison of data obtained with recessed-tip solid-state  $\text{Ag}/\text{AgCl}$  electrodes (McCaig & Leader, 1984). It seems unlikely, although not impossible, that inwardly directed  $\text{Cl}^-$  transport mechanisms can maintain  $a_{\text{Cl}}^i$  at about 4 mM in an almost  $\text{Cl}^-$ -free extracellular solution. (The almost complete absence of extracellular  $\text{Cl}^-$




Fig. 4. Effect of a  $Cl^-$ -free solution on intragial  $Cl^-$  activity. In this experiment, the recordings of an extracellularly (A) and an intracellularly positioned (B)  $Cl^-$ -sensitive micro-electrode are illustrated during three trains of stimuli in normal extracellular bathing solution and during three trains of stimuli after the extracellular  $Cl^-$  was completely replaced by gluconate. Note that the apparent  $a_{Cl}^i$  did not decrease below 4 mM at an  $a_{Cl}^e$  of about the same value. This reveals the importance of intracellular interfering anions. For further explanation see text.

around the glial cell is also indicated by the lack of the usual stimulus-related rise of  $a_{Cl}^i$ .)

#### Intracellular $Na^+$ activity

The mean intracellular  $Na^+$  activity as measured in twelve glial cells with a mean  $E_m$  of  $-83.3 \pm 2.8$  mV was  $25.2 \pm 4.6$  mM (mean  $\pm$  S.D.). Stimulation of the l.o.t. (20 Hz, 15 s) resulted in an average decrease of  $a_{Na}^i$  in these cells of  $6.0 \pm 1.9$  mM. In the experiment illustrated in Fig. 5, a  $Na^+$ -sensitive micro-electrode was used to follow changes of intragial  $a_{Na}^i$  when the l.o.t. was stimulated with pulse trains of constant frequency and variable duration (5–50 s). Each of these stimuli resulted in a decrease of  $a_{Na}^i$ . After the end of the stimulation periods,  $a_{Na}^i$  returned to its resting level within about 2 min. The mechanisms underlying the stimulus-related decrease of glial  $a_{Na}^i$  were not studied in detail in the present paper. One important factor, however, seems to be the activation of the  $Na^+/K^+$  pump (see Discussion).

#### Effects of $Ba^{2+}$

In order to differentiate between several possible mechanisms of  $K^+$  uptake into cortical glia, the action  $Ba^{2+}$  was explored.  $Ba^{2+}$  is known to block  $K^+$  channels (Hille, 1984) and has been previously used as a tool to differentiate between passive and active  $K^+$  uptake mechanisms in skeletal muscle fibres (Sjodin & Ortiz, 1975).




Fig. 5. Effects of repetitive stimulation on intragial  $\text{Na}^+$  activity ( $a_{\text{Na}}^i$ ). In this recording, a double-barrelled  $\text{Na}^+$ -sensitive micro-electrode was used to measure changes in  $a_{\text{Na}}^i$  and membrane potential ( $E_m$ ) during repetitive stimulation of the l.o.t. with stimulation trains of constant frequency (20 Hz) and different durations. At the beginning of the recording, a d.c. voltage pulse (20 mV) was added to the ground potential in order to test the gain adjustment of the differential amplifier.




Fig. 6. Effects of  $\text{Ba}^{2+}$  and ouabain on extracellular  $\text{K}^+$  activity ( $a_{\text{K}}^e$ ) and glial membrane potential ( $E_m$ ). After three trains of stimuli (20 Hz, 10 s) had revealed the typical relationship between changes of  $a_{\text{K}}^e$  and  $E_m$ ,  $\text{Ba}^{2+}$  (0.5 mM) was added. This resulted in a decrease of  $a_{\text{K}}^e$  base-line and a glial depolarization. Stimulation under these conditions now resulted in a glial hyperpolarization (see central inset), whereas the stimulus-induced  $a_{\text{K}}^e$  increases were similar to the control situation. Subsequent addition of ouabain (5  $\mu\text{M}$ ) resulted in a rise of  $a_{\text{K}}^e$ , a further glial depolarization and the disappearance of the previously observed stimulus-related membrane hyperpolarization.

These ions are also known to interfere with the  $K^+$  homeostasis of cultured glial cells (Walz *et al.* 1984a). Fig. 6 illustrates a typical experiment, in which the effects of repetitive stimulation of the l.o.t. on the membrane potential of a glial cell and on  $a_K^e$  were investigated in the presence of  $Ba^{2+}$ . After three trains of stimuli had revealed the typical relationship between changes of  $a_K^e$  and the glial membrane




Fig. 7. Effects of  $Ba^{2+}$  on stimulus-induced increases in intracellular  $K^+$  activity ( $a_K^i$ ), extracellular  $K^+$  ( $a_K^e$ ) and glial membrane potential ( $E_m$ ). The Figure shows three sections taken from a continuous recording from a single glial cell. The left column shows the typical rise of  $a_K^i$  during a stimulus-induced increase of  $a_K^e$  in normal solution. In the presence of  $Ba^{2+}$  (0.5 mM, central column), an even greater rise in  $a_K^e$  led to a much smaller increase of  $a_K^i$ , and was accompanied by a membrane hyperpolarization. Right column shows a stimulation during the recovery period from  $Ba^{2+}$ .

potential,  $Ba^{2+}$  (0.5 mM) was added to the normal bathing solution. This resulted in a decrease of the  $a_K^e$  base line (probably due to reduced resting  $K^+$  efflux from neurones and glia; see also Nicholson, ten Bruggencate & Senekowitsch, 1976) and a glial membrane depolarization. The most striking effect, however, was a complete reversal in the relationship between stimulus-related rise of  $a_K^e$  and the behaviour of the glial membrane potential. In spite of a stimulus-induced rise in  $a_K^e$ , which normally resulted in a membrane *depolarization*, the glia now responded with a prominent membrane *hyperpolarization*. Only at the onset of the stimulus, a short-lasting, transient depolarization was visible (see central inset in Fig. 6). With the idea in mind that the  $K^+$ -induced hyperpolarization may be the consequence of an electrogenic  $K^+$  transport mechanism, ouabain was added to the  $Ba^{2+}$ -containing bathing solution. This procedure resulted in a rise of  $a_K^e$ , a further glial membrane depolarization and a disappearance of the previously observed stimulus-related membrane hyperpolarization (see right inset in Fig. 6).

The next three Figures illustrate the behaviour of the intragial ion activities in the presence of  $Ba^{2+}$ . The first Figure of this series (Fig. 7) illustrates that  $Ba^{2+}$

largely diminished the stimulus-induced rise of  $a_K^i$ . This reduction was evident despite an even bigger increase in  $a_K^e$  that occurred during the l.o.t. stimulation. Additional experiments revealed that this  $Ba^{2+}$ -resistant rise of  $a_K^i$  was completely blocked by ouabain (not illustrated). Another aspect concerns the resting  $a_K^i$  base line of glial cells in the presence of  $Ba^{2+}$  (Fig. 8). In the seven cells, in which  $Ba^{2+}$  (0.5–1 mM) was tested, an increase in  $a_K^i$  of between 5 and 23 mM ( $15.3 \pm 6.3$  mM,




Fig. 8. Effects of  $Ba^{2+}$  on extra- and intracellular  $K^+$  activities ( $a_K^e$ ,  $a_K^i$ ). In this Figure, the recordings of an extracellularly (A) and an intracellularly positioned (B)  $K^+$ -sensitive micro-electrode are illustrated. At the beginning of the recording, one period of tetanic stimulation (30 Hz, 15 s) revealed the typical glial depolarization accompanied by transient elevations in  $a_K^e$  and  $a_K^i$ . Application of  $Ba^{2+}$  (1 mM) resulted in a membrane depolarization of about 25 mV and an  $a_K^i$  increase of about 20 mM. This  $a_K^i$  increase was mirrored by a fall in  $a_K^e$  base line. Note that after wash-out of  $Ba^{2+}$ , the incomplete return of  $a_K^i$  to base line may be due to an improvement of the cell, as indicated by an increase of  $E_m$ .

mean  $\pm$  s.d.) was seen. A decrease in  $a_K^i$  was never observed, indicating that  $Ba^{2+}$  in the concentration of 0.5–1 mM does not inhibit the  $Na^+/K^+$  pump of the glial cells. This observation does not support an assumption of Walz *et al.* (1984a), who tried to explain the  $Ba^{2+}$ -induced depolarization of glial cells by an inhibition of the  $Na^+/K^+$  pump.

The behaviour of intragial  $a_{Cl}^i$  in the presence of  $Ba^{2+}$  indicates the close correspondence between the glial membrane potential and  $a_{Cl}^i$ . In the experiment illustrated in Fig. 9, a  $Cl^-$ -sensitive micro-electrode was used to record  $a_{Cl}^i$  and  $E_m$  of a glial cell (Fig. 9A), whereas a  $K^+$ -sensitive micro-electrode was positioned extracellularly to measure  $a_K^e$  (Fig. 9B). Two stimulus trains in normal solution at the beginning of this recording revealed the stimulus-related rise of  $a_K^e$ , the membrane depolarization and the accompanying rise of  $a_{Cl}^i$  usually seen in the glial cells.  $Ba^{2+}$  was added to the bathing solution and within a few minutes the membrane




Fig. 9. Effects of  $Ba^{2+}$  on intracellular  $Cl^-$  activity ( $a_{Cl}^i$ ) and extracellular  $K^+$  ( $a_K^e$ ). After two control stimulation trains,  $Ba^{2+}$  (0.5 mM) was added to the bathing solution. The typical glial depolarization in  $Ba^{2+}$  was accompanied by an increase in  $a_{Cl}^i$  base line. The  $Ba^{2+}$ -induced shift of a stimulus-related depolarization into a hyperpolarization was mirrored by a shift from an  $a_{Cl}^i$  increase to an  $a_{Cl}^i$  decrease. After the wash-out of  $Ba^{2+}$ , the membrane potential as well as  $a_{Cl}^i$  recovered to their base-line values.

depolarized by more than 20 mV and  $a_{Cl}^i$  increased to about 12 mM. Repetitive stimulation of the l.o.t. during  $Ba^{2+}$  induced a small membrane hyperpolarization and a slight fall in  $a_{Cl}^i$ . Therefore, the increase in resting  $a_{Cl}^i$  as well as the stimulus-related decrease of  $a_{Cl}^i$  clearly indicate the close relationship between the membrane potential and  $a_{Cl}^i$ . If  $a_{Cl}^i$  were to follow changes in  $a_K^e$ , then the  $Ba^{2+}$ -induced decrease of the  $a_K^e$  base line and the stimulus-related  $a_K^e$  increase should have altered  $a_{Cl}^i$  in the opposite direction.

#### DISCUSSION

##### *Base-line levels of intragial ion activities*

At the beginning of the discussion, the observed resting levels of intracellular ion activities need consideration. With respect to  $K^+$ , the measured values of  $a_K^i$  and the known level of the extracellular  $K^+$  concentration reveal an  $E_K$  close to  $E_m$ . This confirms previous findings of intragial  $K^+$  activity in glial cells (Kettenmann *et al.* 1983) and is closely related to the high  $K^+$  conductance of glial cells (Orkand *et al.* 1966). The base-line level of  $a_{Na}^i$  (25 mM on an average) is higher than  $a_{Na}^i$  determined in frog motoneurons (7.5 mM, Grafe, Rimpel, Reddy & ten Bruggencate, 1982) or rat sympathetic ganglion cells (5 mM, Ballanyi *et al.* 1984). This confirms a previous report describing higher intracellular  $Na^+$  levels in glia as compared to neurons

(Coles & Orkand, 1985). However, for the accurate determination of  $a_{\text{Na}}^i$  base-line levels, the  $\text{Ca}^{2+}$  sensitivity of the  $\text{Na}^+$  ligand has to be taken into account. Our  $\text{Na}^+$  electrodes were calibrated under the assumption of an intragial free  $\text{Ca}^{2+}$  concentration of  $10^{-7}$  M (see Methods). This  $\text{Ca}^{2+}$  background level might be too low since, very recently, Morris, Krnjević & MacDonald (1985) as well as Coles & Orkand (1985) have determined free intragial  $\text{Ca}^{2+}$  concentrations ( $[\text{Ca}^{2+}]_i$ ) between  $10^{-6}$  and  $10^{-5}$  M. If  $[\text{Ca}^{2+}]_i$  would be  $10^{-5}$  M, a calculation reveals that the voltage reading of our  $\text{Na}^+$  electrodes would indicate an  $a_{\text{Na}}^i$  of 20 mM instead of 25 mM (calculation based on a selectivity ratio for  $\text{Ca}^{2+}:\text{Na}^+:\text{K}^+ = 2.5:1:0.025$ , background 90 mM- $\text{K}^+$ , 25 mM- $\text{Na}^+$ ; Meier, Lanter, Ammann, Steiner & Simon, 1982). Such a result would not change the conclusion about a relatively high  $a_{\text{Na}}^i$  in the glial cells.

The  $\text{Cl}^-$ -sensitive micro-electrodes revealed an intragial  $a_{\text{Cl}}^i$  base-line level of about 6 mM and a chloride equilibrium potential slightly positive to the resting potential. However, experiments using low  $\text{Cl}^-$  concentrations in the extracellular bathing solution (see Fig. 4) indicate that other intracellular anions to which the liquid exchanger is sensitive may result in a voltage reading of the  $\text{Cl}^-$ -sensitive micro-electrodes equivalent to 3–4 mM- $\text{Cl}^-$ . The subtraction of these 3–4 mM- $\text{Cl}^-$  equivalents reveals a nearly equal  $E_{\text{Cl}}$  and  $E_m$  of the glial cell membrane. This result is in accordance with two other studies in which ion-sensitive micro-electrodes were used to measure  $a_{\text{Cl}}^i$  in glial cells (Bührle & Sonnhof, 1983; Coles & Orkand, 1984). An almost identical  $E_{\text{Cl}}$  and  $E_m$  together with the observation of a strong correlation between changes in membrane potential and  $a_{\text{Cl}}^i$  (see Fig. 9) indicates a high  $\text{Cl}^-$  conductance of the membrane. Later in the Discussion we will use this conclusion to explain the stimulus-related rise of intragial  $a_{\text{Cl}}^i$ . One experiment, on the other hand, apparently contradicts the view of a high  $\text{Cl}^-$  conductance. A reduction of the extracellular  $\text{Cl}^-$  concentration (see Fig. 4) was not accompanied by a transient membrane depolarization expected if  $\text{Cl}^-$  was leaving the cell via conductance pathways. Such a transient depolarization was, for example, seen in skeletal muscle fibres after reduction of the  $\text{Cl}^-$  concentration in the extracellular solution (Hodgkin & Horowitz, 1959). However, two possibilities exist to explain this apparent contradiction. First,  $\text{Cl}^-$  could possibly leave these small cells passively as fast as the extracellular  $\text{Cl}^-$  concentration decreases. During such a situation  $E_{\text{Cl}}$  would never deviate from  $E_m$  and no potential change would be visible. To test this, a very rapid extracellular  $\text{Cl}^-$  concentration change would be necessary, an experiment not possible in our slice preparations. Secondly, there are reasons to assume that the anion channels of glial cells are unusually large in size and not completely impermeable to isethionate and gluconate, the  $\text{Cl}^-$  substituents used in the present study (Sonnhof & Schachner, 1984; Gray & Ritchie, 1985). In such a situation,  $\text{Cl}^-$  leaving the cell via conductance pathways would be replaced by the other anions entering the cell passively. Again, no potential change would be visible during the transition of high to low extracellular  $\text{Cl}^-$  concentration.

It should be noted, however, that a high  $\text{Cl}^-$  conductance does not preclude the presence of an active transport of  $\text{Cl}^-$ . Rather the effects of such a system on  $a_{\text{Cl}}^i$  will be largely short-circuited through the membrane because of its high  $\text{Cl}^-$  permeability (Vaughan-Jones, 1982). Indeed, previous studies, using radiotracer methods, have revealed inwardly directed  $\text{Cl}^-$  pump mechanisms in cultures from either astroglial

cells or from a glioma cell line (Kimelberg, 1981; Walz & Hertz, 1984; Wolpaw & Martin, 1984; see also Ballanyi & Grafe, 1985). It might be possible that such  $\text{Cl}^-$  pumps are involved in the pH regulation by the coupling to the transport of  $\text{HCO}_3^-$ . However, it is unlikely that active  $\text{Cl}^-$  transport contributes significantly to the stimulus-related rise of intragial  $a_{\text{Cl}}^i$ . This observation is better explained by the high  $\text{Cl}^-$  conductance of the membrane (see below).

#### *Stimulus-related changes of intracellular ion activities*

Our experiments demonstrate that the enhanced neuronal activity induced by the stimulation of the l.o.t. resulted in an increase in intragial  $a_{\text{K}}^i$  and  $a_{\text{Cl}}^i$ , whereas  $a_{\text{Na}}^i$  was decreased. The increase of  $a_{\text{K}}^i$  is similar to data from other glial cells (for references see Introduction), however, less is known about the behaviour of  $a_{\text{Cl}}^i$  and  $a_{\text{Na}}^i$ . Radiotracer data from mouse astrocytes in culture show that  $[\text{Cl}^-]_i$  increased and  $[\text{Na}^+]_i$  decreased when the extracellular  $\text{K}^+$  concentration was raised from 3 to 12 mM (Walz & Hertz, 1983b). Similar data was described for the bee retina during light stimulation (Coles & Orkand, 1984, 1985). Qualitatively, our data resemble that obtained from these studies.

What are the mechanisms underlying the stimulus-related changes of intracellular ion activities? With respect to the  $\text{K}^+$  uptake, at least two uptake processes have to be considered, which will be discussed in detail below. Concerning the increase in  $a_{\text{Cl}}^i$ , we first explored the possibility of a 'pump'-mediated uptake, since the described exclusive  $\text{K}^+$  conductance of the glial membrane (Kettenmann *et al.* 1983; Walz, Wuttke & Hertz, 1984b) apparently seems to exclude a passive increase. It should be noted, however, that in neuropile glial cells of the leech, a  $\text{Cl}^-$  permeability contributes significantly to the input conductance of the cell membrane (Walz & Schlue, 1982). Our experiments designed to reveal a  $\text{K}^+/\text{Cl}^-$  co-transport as one likely candidate failed to give a positive result. In particular, in the presence of  $\text{Ba}^{2+}$ , the usual  $a_{\text{Cl}}^i$  increase reversed to a small  $a_{\text{Cl}}^i$  decrease. Since  $\text{Ba}^{2+}$  did not inhibit the stimulus-related rise of  $a_{\text{K}}^e$ , which should be the important factor in activating a  $\text{K}^+/\text{Cl}^-$  co-transport, the latter result is not consistent with such a carrier-mediated increase of  $a_{\text{Cl}}^i$ . The experiments using  $\text{Ba}^{2+}$  did, on the other hand, reveal a strong correspondence between changes of  $E_m$  and  $a_{\text{Cl}}^i$ , which is good evidence for a high  $\text{Cl}^-$  conductance of the membrane (see above). Consequently, we favour the idea that stimulus-related changes of  $a_{\text{Cl}}^i$  are due to  $\text{Cl}^-$  movements via a conductive pathway.

Several possible mechanisms may explain the stimulus-related decrease of  $a_{\text{Na}}^i$ . These include activation of a  $\text{Na}^+/\text{K}^+$  pump, a decrease of an inwardly directed driving force for  $\text{Na}^+$  due to the membrane depolarization, a redistribution of  $\text{Na}^+$  within the glial syncytium as a result of spatial buffer currents, and/or cell swelling during  $\text{KCl}$  uptake. At the moment, we are not able to estimate the relative contribution of these possibilities. One important factor, however, seems to be the activation of a  $\text{Na}^+/\text{K}^+$  pump (see below).

#### *$\text{K}^+$ uptake via a $\text{Ba}^{2+}$ -sensitive mechanism*

$\text{Ba}^{2+}$  is known to block  $\text{K}^+$  channels in a variety of excitable cells (for review see Hille, 1984) and, as recently described, also in oligodendroglia (Corey, Barres & Chun, 1985). This effect has been previously used as a tool to differentiate between passive

and active  $K^+$  uptake mechanisms (Sjodin & Ortiz, 1975). The membrane depolarization, the partial block of  $K^+$  uptake, and the  $K^+$ -induced membrane hyperpolarization, observed in the presence of  $Ba^{2+}$ , are understandable in view of the present knowledge regarding the effects of this ion. The membrane depolarization can be explained by the closure of  $K^+$  channels which shifts the membrane potential towards the equilibrium potential of another ion species.  $Na^+$  is the most likely candidate, since  $E_{Cl}$  and  $E_m$  are almost identical.

The decrease in the  $K^+$ -induced rise of  $a_K^i$  during exposure to  $Ba^{2+}$  reveals an important passive  $K^+$  uptake into glial cells. One possibility as to how this uptake can take place is via a mechanism suggested by Boyle & Conway (1941) and Hodgkin & Horowicz (1959) to explain a passive uptake of  $K^+$  into muscle fibres. According to these authors, during an elevation of  $a_K^e$ , a transient deviation of  $E_m$  to the more depolarized  $E_K$  (due to the  $Cl^-$  conductance of the membrane) would result in an influx of  $K^+$  and  $Cl^-$ . Our data are compatible with this view. Another mechanism by which  $K^+$  uptake could occur through  $Ba^{2+}$ -sensitive channels would be via spatial buffer currents (Orkand *et al.* 1966; Dietzel *et al.* 1982; Coles & Orkand, 1983; Gardner-Medwin, 1983). According to this view,  $K^+$  would enter the glial syncytium through  $K^+$  channels as a result of a spatial potential gradient along the membrane of the electrically coupled glia; the rise of intragial  $a_{Cl}^i$  would be explained by a redistribution of  $Cl^-$  within the linked glial cells. Our experiments do not exclude such a mechanism. However, the existence of a passive KCl uptake would allow the  $K^+$  to be stored in the glial cell in the immediate neighbourhood of the active neurone, where it might be the more readily available to the neurone during recovery from  $K^+$  loss (Gray & Ritchie, 1985).

The third phenomenon observable in the presence of  $Ba^{2+}$  is a  $K^+$ -induced membrane hyperpolarization. This may be due to the activation of a  $K^+$ -induced electrogenic pump current (see below).

#### *$K^+$ uptake via the $Na^+/K^+$ pump*

Our data show that a small part of the stimulus-induced rise in  $a_K^i$  in the glial cells was  $Ba^{2+}$  insensitive. This component disappeared in the presence of ouabain. This finding is consistent with previous reports of a ouabain-sensitive,  $Na^+/K^+$  pump-mediated  $K^+$  uptake in glial cells from invertebrates (Tang, Cohen & Orkand, 1980; Walz, Wuttke & Schlue, 1983) and mammalian glia in culture (Kukes *et al.* 1976; Walz & Hertz, 1982). However, in contrast to other electrophysiological investigations (Tang *et al.* 1983; Walz *et al.* 1983), our results reveal that the electrogenic nature of this glial  $Na^+/K^+$  pump is a result of excess extracellular  $K^+$ ; raised intracellular  $Na^+$  is not essential. Since the membrane hyperpolarization is only seen in the presence of  $Ba^{2+}$ , it is likely that the high membrane conductance of the glia in the normal extracellular fluid shunts the pump-mediated outward current.

We would like to thank Drs H. Pusch and Th. Schumacher for their help in preparing a  $Na^+$ -sensitive cocktail and Dr K. Feasey for reading the manuscript. Ms G. Schneider, Mrs C. Müller and Ms I. Englmaier provided expert technical and secretarial assistance. The work was supported by the Deutsche Forschungsgemeinschaft (SFB 220).

## REFERENCES

- BALLANYI, K. & GRAFE, P. (1985). An intracellular analysis of  $\gamma$ -aminobutyric-acid-associated ion movements in rat sympathetic neurones. *Journal of Physiology* **365**, 41–58.
- BALLANYI, K., GRAFE, P., REDDY, M. M. & TEN BRUGGENCATE, G. (1984). Different types of potassium transport linked to carbachol and GABA actions in rat sympathetic neurons. *Neuroscience* **12**, 917–927.
- BALLANYI, K., GRAFE, P. & TEN BRUGGENCATE, C. (1985). Ion activities in glial cells of guinea pig olfactory cortex *in vitro*. *Pflügers Archiv* **405**, R 28.
- BOYLE, P. J. & CONWAY, E. J. (1941). Potassium accumulation in muscle and associated changes. *Journal of Physiology* **100**, 1–63.
- BÜHRLE, C. P. & SONNHOF, U. (1983). Intracellular ion activities and equilibrium potentials in motoneurons and glia cells of the frog spinal cord. *Pflügers Archiv* **396**, 144–153.
- COLES, J. A. & ORKAND, R. K. (1983). Modification of potassium movement through the retina of the drone (*Apis mellifera*) by glial uptake. *Journal of Physiology* **340**, 157–174.
- COLES, J. A. & ORKAND, R. K. (1984). Chloride enters receptors and glia in response to light or raised extracellular potassium in the retina of the bee drone, *Apis mellifera*. *Journal of Physiology* **357**, 11P.
- COLES, J. A. & ORKAND, R. K. (1985). Changes in sodium activity during light stimulation in photoreceptors, glia and extracellular space in drone retina. *Journal of Physiology* **362**, 415–435.
- COLES, J. A. & TSACOPOULOS, M. (1979). Potassium activity in photoreceptors, glial cells and extracellular space in the drone retina: changes during photostimulation. *Journal of Physiology* **290**, 525–549.
- COREY, D. P., BARRES, B. A. & CHUN, L. L. Y. (1985). An inwardly rectifying potassium channel in oligodendroglia. *Neuroscience Abstracts* **11**, 148.
- DIETZEL, I., HEINEMANN, U., HOFMEIER, G. & LUX, H. D. (1980). Transient changes in the size of the extracellular space in the sensorimotor cortex of cats in relation to stimulus induced changes in potassium concentration. *Experimental Brain Research* **40**, 432–439.
- DIETZEL, I., HEINEMANN, U., HOFMEIER, G. & LUX, H. D. (1982). Stimulus-induced changes in extracellular  $\text{Na}^+$  and  $\text{Cl}^-$  concentration in relation to changes in the size of the extracellular space. *Experimental Brain Research* **46**, 73–84.
- GARDNER-MEDWIN, A. R. (1983). Analysis of potassium dynamics in mammalian brain tissue. *Journal of Physiology* **335**, 393–426.
- GRAFE, P. (1985). Cellular mechanisms of potassium homeostasis in the vertebrate nervous system. *Pflügers Archiv* **405**, R 9.
- GRAFE, P., BALLANYI, K. & TEN BRUGGENCATE, G. (1985). Changes of intracellular free ion concentrations, evoked by carbachol or GABA, in sympathetic neurons. In *Ion Measurements in Physiology and Medicine*, ed. KESSLER, M., HARRISON, D. K. & HÖPER, J., pp. 184–188. Berlin, Heidelberg, New York: Springer.
- GRAFE, P., REDDY, M. M., EMMERT, H. & TEN BRUGGENCATE, G. (1983). Effects of lithium on electrical activity and potassium ion distribution in the vertebrate central nervous system. *Brain Research* **279**, 65–76.
- GRAFE, P., RIMPEL, J., REDDY, M. M. & TEN BRUGGENCATE, G. (1982). Changes of intracellular sodium and potassium ion concentrations in frog spinal motoneurons induced by repetitive synaptic stimulation. *Neuroscience* **7**, 3213–3220.
- GRAY, P. T. A. & RITCHIE, J. M. (1985). Ion channels in Schwann and glial cells. *Trends in Neurosciences* **8**, 411–415.
- GRISAR, T., FRANCK, G. & DELGADO-ESCUETA, A. V. (1983). Glial contribution to seizure:  $\text{K}^+$  activation of ( $\text{Na}^+$ ,  $\text{K}^+$ )-ATPase in bulk isolated glial cells and synaptosomes of epileptogenic cortex. *Brain Research* **261**, 75–84.
- HILLE, B. (1984). *Ionic Channels of Excitable Membranes*. Sunderland, Massachusetts: Sinauer Associates Inc.
- HODGKIN, A. L. & HOROWICZ, P. (1959). The influence of potassium and chloride ions on the membrane potential of single muscle fibres. *Journal of Physiology* **148**, 127–160.
- KETTENMANN, H., SONNHOF, U. & SCHACHNER, M. (1983). Exclusive potassium dependence of the membrane potential in cultured mouse oligodendrocytes. *Journal of Neuroscience* **3**, 500–505.

- KIMELBERG, H. K. (1981). Active accumulation and exchange transport of chloride in astroglial cells in culture. *Biochimica et biophysica acta* **646**, 179–184.
- KUKES, G., ELUL, R. & DE VELLIS, J. (1976). The ionic basis of the membrane potential in a rat glial cell line. *Brain Research* **104**, 71–92.
- MCCAIG, D. & LEADER, J. P. (1984). Intracellular chloride activity in the extensor digitorum longus (EDL) muscle of the rat. *Journal of Membrane Biology* **81**, 9–17.
- MEIER, P. C., AMMANN, D., MORF, W. E. & SIMON, W. (1980). Liquid-membrane ion-sensitive electrodes and their biomedical applications. In *Medical and Biological Applications of Electrochemical Devices*, ed. KORYTA, J., pp. 13–91. New York: J. Wiley & Sons Ltd.
- MEIER, P. C., LANTER, F., AMMANN, D., STEINER, R. A. & SIMON, W. (1982). Applicability of available ion-selective liquid-membrane microelectrodes to intracellular ion-activity measurements. *Pflügers Archiv* **339**, 23–30.
- MORRIS, M. E., KRNEVIĆ, K. & MACDONALD, J. F. (1985). Changes in intracellular free Ca ion concentration evoked by electrical activity in cat spinal neurons *in situ*. *Neuroscience* **14**, 563–580.
- NICHOLSON, C. (1981). Brain-cell microenvironment as a communication channel. In *The Neurosciences – Fourth Study Program*, ed. SCHMITT, F. O. & WORDEN, F. G., pp. 457–476. Cambridge, MA: MIT Press.
- NICHOLSON, C., TEN BRUGGENCATE, G. & SENEKOWITSCH, R. (1976). Large potassium signals and slow potentials evoked during aminopyridine or barium superfusion in cat cerebellum. *Brain Research* **113**, 606–610.
- ORKAND, R. K., NICHOLLS, J. G. & KUFFLER, S. W. (1966). Effect of nerve impulses on the membrane potential of glial cells in the central nervous system of Amphibia. *Journal of Neurophysiology* **29**, 788–806.
- SCHLUE, W. R. & WUTTKE, W. (1983). Potassium activity in leech neuropile glial cells changes with external potassium concentration. *Brain Research* **270**, 368–372.
- SJODIN, R. A. & ORTIZ, O. (1975). Resolution of the potassium ion pump in muscle fibers using barium ions. *Journal of General Physiology* **66**, 269–286.
- SOMJEN, G. G. (1979). Extracellular potassium in the mammalian central nervous system. *Annual Reviews of Physiology* **41**, 159–177.
- SONNHOF, U. & SCHACHNER, M. (1984). A high conductance anion-cation channel in cultured astrocytes. *Pflügers Archiv* **402**, R31.
- SYKOVA, E. (1983). Extracellular K<sup>+</sup> accumulation in the central nervous system. *Progress in Biophysics and Molecular Biology* **42**, 135–190.
- TANG, C. M., COHEN, M. W. & ORKAND, R. K. (1980). Electrogenic pumps in axons and neuroglia and extracellular potassium homeostasis. *Brain Research* **194**, 283–286.
- VAUGHAN-JONES, R. D. (1982). Chloride activity and its control in skeletal and cardiac muscle. *Philosophical Transactions of the Royal Society of London B* **299**, 537–548.
- WALZ, W. & HERTZ, L. (1982). Ouabain-sensitive and ouabain-resistant net uptake of potassium into astrocytes and neurons in primary cultures. *Journal of Neurochemistry* **39**, 70–77.
- WALZ, W. & HERTZ, L. (1983a). Functional interactions between neurones and astrocytes. II. Potassium homeostasis at the cellular level. *Progress in Neurobiology* **20**, 133–183.
- WALZ, W. & HERTZ, L. (1983b). Intracellular ion changes of astrocytes in response to extracellular potassium. *Journal of Neuroscience Research* **10**, 411–423.
- WALZ, W. & HERTZ, L. (1984). Intense furosemide-sensitive potassium accumulation in astrocytes in the presence of pathologically high extracellular potassium levels. *Journal of Cerebral Blood Flow and Metabolism* **4**, 301–304.
- WALZ, W. & SCHLUE, W. R. (1982). External ions and membrane potential of leech neuropile glial cells. *Brain Research* **239**, 119–138.
- WALZ, W., SHARGOOL, M. & HERTZ, L. (1984a). Barium-induced inhibition of K<sup>+</sup> transport mechanisms in cortical astrocytes – its possible contribution to the large Ba<sup>2+</sup> evoked extracellular K<sup>+</sup>-signal in brain. *Neuroscience* **13**, 945–949.
- WALZ, W., WUTTKE, W. & HERTZ, L. (1984b). Astrocytes in primary cultures: membrane potential characteristics reveal exclusive potassium conductance and potassium accumulator properties. *Brain Research* **292**, 367–374.
- WALZ, W., WUTTKE, W. & SCHLUE, W. R. (1983). The Na<sup>+</sup>-K<sup>+</sup> pump in neuropile glial cells of the medicinal leech. *Brain Research* **267**, 93–100.
- WOLPAW, E. W. & MARTIN, D. L. (1984). Cl<sup>-</sup>-transport in a glioma cell line: evidence for two transport mechanisms. *Brain Research* **297**, 317–327.