Environmental Microbiology (2019) 21(11), 4125-4135



# They are young, and they are many: dating freshwater lineages in unicellular dinophytes

# Anže Žerdoner Čalasan <sup>(b)</sup>,<sup>1,2</sup> Juliane Kretschmann<sup>2</sup> and Marc Gottschling <sup>(b)2\*</sup>

<sup>1</sup>Department Biologie/Chemie, Botanik, Universität Osnabrück, Barbarastr. 11, 49076 Osnabrück, Germany. <sup>2</sup>Department Biologie, Systematische Botanik und Mykologie, GeoBio-Center, Ludwig-Maximilians-Universität München, Menzinger Street. 67, 80638 Munich, Germany.

### Summary

Dinophytes are one of few protist groups that have an extensive fossil record and are therefore appropriate for time estimations. However, insufficient sequence data and strong rate heterogeneity have been hindering to put dinophyte evolution into a time frame until now. Marine-to-freshwater transitions within this group are considered geologically old and evolutionarily exceptional due to strong physiological constraints that prevent such processes. Phylogenies based on concatenated rRNA sequences (including 19 new GenBank entries) of two major dinophyte lineages, Gymnodiniaceae and Peridiniales, were carried out using an uncorrelated molecular clock and five calibration points based on fossils. Contrarily to previous assumptions, marine-to-freshwater transitions are more frequent in dinophytes (i.e. five marine-freshwater transitions in Gymnodiniaceae, up to ten but seven strongly supported transitions in Peridiniales), and none of them occurred as early as 140 MYA. Furthermore, most marine-to-freshwater transitions, and the followed diversification, took place after the Cretaceous-Paleogene boundary. Not older than 40 MYA, the youngest transitions within Gymnodiniaceae and Peridiniales occurred under the influence of the Eocene climate shift. Our evolutionary scenario indicates a gradual diversification of dinophytes without noticeable impact of catastrophic events, and their freshwater lineages have

Received 1 March, 2019; revised 24 July, 2019; accepted 29 July, 2019. \*For correspondence. Tel. (+49) 89 17861 271; Fax (+49) 89 172 638. E-mail gottschling@bio.lmu.de

originated several times independently at different points in time.

#### Introduction

The majority of the world's surface is covered with water (Simon et al., 2015), but the difference between marine and fresh water bodies in terms of volume is immense. Oceans comprise around 71% of the globe's surface (Costanza, 1999), and not more than 3% of it is covered in freshwater (Gleick, 1999; Downing et al., 2006). Marine and freshwater ecosystems differ greatly in their physicochemical characteristics, subsequently dictating physiological adaptations developed in the inhabiting organisms. Only a handful of organisms, such as migratory fish (Myers, 1949), have considerable salinity tolerance ranges, whereas most species show a remarkably high fidelity to their respective habitats. On a more general level, a number of eukarvotic lineages often group into distinct and species-rich marine or freshwater phylogenetic clusters, a phenomenon known not only from macroorganisms but also from the microbiome (Shalchian-Tabrizi et al., 2008; Logares et al., 2009, 2010; Heger et al., 2010).

Some of the important questions in evolutionary biology are the origin, number and age of evolutionary transitions between freshwater and marine environments. Aquatic, especially marine environments lack apparent obstacles for pelagic organisms (Paulay and Meyer, 2002; Bråte et al., 2010), thus reducing the impact of allopatric diversification. However, the fundamental osmotic difference between the oceans and diverse freshwater habitats acts as a strong physiological barrier (Pokorný, 2009). Only a few organisms are physiologically adapted to successfully overcome such a barrier and establish new populations. Subsequently, such marine-to-freshwater (or vice versa) transitions are considered to be exceptional and uncommon (100 transitions in 800 MYA for all the eukaryotes: Cavalier-Smith, 2009). Regardless, microorganisms form much larger populations, have higher reproductive rates (Dolan, 2005; Snoke et al., 2006), and are significantly smaller than their macroscopic counterparts. Therefore, it is assumed that barrier-crossing between marine and

# 4126 A. Žerdoner Čalasan, J. Kretschmann and M. Gottschling

freshwater environments takes place more often in the microbiome (Finlay *et al.*, 2006; Bass *et al.*, 2007). More specifically, several (albeit infrequent) transitions between marine and freshwater environments are inferred for microbial groups such as chlorophytes, cryptophytes, dinophytes and haptophytes (Lewis and McCourt, 2004; Logares *et al.*, 2007b; Shalchian-Tabrizi *et al.*, 2008, 2011; Leliaert *et al.*, 2012), indicating that some groups are indeed capable of crossing such a strong physiological barrier.

Dominating the freshwater and marine aquatic ecosystems, microbial eukaryotes play an important role in carbon fixation (Jardillier *et al.*, 2010) and climate regulation (Simó, 2001). Dinophytes are one of major phytoplankton groups and inhabit a continuum of water environments from open seas, shallow coastal waters, estuaries, rivers and lakes to puddles (Fensome *et al.*, 1993; Taylor *et al.*, 2008). The majority of dinophyte species live in marine environments, but approximately 350 species can be found in freshwater habitats (Mertens *et al.*, 2012; Moestrup and Calado, 2018; Fig. 1). Quantification and dating of marine-to-freshwater transitions is the basis toward understanding the past evolutionary processes leading to the high biodiversity in dinophytes. Such studies are rare because of the limited taxon sample availability (compared with the total known diversity), insufficient sequence data and strong rate heterogeneity (particularly of rRNA sequences; Saldarriaga et al., 2004; Murray et al., 2005: Gu et al., 2013). Nevertheless, preliminary data are available for some subordinate groups. Comprised of approximately 50 extant species (Elbrächter et al., 2008), the Thoracosphaeraceae (Peridiniales) can be found in either freshwater or marine habitats. The group clearly prefers the latter, but at least three independent freshwater lineages are recognized within this group (Moestrup and Daugbjerg, 2007; Logares et al., 2007a; Craveiro et al., 2013; Gottschling and Söhner, 2013). Similar dynamics is inferred from the Gymnodiniaceae, as at least five lineages have independently colonized freshwater habitats (Kretschmann et al., 2015).

To deepen our knowledge of marine-to-freshwater transitions of all kinds of organisms, the time component placing transition events into a certain geological era plays an important role. Evidence based on the fossil record (Gray and Taylor, 1988; Batten, 1989) suggests that marine-tofreshwater transitions within dinophytes started to happen



Fig. 1. Diversity of freshwater dinophytes (all cells on the same scale).

- A. Chimonodinium lomnickii var. wierzejskii (Thoracosphaeraceae).
- B. Durinskia oculata (Kryptoperidiniaceae).
- C. Parvodinium umbonatum (Peridiniopsidaceae).
- D. Palatinus apiculatus (Peridiniopsidaceae).
- E. Peridinium cinctum (Peridiniaceae).
- F. Peridinium bipes (Peridiniaceae).
- G. Peridinium willei (Peridiniaceae).
- H. Spiniferodinium limneticum (Gymnodiniaceae). [Color figure can be viewed at wileyonlinelibrary.com]

as early as 140 MYA. Molecular clock analyses shed light on certain previously unexplainable evolutionary events (Medlin, 2008). Based on the integrative fossil record and a relaxed molecular clock approach dinophytes diverged from their sister group Apicomplexa around 1100 MYA (Parfrev et al., 2011; Feulner et al., 2015), but did not diversify until the beginning of the Mesozoic (i.e., after the 'phytoplankton blackout': Riegel. 2008: Servais et al., 2016: and when Pangea started to break apart ~250 MYA: Fensome et al., 1996, 1999). The downfall resulted in an increase of continental shelf areas providing many new habitats for coastal and benthic dinophytes (Fensome et al., 1996; Hoppenrath et al., 2014). Time estimates focusing exclusively on dinophytes are only sporadically available and focus on rather lower taxonomic levels (Alexandrium: John et al., 2003: Symbiodinium: LaJeunesse, 2004: LaJeunesse et al., 2018; Pochon et al., 2006; calcareous dinophytes: Gottschling et al., 2008; Kryptoperidiniaceae: Žerdoner Čalasan et al., 2018) yet to understand the marine-tofreshwater transitions to a greater extent, putting them into a time frame is of the essence.

A considerably higher coverage of genetic information and well-resolved phylogenetic relations between dinophyte taxa on a higher level (Gottschling et al., 2012; Orr et al., 2012; Gu et al., 2013; Kretschmann et al., 2015, 2018; Janouškovec et al., 2017; Price and Bhattacharya, 2017) are the first steps toward a better understanding of complex processes such as marine-to-freshwater transitions. Together with their rich high-quality fossil record (Fensome et al., 1993, 1996), we can use an integrative approach to assign these transitions to geological events (Logares et al., 2009). The aim of our study was to provide the number and the time frame of marine-tofreshwater transitions by carrying out a molecular clock analysis for the Gymnodiniaceae and the Peridiniales (Fig. 1) - two groups with the highest number of marineto-freshwater transitions known in dinophytes.

#### Results

The alignment was comprised of 2 050 parsimonyinformative positions (32%, mean of 9.2 per terminal taxon) and included small ribosomal subunit (SSU), internal transcribed spacer region (ITS1, 5.8S and ITS2) and large ribosomal subunit (LSU). Our data set provided a well-resolved backbone phylogeny using a representative taxon sample of marine and freshwater representatives of the Peridiniales (Fig. 2) and the Gymnodiniaceae (Fig. 3; both groups were considered as the out groups for each other). Maximum Likelihood (likelihood bootstrap support; LBS) and Bayesian analyses (Bayesian posterior probabilities; BPP) recovered phylogenetic trees with highly congruent topologies. The Peridiniales (100 LBS,

#### Marine/freshwater transitions in protists 4127

1.00 BPP) segregated into eight highly diverse monophyletic lineages, including *Scrippsiella sensu lato* (97 LBS, 1.00 BPP), a clade comprising *Thoracosphaera* and *Pfiesteria* with relatives (T/Pf clade; 89 LBS, 1.00 BPP), the Kryptoperidiniaceae (93 LBS), a clade comprising *Ensiculifera* and *Pentapharsodinium* with relatives (98 LBS, 0.99 BPP), the heterotrophic Protoperidiniaceae (94 LBS), the Peridiniaceae (100 LBS, 1.00 BPP), the Peridiniopsidaceae (92 LBS, 1.00 BPP) and the Heterocapsaceae (100 LBS, 1.00 BPP). The relationships among such lineages were not always clear, but some nodes were retrieved with high support, for example, the close relationship between *Scrippsiella s.I.* and the T/Pf clade (97 LBS, 1.00 BPP).

To assess the influence of highly variable regions, two separate alignments were used for the time estimation analysis. Due to a high variability of the ITS region, certain positions were namely aligned unreliably, which might cause artefacts in the topologies or time spans. Hence, to further check for the influence of potentially poorly aligned positions, we carried out the analyses once using the ITS region (Fig. 3, Supporting Information Fig. S2) and once excluding it from the alignment (Supporting Information Figs S1 and S3). No significant differences in topologies, time frames or in effective sample size (ESS) values were recovered, which underlines the solidity of our approach. Furthermore, two different tree priors (Speciation: Yule process; Figs 2 and 3, Supporting Information Fig. S1; and Coalescence: constant size; Supporting Information Figs S2 and S3) were used. For representative time estimation analysis, one should namely use Yule process, in cases of having one species per sample in the taxon set, or alternatively Coalescence Model, in cases of having more accessions per species included in the analysis (see BEAST tutorials 2012). However, we cannot be sure either of the criteria is fulfilled by our taxon sample due to the unresolved taxonomy hindering species delimitation in dinophytes. Hence, both tree priors were tested. Nevertheless, the final dated phylogeny in terms of time spans and topology again hardly differed (the greatest difference referred to the Peridiniaceae crown group that was dated to 64 and 70 MYA, respectively). The only topological change worth mentioning the paraphyly was Speciation: Yule process) or monophyly (in (in Coalescence: constant size) between Chimonodinium and Fusiperidinium from the Thoracosphaeraceae. Nevertheless, the statistical support of both scenarios was not well supported in either of the analyses (BPP < 0.90).

Based on highly supported sister group relationships between marine and freshwater clades, seven independent marine-to-freshwater transitions within the Peridiniales were recovered. Three additional lineages (i.e., the Peridiniaceae, an unnamed clade that included *Gloeodinium* with a relative,



**Fig. 2.** Dated phylogeny of Peridiniales. Dated phylogeny (ultrametric maximum clade credibility tree with node ages from the Bayesian uncorrelated lognormal analysis, under the Yule model) using concatenated rRNA sequences (i.e., SSU, ITS, LSU). Median rate is given in units of substitutions per million years (including 95% confidence intervals). Absolute ages are in million years, and epochs are indicated in the same colors as in Gradstein *et al.* (2012). The numbers on the branches are statistical support values (above: ML bootstrap values, values < 50 are not shown; below: Bayesian posterior probabilities, values < 0.90 are not shown; asterisks indicate maximal support). Note the presence of up to ten independent marine-to-freshwater transitions, six of which happened after the K-Pg boundary, depicted with an orange line. The calibration points are indicated with yellow stars. Freshwater taxa are colored in red. Abbreviations: E/Pe, clade including *Ensiculifera* and *Pentapharsodinium*; HET, Heterocapsaceae; POP, Peridiniopsidaceae. The used outgroup Gymnodiniaceae is shown separately in the Fig. 3. [Color figure can be viewed at wileyonlinelibrary.com]



Fig. 3. Dated phylogeny of Gymnodiniaceae. Dated phylogeny (ultrametric maximum clade credibility tree with node ages from the Bayesian uncorrelated lognormal analysis, under the Yule model) using concatenated rRNA sequences (i.e., SSU, ITS, LSU). Median rate is given in units of substitutions per million years (including 95% confidence intervals). Absolute ages are in million years, and epochs are indicated in the same colors as in Gradstein *et al.* (2012). The numbers on the branches are statistical support values (above: ML bootstrap values, values < 50 are not shown; below: Bayesian posterior probabilities, values < 0.90 are not shown; asterisks indicate maximal support). Note the presence of five independent marine-to-freshwater transitions, all of which happened after the K-Pg boundary, depicted with an orange line. Freshwater taxa are colored in red. The used outgroup Peridiniales as well as the calibration points indicated with yellow stars are shown separately in Fig. 2. [Color figure can be viewed at wileyonlinelibrary.com]

and the Peridiniopsidaceae) may have colonized the freshwater habitat independently as well, but statistical support for deep nodes was overall low. An alternative scenario would assume an initial marine-to-freshwater transition at the base of the Peridiniales, with at least one transition back into the marine environment and thus requiring at least two evolutionary steps. Four distantly related freshwater lineages - the Peridiniaceae, the Gloeodinium lineage, the Peridiniopsidaceae, and Unruhdinium of the Kryptoperidiniaceae - originated prior to the Cretaceous-Paleogene (K-Pg) boundary (though switches to the freshwater environment could not be precisely dated). However, diversification of such lineages mostly took place after the K-Pg boundary (Supporting Information Fig. S4), except for the Peridiniopsidaceae, for which it may have started already in the late Cretaceous. Four freshwater lin-eages originated after the K-Pg boundary and were usually nested deeply in marine clades (e.g., Durinskia from the Kryptoperidiniaceae, Naiadinium and Theleodinium from Scrippsiella s.l.) and for the remaining two clades (Chimonodinium and Fusiperidinium), it is unclear whether the transition took place after or before the K-Pg boundary. The T/Pf clade (except for the strictly marine *Leonella*, †*Posoniella* and *Thoracosphaera*) showed a general preference for habitats with reduced salinity. Within this clade, four lineages moved into the freshwater environment, namely *Apocalathium*, *Chimonodinium*, *Fusiperidinium* and *Tyrannodinium*. Stem ages, however, differed greatly.

Resolution within Gymnodiniaceae (100 LBS, 1.00 BPP) was poorer and only some groups at higher taxonomic levels could be identified, such as the *Nusuttodinium/Spiniferodinium*-clade (100 LBS, 1.00 BPP) and the warnowians (81 LBS, 0.95 BPP). Five transitions were still inferred, three alone in the *Nusuttodinium/Spiniferodinium*-clade (100 LBS, 1.00 BPP): Freshwater *Spiniferodinium* palustre constituted the sister species of marine "*Gymnodinium*" cf. placidum (100 LBS, 1.00 BPP), freshwater *Spiniferodinium* constituted the sister species of marine *Spiniferodinium palauense* (63 LBS) and freshwater and marine representatives of *Nusuttodinium* constituted sister species sister species of marine spiniferodinium palauense (53 LBS) and freshwater and marine representatives of *Nusuttodinium* constituted sister species sister species of marine spiniferodinium palauense (53 LBS) and freshwater and marine representatives of *Nusuttodinium* constituted sister species sister species of marine ter groups as well (79 LBS, 0.99 BPP). The other two

transitions were represented by the isolated species *Gymnodinium fuscum* and in an undetermined gymnodiniacean clade isolated from freshwater sponges. All freshwater transitions and consequent radiations within Gymnodiniaceae occurred long after the K-Pg boundary (Supporting Information Fig. S1). All five lin-eages originated and diversified between late Miocene and early Oligocene (between 30 and 15 MYA), except for the *Nusuttodinium aeruginosum* species group, which possibly transitioned in mid-Miocene already (around 40 MYA).

#### Discussion

There is a growing body of evidence that some protist lineages may have originated in freshwater (Logares et al., 2009; Mulkidjanian et al., 2012; Dittami et al., 2017; Sánchez-Baracaldo et al., 2017) and that evolutionary transitions followed the direction from freshwater into marine habitats (Cavalier-Smith, 2009). In such studies, it is also explained that large and species-rich (sister) groups with fundamentally different ecological preferences to either the marine or the freshwater environment developed parallelly in many eukaryotic lineages. Dinophytes appear different in such aspects: The last common ancestor most likely lived in a marine environment, and freshwater dinophyte groups are relatively species-poor and deeply nest in primarily marine groups (Moestrup and Daugbjerg, 2007; Gottschling and Söhner, 2013; Kretschmann et al., 2015, 2018). Before our study, marine-to-freshwater transitions have been considered infrequent in dinophytes (Logares et al., 2007b), although terms such as 'infrequent' or 'rare' are not objectifiable. Nevertheless, we account for approximately five lineages within Gymnodiniaceae and up to ten lineages within Peri-diniales, all of which independently and successfully colonized different freshwater habitats. More and more freshwater lineages with close relatives among marine representatives are known from other groups as well, such as Gonyaulacales, †Suessiales, and Tovelliaceae (Hansen et al., 2007: Fawcett and Parrow, 2012), and their precise phylogenetic positions will be determined once we improve the molecular trees of dinophytes.

The present time estimates are in agreement with previous analyses investigating Peridiniales (Gottschling *et al.*, 2008; Žerdoner Čalasan *et al.*, 2018). Merely, the Kryptoperidiniaceae have been dated slightly older before (Late: between 100 MYA and 66 MYA, versus Upper: between 145 and 100 MYA, Cretaceous stemgroup ages), but this can be easily explained by the inclusion of long-branching *Kryptoperidinium* that was omitted from the present analysis. The overall high number of freshwater lineages, even in subsets of dinophytes, is indicative of a more dynamic biotic exchange between marine and freshwater environments. In the aspect of transition polarity, dinophytes are rather similar to haptophytes (Shalchian-Tabrizi *et al.*, 2011), diatoms (Alverson *et al.*, 2007) and cercozoan amoebae (Berney *et al.*, 2013), which presumably also originated in marine environments and migrated into freshwater habitats to a certain extent. Although marine-to-freshwater transitions are physiologically more challenging than the reverse process (Cavalier-Smith, 2009), not a single doubtless case of such a recolonization of the marine environment could be inferred for dinophytes.

In addition to the higher number of different freshwater lineages (albeit still lower in comparison to the marine lineages), we also present new insights about the geological age of the processes. Since the Cretaceous, marineto-freshwater transitions occur continuously in a more gradual process and without noticeable impacts of catastrophic events. Most freshwater stem nodes are dated to after the K/Pg-boundary (< 65 MYA), and diversification within such lineages has mostly taken place during the last 10 million years. However, prior studies have posited that most marine-to-freshwater transitions in dinophytes took place prior to the K/Pg-boundary (Cavalier-Smith, 2009; Logares et al., 2009). This conclusion seems to be corroborated with tree topology and with the presence of relatively long branches separating several freshwater lineages from marine groups (Logares et al., 2007b), but it is not based on a time-calibrated phylogenetic analysis as presented here. The fossil record placing freshwater dinophytes into the Mesozoic (Gray and Taylor, 1988; Batten, 1989) also seems to support the idea of freshwater dinophyte lineages being consistently old, but the precise systematic affiliation of such fossils has not yet been determined. We cannot exclude that such Mesozoic fossils in fact correspond to (also extant) freshwater dinophyte lineages, but the data pre-sented here rather suggest that they are representatives of a stem lineage or even of an extinct lineage. The fossil record of freshwater dinophytes is scarce (Fensome et al., 1993), which means that it should be considered with caution. An incomplete or lost fossil record combined with often overlooked high rates of heterogeneity (not necessarily indicating high ages) could be the reason for the assumption of predominantly Mesozoic marine-to-freshwater transitions to have persisted until now.

Our analysis challenges previous evolutionary scenarios, as it shows that marine-to-freshwater transitions are not infrequent and are geologically rather young. Mostly not older than 40 MYA, the origin and diversification of gymnodiniacean and peridinialean freshwater dinophytes correlate with the Eocene climate dynamic changes. Approximately 37 MYA, the Earth started to cool down,

and generally warm and humid environments were re-placed by continental ice sheets and/or glaciers, which stretched as far as to the sea level (Barron and Baldauf. 1989; Miller et al., 1991; Lear et al., 2008). A general decline in the sea level caused by cooling resulted in the isolation of lagoons, which became isolated from the rest of the open sea (Matthews, 2013). If those lagoons were supplied with fresh water from the land, then the concentration of chloride and sodium ions would decline sharply, resulting in many new and initially uncolonized freshwater habitats. In such cases, the organisms trapped in such lagoons had to either adapt to new conditions or become extinct. The cooling of the Earth was already connected with the diversification processes in Symbiodinium, an endosymbiotic dinophyte group found in marine invertebrates and protists (Pochon et al., 2006).

The origin and diversification of the few freshwater lin-eages older than the Eocene may also correlate with significant geological events within the Earth's history. Considering the age of marine-to-freshwater transitions, only three distantly related groups (namely the Peridiniaceae, the Peridiniopsidaceae, and the Gloeodinium lineage) could be older than the K-Pg boundary and of Cretaceous origin. For this era, major environmental perturbations and significant eustatic sea-level falls are assumed, which had immense effects on shallow marine shelf environments (Sims et al., 2006; Tennant et al., 2017). It appears that dinophytes were not greatly affected by catastrophic events at, for example, the K-Pg boundary, as numerous lineages passed through this massive extinction event (Gottschling et al., 2008; Figs 2 and 3, Supporting Information Figs S1-S3). Because of large confidence intervals in this part of the dating analysis (spanning on average 75 MYA), such correlations should be considered with caution. However, most freshwater dinophyte diversification appears to have taken place during the Neogene (< 25 MYA; Supporting Information Fig. S4). The impact of the catastrophic flood of the Mediterranean after the Messinian salinity crisis (Bison et al., 2007; Popescu et al., 2009), or repeated Pleistocene glaciations (Pelzer, 1991; Lindner et al., 2004), remains to be determined. For robust conclusions, a (nearly) complete taxon sample is required, but we are still missing parts in most dinophyte lineages.

The question arises, why would dinophytes cross the marine/freshwater barrier more frequently than other protist groups such as chlorophytes and rhodophytes. A general explanation is that this ecologically and physiologically diverse group copes much better with extreme temperatures and salinity ranges (Taylor *et al.*, 2008). As living models, species such as *Alexandrium ostenfeldii* and *Huia caspica* might be a key in this respect, as they are found in both marine and freshwater habitats (Kremp *et al.*, 2014; Gu *et al.*, 2016; Martens *et al.*, 2016). Something similar can be said for pfiesterian dinophytes. a group with a reduced salinity preference (Litaker et al., 2005: Calado et al., 2009: Burkholder and Marshall, 2012). Another example is the phenotypically differentiated species pair, freshwater Apocalathium aciculiferum and brackish Apocalathium malmogiense, sharing (almost) identical rRNA sequences (Gottschling et al., 2005: Logares et al., 2007a). Thus, it is argued that the marine-to-freshwater transition in this species pair is ongoing (Logares et al., 2008; Annenkova et al., 2015). Finally, many dinophytes produce benthic resting stages (Fensome et al., 1993; Mertens et al., 2012; Gu et al., 2016), which enable them to withstand long periods of unfavourable conditions. This particular ability could be of great evolutionary advantage if not a necessary prerequisite to surviving intermediate stages during such marineto-freshwater transitions.

# **Experimental procedures**

The inference of dinophyte evolution has been improved using next-generation sequencing (NGS; Price and Bhattacharya, 2017; Janouškovec et al., 2017), but a wide range of information of the taxon sample continues to be recruited from the rRNA operon sequences only. Due to restrictions such as limited taxon sample, insufficient genetic data, complex phylogenetic methods and rate heterogeneity, a representative taxon sample covering the currently known dinophyte diversity is not present at this moment in time (Gottschling et al., 2012; Gu et al., 2013). However, the phylogeny of the Peridiniales (Gottschling et al., 2017) and the Gymnodiniaceae (Kretschmann et al., 2015), including many freshwater presenters, is relatively well-understood, which is why these two lineages were selected for the time estimation analysis.

For establishment of own monoclonal strains (Supporting Information Table S1), single motile cells were isolated and placed in 24-well microplates (Zefa; Munich, Germany) containing freshwater WC growth medium (Woods Hole Combo, modified after Guillard and Lorenzen, 1972) without silicate. The plates were stored in climate chambers at 12°C or 18°C and a 12:12 h L:D photoperiod. The established monoclonal strains are currently held in the culture collection at the Institute of Systematic Botany and Mycology (University of Munich) and are available upon request. Nineteen new nuclear rRNA sequences of dinophyte were obtained as described previously (Gottschling *et al.*, 2012; Kretschmann *et al.*, 2018).

For the phylogenetic analysis, separate loci-based SSU, ITS, and LSU matrices were constructed, aligned using 'MAFFT' version 6.502a (Katoh and Standley, 2013), and concatenated with SeaView version 4.6.1 (Gouy *et al.*, 2010). We compiled a taxon sample that was

# 4132 A. Žerdoner Čalasan, J. Kretschmann and M. Gottschling

representative with respect to the known molecular sequence diversity of the Gymnodiniaceae and the Peri-diniales (functioning as mutual outgroups) and included at least one (or more if available) representatives for every freshwater lineage (see the taxon sample used in this study: Supporting Information Table S1). Though they are of paleontological importance (Fensome et al., 1993, 1996), the exclusively marine Protoperidiniaceae were partially excluded because of high substitution rates resulting in long branches (such as Protoperidinium abei and Protoperidinium denticulatum: Gu et al., 2013; as well as long-branching Kryptoperidinium from the Kryptoperidiniaceae: Kretschmann et al., 2018). Not all Protoperidiniaceae are subjected to this phenomenon, and this lineage is nevertheless well represented in our taxon samples including only representatives exhibiting branches of similar length in comparison to the rest of taxa.

Phylogenetic analyses were carried out using both the Maximum Likelihood (ML) and the Bayesian inference approach (BI). Maximum Likelihood-based analyses were conducted using RAxML RAxML-HPC version 8.2.10 (Stamatakis, 2014) with applied GTR +  $\Gamma$  substitution matrix. To determine best-fitted ML-trees, we carried out 10-tree searches from distinct random stepwise addition sequence Maximum Parsimony starting trees and 10 000 non-parametric bootstrap replicates. Bayesian analyses were carried out using MrBayes version 3.2.6 (Ronquist et al., 2012) under the GTR +  $\Gamma$  substitution model using the random-addition-sequence method with 10 replicates. We ran two independent analyses of four chains (one cold and three heated) with 20 000 000 cycles, sampling every 1000th cycle, with an appropriate burn-in (10%) as inferred from the evaluation of the trace files using Tracer version 1.5 (http://tree.bio.ed.ac.uk/software/tracer/). The statistical support values were drawn on the presented chronograms. Marine-to-freshwater transitions were reliably inferred based on sister group relationships between marine and freshwater clades (or freshwater clades nesting within marine clades) with statistical support  $\geq 80$ LBS and/or ≥0.90 BPP respectively. However, pure topology (of, e.g., less well resolved deeper nodes) was taken into account as well. All phylogenetic inferences were carried out at the CIPRES Science Gateway computing facility (Miller et al., 2010). The aligned matrices are available as \*.nex files upon request.

To estimate divergence times in the Gymnodiniaceae and the Peridiniales, the aforementioned alignment was used and the phylogeny was dated using BEAST version 1.8.3 (Drummond *et al.*, 2012). We carried out several different analyses using different evolutionary models with corresponding parameters (e.g., Yule process versus Coalescence) and different alignments (e.g., including and excluding heterogeneous sites). The final decision on the analysis was made based on ESS values, which had to exceed 200. The most fitting evolutionary model (GTR +  $\Gamma$  + i) and the rate heterogeneity were assessed with jModelTest (Darriba *et al.*, 2012); the +i option was ignored, as it leads to overparameterization (see RAxML manual). Unlike some other generally scarce dinophyte dating studies (John *et al.*, 2003; LaJeunesse, 2004; Pochon *et al.*, 2006), we used in-group fossils (the advantages for this strategy are discussed in Gottschling *et al.*, 2008).

For the presented analysis, a Yule branching process with lognormal priors and empirically defined base frequencies was adopted using the complete alignment and following five calibration points (minimal ages for stem groups, see Gottschling et al., 2008): Peridiniales were estimated at 200  $\pm$  1 MYA (Fensome et al., 1996), Peridiniales excluding Heterocapsaceae were estimated at 160  $\pm$  4 MYA based on the first fossil occurrence of (calcareous) coccoid cells with tabulation (Keupp, 1984; Keupp and Ilg, 1989), Scrippsiella s.I. was estimated at 70  $\pm$  0.5 MYA based on the combination archaeopyle (Streng et al., 2004) as apomorphic trait, the Thoracosphaera/Pfiesteria clade at 70  $\pm$  0.5 MYA based on the first occurrence of Thoracosphaera in the fossil record (Fensome et al., 1996), and *†Calciodinellum* at 37.5  $\pm$  0.5 MYA based on its tabulated coccoid cells (Hildebrand-Habel and Streng, 2003). For the GTR +  $\Gamma$ substitution model with four discrete categories, we applied an uncorrelated relaxed molecular clock with a diffuse gamma distribution of shape 0.001, scale 1 000.0 and offset 0.0 for the ucld.mean rate. The starting tree was constructed at random, and the final topology was estimated by combining 15 independent chains of 50 million generations each, sampling every 10 000th iteration. TRACER version 1.6 (http://tree.bio.ed.ac.uk/software/ tracer/) was used to evaluate ESS values and to confirm adequate combining of the Markov chain Monte Carlo chains with an appropriate burn-in (10%). All dating analyses were again performed at the CIPRES Science Gateway computing facility (Miller et al., 2010).

#### Acknowledgements

We thank Sylvia Söhner (Einbeck) for the establishment of strains and the generation of sequences, Nils Kleemeier (Munich) for assistance in the molecular lab, and Nina Simanovic (Maribor) for improving the English version of the manuscript. This work was supported by the Deutsche Forschungsgemeinschaft (grant GO 1459 10-1).

#### References

Alverson, A.J., Jansen, R.K., and Theriot, E.C. (2007) Bridging the Rubicon: phylogenetic analysis reveals repeated colonizations of marine and fresh waters by thalassiosiroid diatoms. *Mol Phylogenet Evol* **45**: 193–210.

- Annenkova, N.V., Hansen, G., Moestrup, Ø., and Rengefors, K. (2015) Recent radiation in a marine and freshwater dinoflagellate species flock. *ISME J* 9: 1821–1834.
- Barron, J., and Baldauf, J.G. (1989) Tertiary cooling steps and paleoproductivity as reflected by diatoms and biosiliceous sediments. In *Productivity of the Oceans: Present and Past*, Berger, W.H., Smetacek, V.S., and Wefer, G. (eds). New York: Wiley, pp. 341–354.
- Bass, D., Richards, T.A., Matthai, L., Marsh, V., and Cavalier-Smith, T. (2007) DNA evidence for global dispersal and probable endemicity of protozoa. *BMC Evol Biol* **7**: 162.
- Batten, D.J. (1989) Cretaceous freshwater dinoflagellates. *Cretac Res* **10**: 271–274.
- Berney, C., Romac, S., Mahé, F., Santini, S., Siano, R., and Bass, D. (2013) Vampires in the oceans: predatory cercozoan amoebae in marine habitats. *ISME J* 7: 2387–2399.
- Bison, K.M., Versteegh, G.J.M., Hilgen, F.J., and Willems, H. (2007) Calcareous dinoflagellate turnover in relation to the Messinian salinity crisis in the eastern Mediterranean Pissouri Basin, Cyprus. *J Micropalaeontol* **26**: 103–116.
- Bråte, J., Klaveness, D., Rygh, T., Jakobsen, K.S., and Shalchian-Tabrizi, K. (2010) Telonemia-specific environmental 18S rDNA PCR reveals unknown diversity and multiple marine-freshwater colonizations. *BMC Microbiol* **10**: 168.
- Burkholder, J.M., and Marshall, H.G. (2012) Toxigenic *Pfiesteria* species updates on biology, ecology, toxins, and impacts. *Harmful Algae* **14**: 196–230.
- Calado, A.J., Craveiro, S.C., Daugbjerg, N., and Moestrup, Ø. (2009) Description of *Tyrannodinium* gen. nov., a freshwater dinoflagellate closely related to the marine *Pfiesteria*-like species. *J Phycol* **45**: 1195–1205.
- Cavalier-Smith, T. (2009) Megaphylogeny, cell body plans, adaptive zones: causes and timing of eukaryote basal radiations. *J Eukaryot Microbiol* **56**: 26–33.
- Costanza, R. (1999) The ecological, economic, and social importance of the oceans. *Ecol Econ* **31**: 199–213.
- Craveiro, S.C., Pandeirada, M.S., Daugbjerg, N., Moestrup, Ø., and Calado, A.J. (2013) Ultrastructure and phylogeny of *Theleodinium calcisporum* gen. et sp. nov., a freshwater dinoflagellate that produces calcareous cysts. *Phycologia* **52**: 488–507.
- Darriba, D., Taboada, G.L., Doallo, R., and Posada, D. (2012) jModelTest 2: more models, new heuristics and parallel computing. *Nat Methods* **9**: 772.
- Dittami, S.M., Heesch, S., Olsen, J.L., and Collen, J. (2017) Transitions between marine and freshwater environments provide new clues about the origins of multicellular plants and algae. *J Phycol* **53**: 731–745.
- Dolan, J.R. (2005) An introduction to the biogeography of aquatic microbes. *Aquat Microb Ecol* **41**: 39–48.
- Downing, J.A., Prairie, Y.T., Cole, J.J., Duarte, C.M., Tranvik, L.J., Striegl, R.G., *et al.* (2006) The global abundance and size distribution of lakes, ponds, and impoundments. *Limnol Oceanogr* **51**: 2388–2397.
- Drummond, A.J., Suchard, M.A., Xie, D., and Rambaut, A. (2012) Bayesian phylogenetics with BEAUti and the BEAST 1.7. *Mol Biol Evol* **29**: 1969–1973.
- Elbrächter, M., Gottschling, M., Hildebrand-Habel, T., Keupp, H., Kohring, R., Lewis, J., *et al.* (2008)

Establishing an Agenda for Calcareous Dinoflagellate Research (Thoracosphaeraceae, Dinophyceae) including a nomenclatural synopsis of generic names. *Taxon* **57**: 1289–1303.

- Fawcett, R.C., and Parrow, M.W. (2012) Cytological and phylogenetic diversity in freshwater *Esoptrodinium/Bernardinium* species (Dinophyceae). J Phycol 48: 793–807.
- Fensome, R.A., Taylor, F.J.R., Norris, G., Sarjeant, W.A.S., Wharton, D.I., and Williams, G.L. (1993) A classification of living and fossil dinoflagellates. Micropaleontology Special Publication, Vol. 7. New York, NY: American Museum of Natural History, pp. 1–245.
- Fensome, R.A., MacRae, R.A., Moldowan, J.M., Taylor, F.J. R., and Williams, G.L. (1996) The early Mesozoic radiation of dinoflagellates. *Paleobiology* **22**: 329–338.
- Fensome, R.A., Saldarriaga, J.F., and Taylor, F.J.R. (1999) Dinoflagellate phylogeny revisited: reconciling morphological and molecular based phylogenies. *Grana* **38**: 66–80.
- Feulner, G., Hallmann, C., and Kienert, H. (2015) Snowball cooling after algal rise. *Nat Geosci* **8**: 659–662.
- Finlay, B.J., Esteban, G.F., Brown, S., Fenchel, T., and Hoef-Emden, K. (2006) Multiple cosmopolitan ecotypes within a microbial eukaryote morphospecies. *Protist* 157: 377–390.
- Gleick, P.H. (1999) The human right to water. *Water Policy* **1**: 487–503.
- Gottschling, M., and Söhner, S. (2013) An updated list of generic names in the Thoracosphaeraceae. *Microorganisms* **1**: 122–136.
- Gottschling, M., Keupp, H., Plötner, J., Knop, R., Willems, H., and Kirsch, M. (2005) Phylogeny of calcareous dinoflagellates as inferred from ITS and ribosomal sequence data. *Mol Phylogenet Evol* **36**: 444–455.
- Gottschling, M., Renner, S.S., Meier, K.J.S., Willems, H., and Keupp, H. (2008) Timing deep divergence events in calcareous dinoflagellates. *J Phycol* **44**: 429–438.
- Gottschling, M., Söhner, S., Zinßmeister, C., John, U., Plötner, J., Schweikert, M., *et al.* (2012) Delimitation of the Thoracosphaeraceae (Dinophyceae), including the calcare-ous dinoflagellates, based on large amounts of ribosomal RNA sequence data. *Protist* **163**: 15–24.
- Gottschling, M., Kretschmann, J., and Žerdoner Čalasan, A. (2017) Description of Peridiniopsidaceae, fam. nov. (Peridiniales, Dinophyceae). *Phytotaxa* **299**: 293–296.
- Gouy, M., Guindon, S., and Gascuel, O. (2010) SeaView version 4: a multiplatform graphical user interface for sequence alignment and phylogenetic tree building. *Mol Biol Evol* **27**: 221–224.
- Gradstein, F.M., Ogg, J.G., Schmitz, M.D., and Ogg, G. (2012) *The Geologic Time Scale 2012*. Amsterdam: Elsevier.
- Gray, J., and Taylor, D.W. (1988) Evolution of the freshwater ecosystem: the fossil record. *Palaeogeogr Palaeoclimatol Palaeoecol* **62**: 1–214.
- Gu, H., Kirsch, M., Zinßmeister, C., Söhner, S., Meier, K.J.
  S., Liu, T., and Gottschling, M. (2013) Waking the dead: morphological and molecular characterization of extant *†Posoniella tricarinelloides* (Thoracosphaeraceae, Dinophyceae). *Protist* **164**: 583–597.
- Gu, H., Mertens, K.N., and Liu, T. (2016) *Huia caspica* gen. & comb. nov., a dinoflagellate species that recently
- © 2019 Society for Applied Microbiology and John Wiley & Sons Ltd., Environmental Microbiology, 21, 4125–4135

4134 A. Žerdoner Čalasan, J. Kretschmann and M. Gottschling

crossed the marine-freshwater boundary. *Phycolological Res* **64**: 251–258.

- Guillard, R.R., and Lorenzen, C.J. (1972) Yellow-green algae with chlorophyllide c. J Phycol 8: 10–14.
- Hansen, G., Daugbjerg, N., and Henriksen, P. (2007) Baldinia anauniensis gen. et sp. nov.: a 'new' dinoflagellate from Lake Tovel, N. Italy. *Phycologia* 46: 86–108.
- Heger, T.J., Mitchell, E.A.D., Todorov, M., Golemansky, V., Lara, E., Leander, B.S., and Pawlowski, J. (2010) Molecular phylogeny of euglyphid testate amoebae (Cercozoa: Euglyphida) suggests transitions between marine supralittoral and freshwater/terrestrial environments are infrequent. *Mol Phylogenet Evol* **55**: 113–122.
- Hildebrand-Habel, T., and Streng, M. (2003) Calcareous dinoflagellate associations and Maastrichtian–Tertiary climatic change in a high-latitude core (ODP hole 689B, Maud rise, Weddell Sea). *Palaeogeogr Palaeoclimatol Palaeoecol* **197**: 293–321.
- Hoppenrath, M., Murray, S.A., and Chomérat, N. (2014) Marine Benthic Dinoflagellates – Unveiling their Worldwide Biodiversity. Stuttgart: Schweizerbart.
- Janouškovec, J., Gavelis, G.S., Burki, F., Dinh, D., Bachvaroff, T.R., Gomik, S.G., *et al.* (2017) Major transitions in dinoflagellate evolution unveiled by phylo-transcriptomics. *Proc Natl Acad Sci USA* **114**: 171–180.
- Jardillier, L., Zubkov, M.V., Pearman, J., and Scanlan, D.J. (2010) Significant CO<sub>2</sub> fixation by small prymnesiophytes in the subtropical and tropical Northeast Atlantic Ocean. *ISME J* 4: 1180–1192.
- John, U., Fensome, R.A., and Medlin, L.K. (2003) The application of a molecular clock based on molecular sequences and the fossil record to explain biogeographic distributions within the *Alexandrium tamarense* "species complex" (Dinophyceae). *Mol Biol Evol* 20: 1015–1027.
- Katoh, K., and Standley, D.M. (2013) MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Mol Biol Evol* **30**: 772–780.
- Keupp, H. (1984) Revision der kalkigen Dinoflagellaten-Zysten G. DEFLANDREs, 1948. Paläontol Z 58: 9–31.
- Keupp, H., and Ilg, A. (1989) Die kalkigen Dinoflagellaten im Ober-Callovium und Oxfordium der Normandie/Frankreich. Berl Geowiss Abh, Reihe A 106: 165–205.
- Kremp, A., Tahvanainen, P., Litaker, W., Krock, B., Suikkanen, S., Leaw, C.P., and Tomas, C. (2014) Phylogenetic relationships, morphological variation, and toxin patterns in the *Alexandrium ostenfeldii* (Dinophyceae) complex: implications for species boundaries and identities. *J Phycol* **50**: 81–100.
- Kretschmann, J., Filipowicz, N.H., Owsianny, P.M., Zinßmeister, C., and Gottschling, M. (2015) Taxonomic clarification of the unusual dinophyte *Gymnodinium limneticum* Wołosz. (Gymnodiniaceae) from the Tatra Mountains. *Protist* **166**: 621–637.
- Kretschmann, J., Žerdoner Čalasan, A., and Gottschling, M. (2018) Molecular phylogenetics of dinophytes harbouring diatoms as endosymbionts (Kryptoperidiniaceae, Peridiniales), with evolutionary interpretations and a focus on the identity of *Durinskia oculata* from Prague. *Mol Phylogenet Evol* **118**: 392–402.

- LaJeunesse, T.C. (2004) "Species" radiations of symbiotic dinoflagellates in the Atlantic and indo-Pacific since the Miocene-Pliocene transition. *Mol Biol Evol* **22**: 570–581.
- LaJeunesse, T.C., Parkinson, J.E., Gabrielson, P.W., Jeong, H.J., Reimer, J.D., Voolstra, C.R., and Santos, S. R. (2018) Systematic revision of Symbiodiniaceae highlights the antiquity and diversity of coral endosymbionts. *Curr Biol* 28: 2570–2580.
- Lear, C.H., Bailey, T.R., Pearson, P.N., Coxall, H.K., and Rosenthal, Y. (2008) Cooling and ice growth across the Eocene-Oligocene transition. *Geology* **36**: 251–254.
- Leliaert, F., Smith, D.R., Moreau, H., Herron, M.D., Verbruggen, H., Delwiche, C.F., and De Clerck, O. (2012) Phylogeny and molecular evolution of the green algae. *Crit Rev Plant Sci* **31**: 1–46.
- Lewis, L.A., and McCourt, R.M. (2004) Green algae and the origin of land plants. *Am J Bot* **91**: 1535–1556.
- Lindner, L., Gozhik, P., Marciniak, B., Marks, L., and Yelovicheva, Y. (2004) Main climatic changes in the Quaternary of Poland, Belarus and Ukraine. *Geol Q* **48**: 97–114.
- Litaker, R.W., Steidinger, K.A., Mason, P.L., Landsberg, J. H., Shields, J.D., Reece, K.S., *et al.* (2005) The reclassification of *Pfiesteria shumwayae* (Dinophyceae): *Pseudopfiesteria*, gen. nov. *J Phycol* **41**: 643–651.
- Logares, R., Rengefors, K., Kremp, A., Shalchian-Tabrizi, K., Boltovskoy, A., Tengs, T., *et al.* (2007a) Phenotypically different microalgal morphospecies with identical ribosomal DNA: a case of rapid adaptive evolution? *Microb Ecol* **53**: 549–561.
- Logares, R., Shalchian-Tabrizi, K., Boltovskoy, A., and Rengefors, K. (2007b) Extensive dinoflagellate phylogenies indicate infrequent marine-freshwater transitions. *Mol Phylogenet Evol* **45**: 887–903.
- Logares, R., Daugbjerg, N., Boltovskoy, A., Kremp, A., Laybourn-Parry, J., and Rengefors, K. (2008) Recent evolutionary diversification of a protist lineage. *Environ Microbiol* **10**: 1231–1243.
- Logares, R., Bråte, J., Bertilsson, S., Clasen, J.L., Shalchian-Tabrizi, K., and Rengefors, K. (2009) Infrequent marine–freshwater transitions in the microbial world. *Trends Microbiol* **17**: 414–422.
- Logares, R., Bråte, J., Heinrich, F., Shalchian-Tabrizi, K., and Bertilsson, S. (2010) Infrequent transitions between saline and fresh waters in one of the most abundant microbial lineages (SAR11). *Mol Biol Evol* **27**: 347–357.
- Martens, H., Van De Waal, D.B., Brandenburg, K.M., Krock, B., and Tillmann, U. (2016) Salinity effects on growth and toxin production in an *Alexandrium ostenfeldii* (Dinophyceae) isolate from The Netherlands. *J Plankton Res* **38**: 1302–1316.
- Matthews, J.A. (2013) Encyclopedia of Environmental Change: Three Volume Set. Thousand Oaks, CA: SAGE.
- Medlin, L.K. (2008) Proceedings of the International Symposium "the Origin and Evolution of Natural Diversity", 1–5 October 2007; Sapporo, Japan.
- Mertens, K.N., Rengefors, K., Moestrup, Ø., and Ellegaard, M. (2012) A review of recent freshwater dinoflagellate cysts: taxonomy, phylogeny, ecology and palaeocology. *Phycologia* 51: 612–619.

- Miller, K.G., Wright, J.D., and Fairbanks, R.G. (1991) Unlocking the ice house: Oligocene-Miocene oxygen isotopes, eustasy, and margin erosion. J Geophys Res 96: 6829–6848.
- Miller, M.A., Pfeiffer, W., and Schwartz, T. (2010) Proceedings of the Gateway Computing Environments Workshop (GCE). New Orleans, Louisiana; November 16–21, 2014.
- Moestrup, Ø., and Calado, A.J. (2018) *Dinophyceae*. Berlin: Springer.
- Moestrup, Ø., and Daugbjerg, N. (2007) On dinoflagellate phylogeny and classification. In *Unravelling the algae, the past, present, and future of algal systematics*, Brodie, J., and Lewis, J. (eds). Boca Raton, FL: CRC, pp. 215–230.
- Mulkidjanian, A.Y., Bychkov, A.Y., Dibrova, D.V., Galperin, M.Y., and Koonin, E.V. (2012) Origin of first cells at terrestrial, anoxic geothermal fields. *Proc Natl Acad Sci USA* **109**: 821–830.
- Murray, S., Jørgensen, M.F., Ho, S.Y.W., Patterson, D.J., and Jermiin, L.S. (2005) Improving the analysis of dinoflagellate phylogeny based on rDNA. *Protist* **156**: 269–286.
- Myers, G.S. (1949) Usage of anadromous, catadromous and allied terms for migratory fishes. *Copeia* **2**: 89–97.
- Orr, R.J.S., Murray, S.A., Stüken, A., Rhodes, L., and Jakobsen, K.S. (2012) When naked became armored: an eight-gene phylogeny reveals monophyletic origin of theca in dinoflagellates. *PLoS One* **7**: e50004.
- Parfrey, L.W., Lahr, D.J.G., Knoll, A.H., and Katz, L.A. (2011) Estimating the timing of early eukaryotic diversification with multigene molecular clocks. *Proc Natl Acad Sci* USA 108: 13624–13629.
- Paulay, G., and Meyer, C. (2002) Diversification in the tropical pacific: comparisons between marine and terrestrial systems and the importance of founder speciation. *Integr Comp Biol* **42**: 922–934.
- Pelzer, F. (1991) *Polen: Eine geographische Landeskunde.* Darmstadt: Wissenschaftliche Buchgesellschaft.
- Pochon, X., Montoya-Burgos, J.I., Stadelmann, B., and Pawlowski, J. (2006) Molecular phylogeny, evolutionary rates, and divergence timing of the symbiotic dinoflagellate genus *Symbiodinium*. *Mol Phylogenet Evol* **38**: 20–30.
- Pokorný, P. (2009) Evolution and function of freshwater ecosystems. In *Earth System: History and Natural Variability*, Cílek, V. (ed). Oxford: Eolss, pp. 118–137.
- Popescu, S.-M., Dalesme, F., Jouannic, G., Escarguel, G., Head, M.J., Melinte-Dobrinescu, M.C., *et al.* (2009) *Galeacysta etrusca* complex: Dinoflagellate cyst marker of paratethyan influxes to the Mediterranean Sea before and after the peak of the messinian salinity crisis. *Palynology* **33**: 105–134.
- Price, D.C., and Bhattacharya, D. (2017) Robust Dinoflagellata phylogeny inferred from public transcriptome databases. *J Phycol* **53**: 725–729.
- Riegel, W. (2008) The late Palaeozoic phytoplankton blackout – artefact or evidence of global change? *Rev Palaeobot Palynol* **148**: 73–90.
- Ronquist, F., Teslenko, M., van der Mark, P., Ayres, D.L., Darling, A., Höhna, S., *et al.* (2012) MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice across a large model space. *Syst Biol* **61**: 539–542.
- Saldarriaga, J.F., Taylor, F.J.R., Cavalier-Smith, T., Menden-Deuerd, S., and Keeling, P.J. (2004) Molecular data and

the evolutionary history of dinoflagellates. *Eur J Protistol* **40**: 85–111.

- Sánchez-Baracaldo, P., Raven, J.A., Pisani, D., and Knoll, A.H. (2017) Early photosynthetic eukaryotes inhabited low-salinity habitats. *Proc Natl Acad Sci USA* **114**: 7737–7745.
- Servais, T., Martin, R.E., and Nützel, A. (2016) The impact of the 'terrestrialisation process' in the late Palaeozoic: *p*CO<sub>2</sub>, *p*O<sub>2</sub>, and the 'phytoplankton blackout'. *Rev Palaeobot Palynol* **224**: 26–37.
- Shalchian-Tabrizi, K., Bråte, J., Logares, R., Klaveness, D., Berney, C., and Jakobsen, K.S. (2008) Diversification of unicellular eukaryotes: cryptomonad colonizations of marine and fresh waters inferred from revised 18S rRNA phylogeny. *Environ Microbiol* **10**: 2635–2644.
- Shalchian-Tabrizi, K., Reier-Roberg, K., Ree, D.K., Klaveness, D., and Bråte, J. (2011) Marine-freshwater colonizations of haptophytes inferred from phylogeny of environmental 18S rDNA sequences. *J Eukaryot Microbiol* 58: 315–318.
- Simó, R. (2001) Production of atmospheric sulfur by oceanic plankton: biogeochemical, ecological and evolutionary links. *Trends Ecol Evol* **16**: 287–294.
- Simon, M., Jardillier, L., Deschamps, P., Moreira, D., Restoux, G., Bertolino, P., and López-García, P. (2015) Complex communities of small protists and unexpected occurrence of typical marine lineages in shallow freshwater systems. *Environ Microbiol* **17**: 3610–3627.
- Sims, P.A., Mann, D.G., and Medlin, L.K. (2006) Evolution of the diatoms: insights from fossil, biological and molecular data. *Phycologia* **45**: 361–402.
- Snoke, M.S., Berendonk, T.U., Barth, D., and Lynch, M. (2006) Large global effective population sizes in *Paramecium*. *Mol Biol Evol* 23: 2474–2479.
- Stamatakis, A. (2014) RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* **30**: 1312–1313.
- Streng, M., Hildebrand-Habel, T., and Willems, H. (2004) A proposed classification of archeopyle types in calcareous dinoflagellate cysts. *J Paleo* **78**: 456–483.
- Taylor, F.J.R., Hoppenrath, M., and Saldarriaga, J.F. (2008) Dinoflagellate diversity and distribution. *Biodivers Conserv* **17**: 407–418.
- Tennant, J.P., Mannion, P.D., Upchurch, P., Sutton, M.D., and Price, G.D. (2017) Biotic and environmental dynamics through the late Jurassic-early cretaceous transition: evidence for protracted faunal and ecological turnover. *Biol Rev* 92: 776–814.
- Žerdoner Čalasan, A., Kretschmann, J., and Gottschling, M. (2018) Absence of co-phylogeny indicates repeated diatom capture in dinophytes hosting a tertiary endosymbiont. *Org Divers Evol* **18**: 29–38.

# **Supporting Information**

Additional Supporting Information may be found in the online version of this article at the publisher's web-site:

Appendix S1: Supporting Information