

RESEARCH ARTICLE

Schistosoma haematobium infection and environmental factors in Southwestern Tanzania: A cross-sectional, population-based study

Kirsi M. Manz^{1*}, Inge Kroidl^{2,3,4}, Petra Clowes^{2,3}, Martina Gerhardt^{2,3}, Wilbrod Nyembe³, Lucas Maganga³, Weston Assisya³, Nyanda E. Ntinginya³, Ursula Berger¹, Michael Hoelscher^{2,4}, Elmar Saathoff^{2,4}

1 Institute for Medical Information Processing, Biometry, and Epidemiology (IBE), Ludwig-Maximilians University, Munich, Germany, **2** Division of Infectious Diseases and Tropical Medicine, University Hospital, LMU Munich, Germany, **3** NIMR-Mbeya Medical Research Center (MMRC), Mbeya, Tanzania, **4** German Center for Infection Research (DZIF), partner site Munich, Munich, Germany

* manz@ibe.med.uni-muenchen.de



OPEN ACCESS

Citation: Manz KM, Kroidl I, Clowes P, Gerhardt M, Nyembe W, Maganga L, et al. (2020) *Schistosoma haematobium* infection and environmental factors in Southwestern Tanzania: A cross-sectional, population-based study. PLoS Negl Trop Dis 14(8): e0008508. <https://doi.org/10.1371/journal.pntd.0008508>

Editor: Stefanie Knopp, Swiss Tropical and Public Health Institute, SWITZERLAND

Received: October 18, 2019

Accepted: June 22, 2020

Published: August 24, 2020

Copyright: © 2020 Manz et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: The data used for this analysis contain the GPS positions of all participating households, which could be used to find and identify participating households and individuals. Thus making them publicly available would constitute a severe breach of confidentiality. Data are therefore only available on request. Please contact Dr. Friedrich Rieß, (email: Friedrich.Riess@lrz.uni-muenchen.de), Division of Infectious Diseases and Tropical Medicine, University

Abstract

Schistosomiasis is a leading cause of morbidity in Africa. Understanding the disease ecology and environmental factors that influence its distribution is important to guide control efforts. Geographic information systems have increasingly been used in the field of schistosomiasis environmental epidemiology. This study reports prevalences of *Schistosoma haematobium* infection and uses remotely sensed and questionnaire data from over 17000 participants to identify environmental and socio-demographic factors that are associated with this parasitic infection. Data regarding socio-demographic status and *S. haematobium* infection were obtained between May 2006 and May 2007 from 17280 participants (53% females, median age = 17 years) in the Mbeya Region, Tanzania. Combined with remotely sensed environmental data (vegetation cover, altitude, rainfall etc.) this data was analyzed to identify environmental and socio-demographic factors associated with *S. haematobium* infection, using mixed effects logistic regression and geostatistical modelling. The overall prevalence of *S. haematobium* infection was 5.3% (95% confidence interval (CI): 5.0–5.6%). Multivariable analysis revealed increased odds of infection for school-aged children (5–15 years, odds ratio (OR) = 7.8, CI: 5.9–10.4) and the age groups 15–25 and 25–35 years (15–25 years: OR = 5.8, CI: 4.3–8.0, 25–35 years: OR = 1.6, CI: 1.1–2.4) compared to persons above 35 years of age, for increasing distance to water courses (OR = 1.4, CI: 1.2–1.6 per km) and for proximity to Lake Nyasa (<1 km, OR = 4.5, CI: 1.8–11.4; 1–2 km, OR = 3.5, CI: 1.7–7.5; 2–4 km; OR = 3.3, CI: 1.7–6.6), when compared to distances >4 km. Odds of infection decreased with higher altitude (OR = 0.7, CI: 0.6–0.8 per 100 m increase) and with increasing enhanced vegetation index EVI (OR = 0.2, CI: 0.1–0.4 per 0.1 units). When additionally adjusting for spatial correlation population density became a significant predictor of schistosomiasis infection (OR = 1.3, CI: 1.1–1.5 per 1000 persons/km²) and altitude turned non-significant. We found highly focal geographical patterns of *S. haematobium* infection in Mbeya Region in Southwestern Tanzania. Despite low overall prevalence our

Hospital, LMU Munich, Leopoldstr. 5, 80802 Munich, Germany.

Funding: This study was supported by grants SANTE/2004/078-545/130 and SANTE/2006/129-931 to MH from the European Commission (https://ec.europa.eu/info/departments/health-and-food-safety_en). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

spatially heterogeneous results show that some of the study sites suffer from a considerable burden of *S. haematobium* infection, which is related to various socio-demographic and environmental factors. Our results could help to design more effective control strategies in the future, especially targeting school-aged children living in low altitude sites and/or crowded areas as the persons at highest need for preventive chemotherapy.

Author summary

Urinary schistosomiasis caused by the parasitic blood fluke *Schistosoma haematobium* is an important public health concern in the tropics and subtropics. Freshwater snails of the genus *Bulinus* are the intermediate hosts and infection is acquired by contact with fresh water inhabited by infected snails. This means that transmission of the parasite strongly depends on environmental factors, which we wanted to explore in this study. Individually assessed questionnaire and laboratory data from a study in Mbeya Region in Southwestern Tanzania were combined with remotely sensed and other environmental data that were publicly available. We found that the odds of urinary schistosomiasis were higher in school-aged children and young adults than in older age groups and in children below 5 years. Odds of infection were also higher at lower altitudes, in areas with less vegetation and in the vicinity of Lake Nyasa. These results add to the general knowledge regarding *S. haematobium* infection and could help to design more effective control strategies in the future, both locally and on larger scales.

Introduction

Trematodes of the genus *Schistosoma* are among the most common infectious agents of humans. They cause schistosomiasis which occurs in 78 tropical and sub-tropical countries. The highest prevalences are encountered in Sub-Saharan-Africa (SSA), where according to recent estimates there are 112 million cases of *Schistosoma haematobium* infection and 54 million cases of *Schistosoma mansoni* infection [1]. The current main strategy to combat schistosomiasis in SSA and other developing parts of the world is mass administration of praziquantel to school age children, with the aim to reduce the disease burden. According to WHO, at least 221 million people world-wide and 200 million in Africa required preventive mass-treatment for schistosomiasis in 2017, out of which 102 million world-wide and 91 million people in Africa were reported to have been treated [1].

Schistosomiasis infection occurs through contact with fresh water contaminated with the free-swimming larval forms (cercariae) of the parasite. The cercariae penetrate the skin and mature to adulthood inside the human body. Adult worms live in the veins draining the urinary tract (*S. haematobium*) or the intestine (other species). The eggs, which are passed into the environment with the feces or urine, hatch in fresh water and release so-called miracidia, which infect suitable host snails. The snail in turn releases cercariae which infect humans during contact with fresh water. However, many of the eggs are trapped in the human host's tissues and the body's reaction to them can cause massive tissue damage. Thus trapped eggs are the main cause of morbidity in schistosome infection [2].

S. haematobium, the agent causing urinary schistosomiasis, only occurs in Africa and the Middle East, where it is the most common schistosome species [3,4]. Consequences of urinary schistosomiasis include hematuria, dysuria, bladder scarring, chronic urinary tract infection

and possibly bladder cancer [5,6]. Up to three-quarters of women with *S. haematobium* infection are estimated to suffer from female genital schistosomiasis when eggs accumulate in the vagina, cervix, uterus or fallopian tubes [7–9]. Environmental conditions that are permissive to the development of intermediate host snails are important factors for schistosomiasis transmission in endemic regions. Advances in remote sensing (RS) and geographic information systems (GIS) have enabled researchers to explore these environmental and climatic factors in greater depth. Risk mapping, with the aid of RS and GIS applications, is suited to the study of schistosomiasis as the infectious agents and their snail hosts are sensitive to environmental conditions [10, 11]. When exploring the associations between environmental factors and infection with schistosomiasis the spatial correlation should be taken into account by employing geostatistical models [12–20].

To capture the effects of individual factors and especially the effects of the surrounding environment on the risk of schistosomiasis infection, the situation before any large-scale preventive chemotherapy should be considered. The aim of this study was to report pre-treatment prevalences of *S. haematobium* infection and to identify individual factors and local environmental conditions that might influence *S. haematobium* transmission in Mbeya Region in Southwestern Tanzania.

Methods

Ethics

The EMINI (Evaluating and Monitoring the Impact of New Interventions) cohort study was approved by the ethics committee of the Tanzanian National Institute for Medical Research and conducted according to the Declaration of Helsinki. All participants signed/thumb printed a written informed consent before enrollment into the study, with parents consenting for their minor children below 18 years of age. Children between 12 and 18 years additionally signed/thumb printed the consent document and younger children who were old enough to understand the process were asked to participate in the consenting procedure as well.

Study area and epidemiological data collection

The study area is located in the Mbeya Region in Southwestern Tanzania and extends from 32.678° to 33.963° East and from 8.652° to 9.649° South as shown in Fig 1. Fig 2 provides an overview of the study area, the study sites and the participating households. Below we briefly describe the study area and data collection methods, more detailed accounts and results for other helminth infections in this population are provided elsewhere [21–23].

Prior the start of the EMINI study the study area and the nine study sites were preselected to cover a wide variety of economic and environmental conditions. This was followed by a census of the nine selected sites, covering more than 42,000 households. Household positions were determined using handheld GPS devices. After the census of all households in each site was completed, 10% of the households within these sites were chosen as a geographically stratified random sample to participate in the study. Using the geographical position, each household was visited annually starting in 2006, and all household members who were present during each visit were asked to participate in the study. Within the same annual survey, households were only revisited if the initial visit was completely unsuccessful, i.e. if no household member had been present.

During the survey visits we collected blood and urine samples to test for HIV and schistosomiasis, and performed structured interviews in the local language (Kiswahili). Additional collection of stool samples to study soil-transmitted helminth infections started at the third annual visit from 2008 onwards in 50% of the households. For the analysis of *S. haematobium*




Fig 1. Map of Africa showing the location of Tanzania (A) and map of Tanzania indicating where Mbeya Region (green area) and the study area (red square) are located (B). (Original work of the authors).

<https://doi.org/10.1371/journal.pntd.0008508.g001>

infection shown here, we used data from the first annual survey which was conducted between May 2006 and May 2007. To our knowledge no larger schistosomiasis control efforts had been implemented in the region prior to this study.

Urine samples were processed as follows: Morning and spot urine samples were collected from each participant and 10 ml of urine from each of the two samples were filtered through polyamide mesh with 20 μm pore size. The filters were stained with Logol's iodine and microscopically examined for *S. haematobium* eggs. All participants with urine samples with at least one *S. haematobium* egg were regarded as infected. *S. haematobium* infection intensity was recorded in eggs per centiliter (EPC) and classified into no (0 EPC), light intensity (<50 EPC) and heavy intensity infections (≥ 50 EPC) according to World Health Organization guidelines [24]. Infected participants were offered standard treatment with praziquantel. HIV testing was done using the Determine HIV 1/2 rapid test (Abbott Laboratories, Abbott Park, IL) followed by confirmatory ELISA testing (Enzygnost Anti HIV 1/2 Plus, DADE-Behring, Marburg, Germany) if the RDT was positive. Western blot testing (MPD HIV Blot 2.2, MP Biomedicals, and Geneva, Switzerland) was performed to resolve conflicting results [25].

The financial situation of a household does not necessarily reflect the actual socio-economic status (SES) in low resource settings, particularly because it is difficult to get reliable information on household income and expenditure. To get a more reliable estimate of the SES, we applied a modified method initially proposed by Filmer and Pritchett [26, 27] that uses polychoric principal component analysis to combine different proxies for household wealth into an SES score. The following items were used to construct the score: household belongings (clock or watch, radio, television, mobile telephone, refrigerator, hand cart, bicycle, motor




Fig 2. Map of the study area showing the nine distinct study sites. (Original work of the authors).

<https://doi.org/10.1371/journal.pntd.0008508.g002>

cycle, car, savings account), materials used to build the house, sources of energy and drinking water, number of persons per room. The necessary data were obtained during the household interviews.

Ecological data

Data regarding the larger lakes in the area and elevation data, which were also used to calculate slope of the terrain, were retrieved from NASA's Shuttle Radar Topography Mission (SRTM) global digital elevation model, version 2.1 with a nominal resolution of 90 m [28]. Information on water courses was obtained from Vector Map Level 0 (VMAP0) data by the US National Imagery and Mapping Agency. Water courses were defined as elongated flowing water bodies such as streams and rivers, in contrast to ponds or lakes.

Data regarding land surface temperature during the day (LST day) and night (LST night), and green vegetation cover (EVI = enhanced vegetation index) had been collected by NASA's Moderate-Resolution Imaging Spectroradiometer (MODIS) Terra satellite [29]. LST data (version MOD11A2) have 8 days temporal and about 1 km spatial resolution. Vegetation data (version MOD13Q1) have 16 days temporal and 250 m spatial resolution. Data for the years 2003 through 2008 were retrieved from the online data pool, courtesy of the NASA EOSDIS Land Processes Distributed Active Archive Center (LP DAAC), USGS/Earth Resources Observation and Science (EROS) Center, Sioux Falls, South Dakota (<https://lpdaac.usgs.gov/>). Both, LST and vegetation data were processed in the following way to produce long-term averages: data surfaces for every 8-day period (LST) and every 16-day period (EVI) for the years 2003 to 2008 were imported into Idrisi GIS software v.32 (Clark Labs, Worcester, MA, United States of

America). In Idrisi, annual averages, minima and maxima of day- and night-LST and EVI for each of these six years were calculated for every pixel and in turn used to calculate long term averages for each pixel for the whole period 2003 to 2008. The calculated averages, minima and maxima are thus an average of the six annual minima and maxima for each pixel. For this, we utilized only those pixels that were “good data quality” according to the quality assessment layers that are distributed together with the actual raw data. Then LST was converted to °C and EVI was converted back to its native range between -1 and +1.

Mean annual rainfall data with 1 km spatial resolution was obtained from the WorldClim-Global Climate Data website (<http://www.worldclim.org/>).

All above environmental data were then combined with the household position data in a GIS database using Manifold System 8.0 Professional Edition (Manifold Net Ltd, Carson City, NV), which was then used to calculate the following parameters.

Distance to the nearest water course, defined as the shortest distance between the participant’s household and the closest river or stream, and distance to the nearest lake were calculated using the household positions and the above referenced data on lakes and water courses. As elaborated further below, all lakes other than Lake Nyasa were too far away (>4 km) from participant’s homes to play an important role in their daily water contact activities.

Household positions and number of inhabitants that had been collected during the initial population census were used to calculate population densities by dividing the number of inhabitants within a 1 km buffer around the participant’s home by the surface area of this buffer. Similarly, LST, EVI, rainfall and elevation data were averaged for a buffer area of 1000 m radius around each household to characterize the situation in the area where many of the daily activities of participants would be performed. This approach was preferred to using the respective spot values at the household position, because spot data are more prone to random error than averages for a wider area.

Statistical analyses

Descriptive analyses were performed using Stata statistics software (Release 14. College Station, TX: StataCorp LP). Maps exploring the prevalence were generated using Manifold System 8.0 Professional Edition (Manifold Net Ltd, Carson City, NV). Geostatistical models were estimated in R version 3.5.0 [30] using the package “*gamm4*” [31].

Since the majority of *S. haematobium* infections were of light intensity and our primary interest was to identify factors related to the presence/absence of infection, we used a binary (no/yes) infection outcome for most of our models. Mixed effects logistic regression with random effects for the study sites and households was used to report odds ratios, which is a suitable method for analysis of clustered cross-sectional data [32].

The following variable transformations were applied to enhance interpretability of results: The reported odds ratios (ORs) correspond to an increase of 1000 persons/km² for the population density, 100 m for elevation, 100 mm for annual rainfall and 0.1 units for EVI. For EVI, LST day and LST night we considered minimum, average and maximum values and included the representation of the respective factor leading to the model with the lowest Akaike Information Criterion (AIC). Age of the participants was stratified into five categories based on the typical *S. haematobium* infection patterns over age. Distance to lake was also calculated from remotely sensed data, which however only feature large waterbodies, but do not include small ponds and pools, which also play an important role in schistosomiasis transmission. This variable was stratified into four categories based on the assumed relevance with regard to water contact activities; distances of 4 km and more were deemed too far away for daily lake-water contact. Since only participants in Kyela lived closer than 4 km to a lake (in this case Lake

Nyasa), the three lower strata of this variable only include part of the population of Kyela site. All other participants lived more than 4 km away from any lake, including Lake Nyasa. In addition to HIV positive and negative participants, the participants with missing or indecisive HIV test results (371 in total) were included as an additional “no information” stratum into the analysis. A total of 30 observations were discarded due to missing values regarding *S. haematobium* infection status or other information, resulting in complete data from 17,280 participants.

First, univariable mixed effects logistic regression with study site and household as random effects was performed to estimate odds ratios of *S. haematobium* infection with their 95% confidence intervals for each of the covariates of interest. The random effects were included to account for within-household and within-site clustering of infection.

For our initial multivariable “base” model we included the individual factors age, sex, SES and HIV status (see [S1 Table](#)). These variables were included as potential confounders based on their potential relevance for schistosomiasis infection and were left in the model, regardless of their association with *S. haematobium* infection. HIV infection was included not only as a potential confounder, but also because we wanted to explore its relationship with schistosomiasis, since results in the literature regarding this are conflicting. Then, step by step, we included each univariably assessed covariate and left it in the model if the model’s AIC decreased, to identify the most parsimonious model. For more details on the variable selection see [S1 Text](#).

Since environmental data is prone to be highly correlated, we checked for potentially collinear variables by calculating the variance inflation factor (VIF) at each multivariable analysis step. A $VIF > 10$ is mentioned in the literature as an indicator for serious collinearity [33, 34], thus variables with $VIF \geq 10$ were not simultaneously entered into the multivariable models.

Spatial autocorrelation in general refers to situations where nearby observations are more similar than observations further away from each other. Spatial autocorrelation is present in our data, if the probability to be infected with *S. haematobium* is clustered in space, which means that it depends on the location where the participant lives. To account for this, we employed geostatistical modelling [35]. We estimated mixed generalized additive logistic models and included a spatially correlated effects base (*i.e.* a spatial smoother) on the location of the households and additionally adjusted for clustering of infection by including random effects for study site and household level, respectively. Geostatistical models were estimated for all univariable models, for the multivariable model with the lowest AIC and for the full model including all non-collinear variables (for the full model see [S2 Table](#)).

The association of HIV status with the number of excreted *S. haematobium* eggs was investigated by means of uni- and multi-variable mixed effects negative binomial regression in participants infected with *S. haematobium*. The negative binomial model was chosen to account for the overdispersion present in the data, since the variance of the egg counts was much larger than the mean [36]. We used the egg counts (number of eggs per centiliter of urine) as outcome and HIV status as the main predictor with random effects for study site and household and additionally adjusted for age, sex and SES.

Results

Descriptive statistics

Data from 17280 participants from 4189 households were used for this analysis. The median age of the study population was 17 years (interquartile range (IQR): 9–34 years) and 53% (9194/17280) were females ([Table 1](#)). The overall prevalence of *S. haematobium* infection in all nine sites was 5.3% (914/17280, 95% confidence interval (CI): 5.0–5.6%), ranging from 0.0 to 15.8% per site ([Fig 3](#) and [Table 2](#)). In total the infection occurred in 13.8% of all households

Table 1. Characteristics of study participants and environmental conditions at their places of residence.

Variable (Unit)		All sites	Igurusi	Santylia
Number of participants	N	17280	1947	2003
Sex female	%	53	52	55
Age (years)	Median (IQR)	17 (9–34)	19 (9–36)	17 (8–34)
<i>S. haematobium</i> infection prevalence	%	5.3	15.8	0
Infection intensity:				
No infection (0 eggs/10 ml)	%	94.7	84.2	100
Light infection (1–49 eggs/10 ml)	%	4.7	12.7	0
Heavy infection (≥ 50 eggs/10 ml)	%	0.6	3.1	0
HIV infection				
No	%	90.9	88.7	91.8
Yes	%	7.0	6.7	5.6
No information	%	2.1	4.6	2.6
SES score				
	Median (IQR)	-0.10 (-0.57–0.46)	-0.09 (-0.54–0.48)	-0.44 (-0.78 –0.07)
Population density (persons/km²)				
	Median (IQR)	442 (221–2122)	1739 (732–2279)	205 (132–285)
Household with latrine				
No	%	2.4	1.0	1.7
Yes	%	97.6	99.0	98.3
Elevation (m)				
	Median (IQR)	1579 (1216–1728)	1193 (1156–1205)	2025 (1982–2066)
LST Day (°C)				
	Median (IQR)	33.2 (30.4–33.9)	34.6 (34.2–36.2)	29.9 (29.3–30.8)
LST Night (°C)				
	Median (IQR)	14.0 (11.8–16.0)	16.2 (16.0–16.5)	11.7 (10.9–12.0)
Rainfall (mm)				
	Median (IQR)	1254 (1156–1605)	1234 (1229–1241)	1527 (1448–1612)
Slope (°)				
	Median (IQR)	2.3 (1.3–4.4)	1.2 (1.1–1.4)	6.6 (6.0–7.3)
Distance to water course (in km)				
	Median (IQR)	1.1 (0.7–1.7)	0.9 (0.5–1.4)	1.3 (0.8–2.1)
EVI minimum				
	Median (IQR)	0.17 (0.15–0.20)	0.16 (0.15–0.18)	0.20 (0.19–0.21)
Distance to Lake Nyasa* (in km)				
Below 1	%	1.1	0	0
1–2	%	2.2	0	0
2–4	%	2.5	0	0
4 and above	%	94.2	100	100

Results for all study sites combined, for the site with the highest infection prevalence (Igurusi) and the site without infection (Santylia). IQR = interquartile range, SES = socio-economic status, LST = land-surface temperature, EVI = enhanced vegetation index.

*Distances below 4 km only apply to participants from Kyela site

<https://doi.org/10.1371/journal.pntd.0008508.t001>

(577/4189). Restricting the analyses to school-aged children between 5 and 18 years of age only, an overall prevalence of 9.1% (CI: 8.4–9.8%) was found with site-specific prevalences varying between 0.0 and 29.9% (Table 2). For children under 5 years of age the prevalences were low (see S3 Table). Infections were mostly of light intensity (815/914, 89.2%), heavy




Fig 3. Prevalence of *S. haematobium* infection in the nine EMINI study sites in Mbeya region, Tanzania. Households with at least one infected person are represented by yellow to red Voronoi polygons, households without infection are shown in green. (Original work of the authors).

<https://doi.org/10.1371/journal.pntd.0008508.g003>

infections were found in 99 of the 914 infected study participants (10.8%). Prevalences for the following survey rounds are reported in the [S4 Table](#).

Univariable analysis of potential risk factors

Results of the univariable analysis are shown in [Table 3](#). Using non-spatial modelling we found a statistically significant positive association between *S. haematobium* infection and number of persons in the household, land-surface temperature during the night, distance to nearest water course and distance to Lake Nyasa between 2 and 4 km when compared to a distance of 4 km or more. Compared to participants above 35 years of age, younger participants had higher odds of infection. The results were significant for the age groups 5–15, 15–25 and 25–35 years. Children aged 0–5 years were not significantly different from the reference group in terms of *S. haematobium* infection status. The age structure of our cohort and the infection prevalence over age are shown in [Fig 4](#).

In the non-spatial models we found statistically significant negative univariable associations of *S. haematobium* infection with HIV infection, elevation and EVI.

Incorporating the spatial component into the univariable analyses ([Table 3](#)) resulted in the following changes: Population density (*i.e.* number of persons per km²) and LST day became significant positive predictors of *S. haematobium* infection pointing towards higher odds of infection with increasing population and with higher LST during the day. Slope became a significant negative predictor of schistosomiasis infection with steeper slopes corresponding to lower odds of infection. In contrast, LST during the night and elevation turned non-significant when taking the spatial correlation into account.

Multivariable analysis of potential risk factors

In the multivariable non-spatial and spatial models ([Table 4](#)), the effect of age on *S. haematobium* infection remained similar to that of the univariable models: compared to participants

Table 2. Pre-treatment *S. haematobium* infection prevalences in the nine EMINI study sites in Mbeya Region, Tanzania.

Study site	<i>S. haematobium</i> (total population)			<i>S. haematobium</i> (school-aged children)		
	No. of participants (households)	No. of infections	Infection prevalence % (95% CI)	No. of participants	No. of infections	Infection prevalence % (95% CI)
Igurusi	1947 (514)	307	15.8 (14.2–17.5)	695	208	29.9 (26.6–33.4)
Itaka	1924 (369)	225	11.7 (10.3–13.2)	809	171	21.1 (18.5–24.1)
Kyela	2261 (534)	184	8.1 (7.1–9.3)	853	96	11.3 (9.3–13.6)
Mlowo	2139 (494)	100	4.7 (3.9–5.7)	898	82	9.1 (7.4–11.2)
Utengule	1722 (454)	62	3.6 (2.8–4.6)	627	36	5.7 (4.2–7.9)
Iyunga	1503 (336)	19	1.3 (0.8–2.0)	641	10	1.6 (0.8–2.9)
Ruanda	1676 (406)	15	0.9 (0.5–1.5)	669	6	0.9 (0.4–2.0)
Isongole	2105 (542)	2	0.1 (0.0–0.4)	806	1	0.1 (0.0–0.9)
Santylia	2003 (540)	0	0	730	0	0
All Sites	17280 (4189)	914	5.3 (5.0–5.6)	6728	610	9.1 (8.4–9.8)

Shown are the prevalences in the total population (left) and for school-aged children only (5 to 18 years, right). CI = confidence interval.

<https://doi.org/10.1371/journal.pntd.0008508.t002>

above 35 years of age, younger participants between 5 and 35 years were more likely to be infected with *S. haematobium*. Another significant positive association in both modelling approaches was found for the distance to water course, meaning that the odds of *S. haematobium* infection increased with increasing distance from water course. Regarding distance to Lake Nyasa, higher odds of infection were found for short distances to the lake compared to the reference category of 4 km and above for both non-spatial and spatial models, respectively.

We found a negative association of *S. haematobium* infection with minimum EVI for both modelling approaches. For the non-spatial model the odds of infection decreased 4.8 fold (1/0.21) for each 0.1 unit increase in EVI and for the spatial model the decrease of odds of infection was with the factor of 6.7 (1/0.15) for each 0.1 unit increase even higher. Elevation showed non-spatially a highly significant association with *S. haematobium* infection (OR = 0.74, CI: 0.64–0.84), which didn't reach statistical significance after taking the spatial correlation into account (OR = 0.69, CI: 0.46–1.04). In contrast to this, population density became highly significant in the spatial model (non-spatial: OR = 1.08, CI: 0.97–1.20; spatial: OR = 1.27, CI: 1.10–1.46).

Elevation and LST during day and night both had strong univariable associations with *S. haematobium* infection but were also strongly collinear, which prevented inclusion into the same multivariable model. Since elevation is more often used in the literature than LST, we decided to include elevation in our models for better comparability. The negative association of HIV with *S. haematobium* infection, found in the univariable analysis, was not significant anymore in the multivariable models, although the adjusted odds of *S. haematobium* infection in HIV-positive participants were still lower than in HIV-negative participants (non-spatial: OR = 0.64, CI: 0.40–1.03; spatial OR = 0.63, CI: 0.39–1.00). Gender and socio-economic status,

Table 3. Univariable association of socio-demographic and environmental factors with *S. haematobium* infection.

All study sites (N = 17280)			Univariable non-spatial*			Univariable spatial [§]		
Covariate	N	% pos.	OR	95% CI	p value	OR	95% CI	p value
Sex								
Female*	9194	4.95	1.00	-	-	1.00	-	-
Male	8086	5.68	1.11	0.95–1.29	0.1865	1.09	0.94 – 1.27	0.2300
Age (years)								
below 5	2233	2.15	1.16	0.77–1.74	0.4741	1.19	0.79 – 1.79	0.4105
5–15	5534	8.98	8.04	6.05–10.70	<0.0001	8.00	6.04 – 10.60	<0.0001
15–25	3063	7.61	5.97	4.40–8.10	<0.0001	5.96	4.41 – 8.06	<0.0001
25–35	2336	2.65	1.60	1.09–2.35	0.0161	1.63	1.11 – 2.39	0.0128
35 and above*	4114	1.80	1.00	-	-	1.00	-	-
HIV infection								
No*	15700	5.57	1.00	-	-	1.00	-	-
Yes	1209	2.07	0.32	0.21–0.51	<0.0001	0.32	0.21 – 0.51	<0.0001
No information	371	4.04	0.49	0.27–0.88	0.0176	0.51	0.29 – 0.91	0.0219
SES score (per 1 unit)								
			0.92	0.82–1.03	0.1483	0.99	0.88 – 1.10	0.8300
Number of persons in household (per person)								
			1.05	1.02–1.09	0.0024	1.05	1.02 – 1.08	0.0012
Population density (per 1000 persons/km²)								
			1.02	0.93–1.13	0.6510	1.20	1.06 – 1.36	0.0045
Household with latrine								
No*	419	8.35	1.00	-	-	1.00	-	-
Yes	16861	5.21	0.63	0.37–1.09	0.0970	0.64	0.39 – 1.04	0.0709
Elevation (per 100 m)								
			0.71	0.63–0.80	<0.0001	0.72	0.51 – 1.00	0.0526
LST day (per 1°C)								
			1.04	0.96–1.13	0.2950	1.30	1.11 – 1.53	0.0014
LST night (per 1°C)								
			1.34	1.16–1.56	0.0001	1.27	0.92 – 1.75	0.1468
Rainfall (per 100 mm)								
			0.99	0.89–1.10	0.8977	0.88	0.65 – 1.19	0.4030
Slope (per 1°)								
			0.95	0.88–1.04	0.2794	0.88	0.80 – 0.97	0.0091
Distance to water course (in km)								
			1.29	1.15–1.45	<0.0001	1.15	1.02 – 1.30	0.0207
EVI minimum (per 0.1 units)								
			0.48	0.31–0.73	0.0008	0.19	0.09 – 0.37	<0.0001
EVI average (per 0.1 units)								
			0.63	0.41–0.94	0.0255	0.34	0.20 – 0.60	0.0001
EVI maximum (per 0.1 units)								
			0.93	0.67–1.29	0.6585	0.68	0.46 – 1.00	0.0502
Distance to Lake Nyasa (in km)								
below 1	195	9.23	1.63	0.78–3.43	0.1974	3.71	0.78 – 17.60	0.0994
1–2	372	9.14	1.56	0.86–2.83	0.1397	3.26	0.95 – 11.21	0.0611
2–4	428	11.92	2.15	1.25–3.70	0.0059	4.01	1.59 – 10.07	0.0032
4 and above*	16285	4.98	1.00	-	-	1.00	-	-

*Univariable mixed effects logistic regression model with random effects for households and study sites (N = 17,280).

[§]Univariable generalized additive mixed model of the binomial family with spatially correlated effects base on the location of the observation, with additional random effects for households and study sites (N = 17,280). The outcome for both modelling approaches was the binary *S. haematobium* infection status. N = number of participants, % pos. = prevalence of *S. haematobium* infection for categorical variables, OR = odds ratio, CI = confidence interval, SES = socio-economic status, LST = land-surface temperature, EVI = enhanced vegetation index. The reference category for categorical variables is denoted by the asterisk *.

<https://doi.org/10.1371/journal.pntd.0008508.t003>

which were included in the models as potential confounders, were both not significantly associated with *S. haematobium* infection.

HIV and egg excretion

Among the 914 *S. haematobium* infected participants, a total of 25 individuals were co-infected with HIV. The geometric mean egg counts per centiliter of urine were 3.3 (CI: 1.8–6.2) and 4.8




Fig 4. *S. haematobium* prevalence by age. The red line shows LOWESS-smoothed *S. haematobium* infection prevalence, grey bars indicate the number of participants in each age stratum.

<https://doi.org/10.1371/journal.pntd.0008508.g004>

(CI: 4.3–5.4) EPC in HIV positive and HIV negative individuals, respectively. When analyzing the association of egg counts with HIV infection in these 914 participants, the univariable mixed effects negative binomial model showed an egg count ratio of 0.90 (95% CI: 0.69–1.17) for HIV positive compared to HIV negative individuals (see S5 Table). When additionally adjusting for age, sex, and SES, an egg count ratio of 1.68 (95% CI: 0.98–2.90) for HIV positive, compared to HIV negative participants was found, showing a trend towards increased *S. haematobium* egg excretion in HIV infected individuals.

Discussion

Our results show significant associations of individually assessed and remotely sensed factors with *S. haematobium* infection in Mbeya Region in Southwestern Tanzania. Using multivariable modelling, we found increased odds of infection in school-age children and young adults, with increasing distance to water course and decreasing distance to Lake Nyasa, whereas higher vegetation cover was associated with lower odds of infection. After including the spatial component into the analyses, elevation was not significant anymore, whereas population density became a significant predictor of schistosomiasis. The overall prevalence was 5.3%, ranging from 0 to 15.8% in the different study sites showing the spatial heterogeneity which is typical for schistosomiasis. These prevalences are in line with spatial predictions made for Tanzania by Brooker et al., where the prevalence predicted for our study region was also low (*i.e.* the estimated probability for our study area having an infection prevalence >50% was 0–10%) [37]. Considering only school-age children, we found a higher overall prevalence of 9.1% with site-specific prevalences up to 29.9%.

Table 4. Multivariable association of socio-demographic and environmental factors with *S. haematobium* infection.

All study sites (N = 17280)			Multivariable non-spatial*			Multivariable spatial [§]		
Covariate	N	% pos.	OR	95% CI	p value	OR	95% CI	p value
Sex								
Female*	9194	4.95	1.00	-	-	1.00		
Male	8086	5.68	1.03	0.87–1.21	0.7588	1.02	0.88 – 1.20	0.7653
Age (years)								
below 5	2233	2.15	1.15	0.77–1.74	0.4935	1.12	0.74 – 1.68	0.5932
5–15	5534	8.98	7.82	5.85–10.43	<0.0001	7.26	5.50 – 9.60	<0.0001
15–25	3063	7.61	5.84	4.30–7.95	<0.0001	5.51	4.09 – 7.42	<0.0001
25–35	2336	2.65	1.64	1.12–2.40	0.0119	1.57	1.07 – 2.29	0.0213
35 and above*	4114	1.80	1.00	-	-	1.00	-	-
HIV infection								
No*	15700	5.57	1.00	-	-	1.00	-	-
Yes	1209	2.07	0.64	0.40–1.03	0.0637	0.63	0.39 – 1.00	0.0524
No information	371	4.04	0.70	0.36–1.33	0.2752	0.72	0.39 – 1.34	0.3021
SES score (per 1 unit)								
			0.90	0.79–1.02	0.1004	0.89	0.79 – 1.01	0.0631
Population density (per 1000 persons/km²)								
			1.08	0.97–1.20	0.1559	1.27	1.10 – 1.46	0.0008
Elevation (per 100 m)								
			0.74	0.64–0.84	<0.0001	0.69	0.46 – 1.04	0.0750
EVI minimum (per 0.1 units)								
			0.21	0.11–0.39	<0.0001	0.15	0.07 – 0.33	<0.0001
Distance to water course (in km)								
			1.37	1.20–1.57	<0.0001	1.17	1.03 – 1.34	0.0186
Distance to Lake Nyasa (in km)								
below 1	195	9.23	4.53	1.80–11.38	0.0013	4.96	0.91 – 27.08	0.0642
1–2	372	9.14	3.51	1.65–7.47	0.0011	4.11	1.06 – 15.94	0.0413
2–4	428	11.92	3.33	1.68–6.59	0.0006	4.31	1.56 – 11.93	0.0048
4 and above*	16285	4.98	1.00	-	-	1.00	-	-

*Multivariable mixed effects logistic model with random effects for households and study sites (N = 17,280).

[§]Multivariable generalized additive mixed model of binomial family with spatially correlated effects base on the location of the observation, with additional random effects for households and study sites (N = 17,280). The outcome for both modelling approaches was the binary *S. haematobium* infection status. N = number of participants, % pos. = prevalence of *S. haematobium* infection for categorical variables, OR = prevalence ratio, CI = confidence interval, SES = socio-economic status, EVI = enhanced vegetation index. The reference category for categorical variables is denoted by the asterisk *.

<https://doi.org/10.1371/journal.pntd.0008508.t004>

Male sex did not show a statistically significant association with *S. haematobium* infection in our study, even after spatial adjustment. Previous studies report inconsistent associations of sex with *S. haematobium* prevalence [15, 38–42]. Women and men can have different water contact behaviour relating to activities such as swimming, fishing, and doing laundry but the frequency and type of water contact also depend on the water sources available in a community.

Age is a significant predictor of *S. haematobium* infection. In our study population, schistosomiasis mainly affects children and young adults, which is consistent with the literature [2, 43]. Many other studies are concerned with quantifying *S. haematobium* infection in children only, whereas we report results from a population-based study including all age groups and thus show the distribution of *S. haematobium* infection in the whole population.

The assessment of HIV as a factor with potential influence on *S. haematobium* infection is limited by the fact that both diseases show distinct age patterns with peaks at different ages: As opposed to *S. haematobium* infection, which is most prevalent in school age children, the prevalence of HIV peaks at around 35 years of age, when *S. haematobium* prevalence has already dropped to low levels. Thus the strong negative association of HIV and *S. haematobium*

infection in the univariable analysis (OR = 0.40) is at least partly due to the different age peaks that both diseases occur at, although a similar trend towards lower *S. haematobium* prevalence in HIV infected participants is also visible in the age-adjusted multivariable model (non-spatial OR = 0.64). We further investigated the relationship between *S. haematobium* egg excretion and HIV by means of mixed effects negative binomial regression with egg-counts as the outcome in *S. haematobium* infected participants only. The adjusted model showed a trend towards increased *S. haematobium* egg excretion in HIV co-infected individuals (S5 Table). This positive association mainly results from the inclusion of age as a covariate into the model, and again is caused by different age/infection patterns of the two diseases.

Results in the literature regarding the interaction between HIV and schistosomiasis are conflicting. Some studies have found an increased risk of HIV transmission in schistosomiasis infected individuals [44–46], which is especially severe for women affected by genital schistosomiasis [8]. However, microscopic diagnosis of schistosomiasis in HIV infected individuals appears to lack sensitivity [47, 48]. Several articles describe significantly [49–52] or non-significantly [53, 54] decreased *S. mansoni* and/or *S. haematobium* egg excretion in HIV co-infected individuals. Other studies find no association of HIV status with schistosome egg-excretion after adjusting for age [55, 56] or even slightly higher mean egg counts (non-significant) in HIV-positives [57]. Some of these differing findings could be explained by different levels of immune suppression in the populations that were studied [50, 52, 58], and by sex differences regarding the impact HIV has on *Schistosoma* egg excretion [47].

We found a significant positive univariable association of *S. haematobium* infection with the number of household members, but this effect vanished when adjusting for other covariates, thus it was likely caused by confounding, e.g. larger households in an area with high infection prevalence. However, Sady *et al.* found an association between schistosomiasis and presence of other infected family members, which also remained significant in the multivariable analysis [59]. It seems a likely explanation that members of the same household use the same water bodies for their daily activities which leads to household clustering of the infection.

We had expected a significant association between socioeconomic status and *S. haematobium* infection since *S. haematobium* infection is regarded as a disease of poverty. Our findings were not statistically significant, but nevertheless, the odds of infection decreased with increasing SES. Previous studies have reasoned that those living at poverty level are at an increased risk of schistosomiasis due to a greater exposure to unsafe water, less education about transmission and exposure reduction strategies, and limited access to effective treatment [6, 60, 61]. In our study SES showed the strongest association with *S. haematobium* infection in the spatially adjusted models. Another study where SES was calculated in a manner similar to ours, found an association between SES and *S. haematobium* infection [62]. One reason for the lack of association might be that the vast majority of participants in the rural study sites do not have access to safe water for bathing, swimming and washing of clothes, independent of SES.

Population density showed a statistically significant positive association with *S. haematobium* infection only in the spatially adjusted models, indicating increased transmission in densely populated areas. A similar association was found by Koroma *et al.* using geostatistical modelling [17]. In a study conducted by Nagi *et al.*, population density was a significant risk factor for *S. mansoni* infection [63].

Living at higher altitude was associated with reduced odds of urinary schistosomiasis, which is also found in other studies [e.g. 20, 64–66]. Higher altitude is associated with steeper slopes and less stagnant water, which in turn leads to less temporary water bodies and to less suitable habitats for the snails. In our study the site with no infected participants was the highest altitude site. High elevation is associated with lower temperatures, too. In the univariable analysis LST night was highly significantly associated with *S. haematobium* infection indicating

higher odds of infection with increasing night-time temperature. However, after spatial adjustment this effect turned non-significant. This finding is in line with the study by Clements *et al.* [13], where an association between *S. haematobium* and minimum LST was found, which turned non-significant, when including the spatial correlation into the model. Schur *et al.* [20] used spatial modelling and found a non-linear relation between *S. haematobium* with night LST indicating a positive relation and a decrease in risk at the lowest and the highest values LST.

Regarding LST day in the univariable analysis, we found a positive significant effect only after including the spatial correlation structure. While Clark *et al.* also found a positive association between *S. mansoni* and LST using geostatistical models [67], negative associations were found in other studies [20, 68].

Slope of the terrain showed negative odds of infection after spatial adjustment of the univariable model, indicating that steeper slopes lead to decreased *S. haematobium* infection. Kulinkina *et al.* also found a negative association of slope and *S. haematobium* infection [69]. Probable reasons for this negative association of slope with infection are the same as those discussed above for increasing altitude.

As opposed to other studies, we found no stronger associations of rainfall with *S. haematobium* infection [20, 68]. Such an association was also found by Clements *et al.* [13], it however diminished when incorporating the spatial correlation into the analysis. Potential explanations why rainfall did not appear to play an important role in our study, might be a lack of variation in rainfall in our relatively small study area, the low spatial resolution of the rainfall data and the lack of seasonal resolution both for the rainfall and for the schistosomiasis data.

Considering that *S. haematobium* infection requires contact with contaminated water, it was surprising to find an increased prevalence of infection with increasing distance to the nearest water course, i.e. stream or river. Water velocity is an important factor in the transmission of schistosomiasis since the survival of the intermediate host snails can be impacted in fast-flowing water [70]. Even if Rabone *et al.* [71] found a negative association between water speed and *Bulinus spp.* abundance, this does not rule out that the snails may tolerate medium flow speeds for short periods of time. *Bulinus spp.* generally seem to prefer low flow environments [72]. Furthermore, shallow water bodies and still waters are conducive to the growth of algae and aquatic plants on which the snail intermediate hosts feed [73]. Lakes, ponds and pools thus provide the conditions favored by both the snail intermediate host and the free-swimming forms of the schistosome. People living further away from a stream likely more often use ponds or pools to do their washing, swimming and fishing, and thus increase their risk of infection. This is also supported by our finding that *S. haematobium* prevalence increased with decreasing distance to Lake Nyasa in the one site that is situated close to the lake. In the literature, one study found an increased prevalence of schistosomiasis in children living close to streams, springs, pools or ponds [59], whereas in another study, the distance to river was not associated with *S. haematobium* infection [62]. For our study, information regarding smaller stagnant waterbodies would have been interesting, but unfortunately our remotely sensed data only contained information regarding the few larger lakes in the area. We also have no malacological information regarding the presence of suitable host snails in our study area, but at least for Lake Nyasa, *Biomphalaria pfeifferi* (intermediate host snails for *S. mansoni*) have been found in the southern parts of the lake [74].

We found decreased odds of *S. haematobium* infection with increasing amount of vegetation, as measured by the EVI. Sturrock *et al.* [75] also observed significantly decreased odds of *S. haematobium* infection with increasing EVI in their model without spatial component, which however vanished after including the spatial component into the analyses. Some studies have found a significant positive association between *S. haematobium* and/or *S. mansoni* and

the Normalized Difference Vegetation Index (NDVI), which is very similar to EVI [19, 67, 76], whereas in other studies NDVI was not associated with schistosomiasis in the preliminary analyses and thus excluded from further analyses [77, 78]. Another study observed a non-linear association between schistosomiasis and NDVI [20]. However, EVI characterizes the land vegetation cover rather than aquatic vegetation, which is more relevant to the intermediate host snails. A recent study by Wood *et al.* [79] found tight associations between certain kinds of aquatic vegetation and the intermediate host snails, which further hints towards aquatic vegetation being the key for schistosomiasis transmission. Another possible explanation of the reversed association between schistosomiasis and EVI is related to population density. Lower EVI also indicates higher land-clearing and higher population density. Indeed, the two sites with the highest population density had the lowest EVI.

Our study has several limitations: First, data for this study came from a general survey and data collection was not specifically targeted to investigate risk factors of *S. haematobium* infection. Our study would have benefited from more focused questions regarding water contact activities, exposure and occupation. Although some of the environmental factors considered here were strongly related to schistosomiasis, they probably do not fully explain the spatial pattern of *S. haematobium* infection in the study area, since we neither have information regarding the occurrence of suitable snail intermediate hosts, nor data regarding smaller stagnant or temporary waterbodies, which both would be important to get a better picture of transmission potential in an area. While such data collection might be feasible for small-scale studies, it is not easily available across larger spatial scales. Two recent studies have shown that intermediate host snail populations are very patchy in time and space [71, 79] and thus targeting them for control campaigns would require effort exceeding what is feasible. However, Wood *et al.* identified environmental proxies for the snails which were shown to better predict schistosomiasis infection risk than the actual snail abundance data. The area of snail-suitable habitat within a water contact site and the total area of the water contact site were identified as such environmental proxies [79]. Since we are lacking data on small ponds and water contact sites, such analyses are left for future studies.

Due to collinearity with elevation, we couldn't include LST into multivariable models and thus cannot evaluate its impact after adjusting for other important factors. We collected two urine samples (one morning and one spot sample between 10 am and 2 pm for a maximized egg output [80]) but may still have missed some light infections: especially chronically infected adults pass fewer eggs in the urine and where lesions and fibrous tissue have already developed, more eggs are trapped within the body [81].

Despite these limitations, our study also has a couple of unique strengths. The study covers a medium sized area with a large and representative study population of more than 17,000 participants, for which we have individual data regarding *S. haematobium* infection, place of residence and individual risk factors. In contrast, most other studies cited here either have individual data on relatively small study populations from few communities, or they cover entire provinces or countries, and use school prevalences instead of individual outcome data, which means that the population above school age is not considered, and that only pooled data on outcomes and risk factors can be analysed. Regarding some of the different findings of our study compared to others, and also between other studies, one should keep in mind that "scale matters" [82], i.e. that the above mentioned differences in scale between studies might also explain some of the different findings.

Conclusions

S. haematobium infection investigated in this study in Mbeya region of Southwestern Tanzania revealed highly focal infection with prevalences between 0 and 16% in the different study sites.

Our study did not find any *S. haematobium* infection in the site with the highest altitude. The inverse relationship between elevation and *S. haematobium* infection indicates that lower altitude sites are at higher risk of schistosomiasis. We also found a strong negative association of EVI with schistosomiasis and a strongly elevated infection risk in children. Elevation and vegetation density can easily be obtained from remote sensing data that are available free of charge in the public domain, whereas age data are not easy to obtain. Proximity to Lake Nyasa was also associated with increased odds of infection. The positive association of distance to flowing water might suggest that smaller temporary water bodies potentially housing infected intermediate host snails are preferred when the river is too far away for regular water contacts. Population density was another important risk factor for the infection. Regarding public health implications, people living at low altitude and school-aged children living in crowded areas might be considered as the persons at higher risk for *S. haematobium* infection. These findings from our large population-based study of more than 17,000 participants and the reported pre-treatment prevalences could help to evaluate and improve ongoing and future control activities in the region, in Tanzania and elsewhere.

Supporting information

S1 Checklist. STROBE Checklist.

(DOC)

S1 Table. Multivariable association of socio-demographic factors only with *S. haematobium* infection. Results of multivariable mixed effects logistic regression with site and household as random effects and with binary *S. haematobium* infection status as the outcome (N = 17,280). *Multivariable mixed effects logistic model with random effects for households and study sites. §Multivariable generalized additive mixed model of binomial family with spatially correlated effects base on the location of the observation, with additional random effects on households and study sites. OR = odds ratio, CI = confidence interval, SES = socio-economic status. The reference category for stratified variables is denoted by the asterisk *.

(DOCX)

S2 Table. Full model: Association of socio-demographic and environmental factors with *S. haematobium* infection. *Multivariable mixed effects logistic model with random effects for households and study sites. §Multivariable generalized additive mixed model of binomial family with spatially correlated effects base on the location of the observation, with additional random effects for households and study sites.

(DOCX)

S3 Table. *S. haematobium* infection prevalences in children below 5 years of age. CI: confidence interval.

(DOCX)

S4 Table. Prevalence of *S. haematobium* infection over all survey rounds of the EMINI study. CI: confidence interval.

(DOCX)

S5 Table. Association of HIV status with *S. haematobium* infection intensity. Results of multivariable mixed effects negative binomial regression with site and household as random effects and with *S. haematobium* egg counts as the outcome, adjusted for socio-demographic factors. Only includes *S. haematobium* infected participants (N = 914). ECR = egg count ratio, CI = confidence interval, SES = socio-economic status. The reference category for stratified

variables is denoted by the asterisk *.
(DOCX)

S1 Text. Details on variable selection.
(DOCX)

Acknowledgments

We would like to thank the participants of the EMINI study and the EMINI field, laboratory and coordination teams at MMRC for their support throughout the study.

Author Contributions

Conceptualization: Michael Hoelscher, Elmar Saathoff.

Data curation: Elmar Saathoff.

Formal analysis: Kirsi M. Manz, Ursula Berger, Elmar Saathoff.

Funding acquisition: Michael Hoelscher.

Investigation: Inge Kroidl, Petra Clowes, Martina Gerhardt, Wilbrod Nyembe, Lucas Maganga, Weston Assisya, Elmar Saathoff.

Project administration: Petra Clowes, Lucas Maganga, Nyanda E. Ntinginya.

Resources: Petra Clowes, Weston Assisya, Michael Hoelscher, Elmar Saathoff.

Supervision: Inge Kroidl, Petra Clowes, Martina Gerhardt, Lucas Maganga, Weston Assisya, Nyanda E. Ntinginya, Michael Hoelscher, Elmar Saathoff.

Visualization: Kirsi M. Manz, Elmar Saathoff.

Writing – original draft: Kirsi M. Manz, Elmar Saathoff.

Writing – review & editing: Kirsi M. Manz, Inge Kroidl, Petra Clowes, Martina Gerhardt, Wilbrod Nyembe, Lucas Maganga, Weston Assisya, Nyanda E. Ntinginya, Ursula Berger, Michael Hoelscher, Elmar Saathoff.

References

1. World-Health-Organization. Schistosomiasis Fact sheet. Assessed on March 18th, 2020. Available from: <https://www.who.int/en/news-room/fact-sheets/detail/schistosomiasis>.
2. Colley DG, Bustinduy AL, Secor WE, King CH. Human schistosomiasis. *Lancet*. 2014; 383(9936):2253–64. [https://doi.org/10.1016/S0140-6736\(13\)61949-2](https://doi.org/10.1016/S0140-6736(13)61949-2) PMID: 24698483
3. Brooker S, Kabatereine NB, Smith JL, Mupfasoni D, Mwanje MT, Ndayishimiye O, et al. An updated atlas of human helminth infections: the example of East Africa. *Int J Health Geogr*. 2009; 8:42. <https://doi.org/10.1186/1476-072X-8-42> PMID: 19589144
4. Sturrock RF. The intermediate hosts and host-parasite relationships. In: Jordan P, Webber G, Sturrock RF, editors. *Human Schistosomiasis*. Wallingford: CAB International. 1993:33–85.
5. Gray DJ, Ross AG, Li YS, McManus DP. Diagnosis and management of schistosomiasis. *BMJ*. 2011; 342:d2651. <https://doi.org/10.1136/bmj.d2651> PMID: 21586478
6. van der Werf MJ, de Vlas SJ, Brooker S, Looman CW, Nagelkerke NJ, Habbema JD, et al. Quantification of clinical morbidity associated with schistosome infection in sub-Saharan Africa. *Acta Trop*. 2003; 86(2–3):125–39. [https://doi.org/10.1016/s0001-706x\(03\)00029-9](https://doi.org/10.1016/s0001-706x(03)00029-9) PMID: 12745133
7. Downs JA, Mguta C, Kaatano GM, Mitchell KB, Bang H, Simplicio H, et al. Urogenital schistosomiasis in women of reproductive age in Tanzania's Lake Victoria region. *The American journal of tropical medicine and hygiene*. 2011; 84(3):364–9. <https://doi.org/10.4269/ajtmh.2011.10-0585> PMID: 21363971

8. Hotez PJ, Harrison W, Fenwick A, Bustinduy AL, Ducker C, Sabina Mbabazi P, et al. Female genital schistosomiasis and HIV/AIDS: Reversing the neglect of girls and women. *PLoS neglected tropical diseases*. 2019; 13(4):e0007025. <https://doi.org/10.1371/journal.pntd.0007025> PMID: 30946746
9. Kjetland EF, Kurewa EN, Ndhlovu PD, Midzi N, Gwanzura L, Mason PR, et al. Female genital schistosomiasis—a differential diagnosis to sexually transmitted disease: genital itch and vaginal discharge as indicators of genital *Schistosoma haematobium* morbidity in a cross-sectional study in endemic rural Zimbabwe. *Tropical medicine & international health: TM & IH*. 2008; 13(12):1509–17.
10. Simoonga C, Utzinger J, Brooker S, Vounatsou P, Appleton CC, Stensgaard AS, et al. Remote sensing, geographical information system and spatial analysis for schistosomiasis epidemiology and ecology in Africa. *Parasitology*. 2009; 136(13):1683–93. <https://doi.org/10.1017/S0031182009006222> PMID: 19627627
11. Walz Y, Wegmann M, Dech S, Vounatsou P, Poda JN, N'Goran EK, et al. Modeling and Validation of Environmental Suitability for Schistosomiasis Transmission Using Remote Sensing. *PLoS neglected tropical diseases*. 2015; 9(11):e0004217. <https://doi.org/10.1371/journal.pntd.0004217> PMID: 26587839
12. Vounatsou P, Raso G, Tanner M, N'goran EK, Utzinger J. Bayesian geostatistical modelling for mapping schistosomiasis transmission. *Parasitology*. 2009 Nov; 136(13):1695–705. <https://doi.org/10.1017/S003118200900599X> PMID: 19490724
13. Clements AC, Lwambo NJ, Blair L, Nyandindi U, Kaatano G, Kinung'hi S, et al. Bayesian spatial analysis and disease mapping: tools to enhance planning and implementation of a schistosomiasis control programme in Tanzania. *Tropical medicine & international health*. 2006; 11(4):490–503.
14. Chamartin F, Hougbedji CA, Hürlimann E, Yapi RB, Silué KD, Soro G, et al. Bayesian risk mapping and model-based estimation of *Schistosoma haematobium*-*Schistosoma mansoni* co-distribution in Côte d'Ivoire. *PLoS Negl Trop Dis*. 2014 Dec 18; 8(12):e3407. <https://doi.org/10.1371/journal.pntd.0003407> PMID: 25522007
15. Clements AC, Barnett AG, Nyandindi U, Lwambo NJ, Kihamia CM, Blair L. Age and gender effects in self-reported urinary schistosomiasis in Tanzania. *Tropical medicine & international health*. 2008; 13(5):713–21.
16. Beck-Wörner C, Raso G, Vounatsou P, N'Goran EK, Rigo G, Parlow E, et al. Bayesian spatial risk prediction of *Schistosoma mansoni* infection in western Côte d'Ivoire using a remotely-sensed digital elevation model. *Am J Trop Med Hyg*. 2007 May; 76(5):956–63. PMID: 17488922
17. Koroma JB, Peterson J, Gbakima AA, Nylander FE, Sahr F, et al. Geographical Distribution of Intestinal Schistosomiasis and Soil-Transmitted Helminthiasis and Preventive Chemotherapy Strategies in Sierra Leone. *PLoS Negl Trop Dis* 2010; 4(11): e891. <https://doi.org/10.1371/journal.pntd.0000891> PMID: 21124881
18. Peng WX, Tao B, Clements A, Jiang QL, Zhang ZJ, Zhou YB, et al. Identifying high-risk areas of schistosomiasis and associated risk factors in the Poyang Lake region, China. *Parasitology*. 2010 Jun; 137(7):1099–107. <https://doi.org/10.1017/S003118200999206X> PMID: 20128946
19. Schur N, Hürlimann E, Garba A, Traore MS, Ndir O, Ratard RC, et al. Geostatistical model-based estimates of Schistosomiasis prevalence among individuals aged ≤ 20 years in West Africa. *PLoS neglected tropical diseases*. 2011; 5(6):e1194. <https://doi.org/10.1371/journal.pntd.0001194> PMID: 21695107
20. Schur N, Hürlimann E, Stensgaard AS, Chimfwembe K, Mushingi G, Simoonga C, et al. Spatially explicit *Schistosoma* infection risk in eastern Africa using Bayesian geostatistical modelling. *Acta Trop*. 2013 Nov; 128(2):365–77. <https://doi.org/10.1016/j.actatropica.2011.10.006> PMID: 22019933
21. Manz KM, Clowes P, Kroidl I, Kowuor DO, Geldmacher C, Ntinginya NE, et al. *Trichuris trichiura* infection and its relation to environmental factors in Mbeya region, Tanzania: A cross-sectional, population-based study. *PLoS One*. 2017; 12(4):e0175137. <https://doi.org/10.1371/journal.pone.0175137> PMID: 28384306
22. Riess H, Clowes P, Kroidl I, Kowuor DO, Nsojo A, Mangu C, et al. Hookworm infection and environmental factors in mbeya region, Tanzania: a cross-sectional, population-based study. *PLoS neglected tropical diseases*. 2013; 7(9):e2408. <https://doi.org/10.1371/journal.pntd.0002408> PMID: 24040430
23. Schüle SA, Clowes P, Kroidl I, Kowuor DO, Nsojo A, Mangu C, et al. *Ascaris lumbricoides* infection and its relation to environmental factors in the Mbeya region of Tanzania, a cross-sectional, population-based study. *PLoS One*. 2014; 9(3):e92032. <https://doi.org/10.1371/journal.pone.0092032> PMID: 24643023
24. Prevention and control of schistosomiasis and soil-transmitted helminthiasis. *World Health Organ Tech Rep Ser*. 2002; 912:i-vi, 1–57.
25. Kroidl I, Clowes P, Mwalongo W, Maganga L, Maboko L, Kroidl AL, et al. Low specificity of determine HIV1/2 RDT using whole blood in south west Tanzania. *PLoS One*. 2012; 7(6):e39529. <https://doi.org/10.1371/journal.pone.0039529> PMID: 22768086

26. Filmer D, Pritchett LH. Estimating wealth effects without expenditure data—or tears: an application to educational enrollments in states of India. *Demography*. 2001; 38(1):115–32. <https://doi.org/10.1353/dem.2001.0003> PMID: 11227840
27. Kolenikov S, Angeles G. Socioeconomic status measurement with discrete proxy variables: Is principal component analysis a reliable answer? *Review of Income and Wealth*. 2009; 55(1):128–65.
28. Farr TG, Rosen PA, Caro E, Crippen R, Duren R, Hensley S, et al. The Shuttle Radar Topography Mission. *Reviews of Geophysics*. 2007; 45(2):RG2004.
29. NASA LP DAAC. MODIS 11A2 & MODIS 13Q1. NASA EOSDIS Land Processes DAAC, USGS Earth Resources Observation and Science (EROS) Center, Sioux Falls, South Dakota 2001.
30. R Core Team (2018). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL: <https://www.R-project.org>.
31. Simon Wood and Fabian Scheipl (2017). *gamm4: Generalized Additive Mixed Models using 'mgcv' and 'lme4'*. R package version 0.2–5. <https://CRAN.R-project.org/package=gamm4>.
32. Santos CA, Fiaccone RL, Oliveira NF, Cunha S, Barreto ML, do Carmo MB, et al. Estimating adjusted prevalence ratio in clustered cross-sectional epidemiological data. *BMC Med Res Methodol*. 2008 Dec 16; 8:80. <https://doi.org/10.1186/1471-2288-8-80> PMID: 19087281
33. Alin A. Multicollinearity. *Wiley Interdisciplinary Reviews: Computational Statistics*. 2010; 2(3):370–4.
34. O'brien RM. A Caution Regarding Rules of Thumb for Variance Inflation Factors. *Quality & Quantity*. 2007; 41(5):673–90.
35. Wood SN. *Generalized Additive Models: An Introduction with R* (2nd edition). CRC/Taylor& Francis. 2017.
36. Wilson K, Grenfell BT. Generalized linear modelling for parasitologists. *Parasitol Today*. 1997; 13(1):33–8. [https://doi.org/10.1016/s0169-4758\(96\)40009-6](https://doi.org/10.1016/s0169-4758(96)40009-6) PMID: 15275165
37. Brooker S, Hay SI, Issae W, Hall A, Kihamia CM, Lwambo NJ, et al. Predicting the distribution of urinary schistosomiasis in Tanzania using satellite sensor data. *Tropical medicine & international health*. 2001; 6(12):998–1007.
38. Angelo T, Buza J, Kinung'hi SM, Kariuki HC, Mwanga JR, Munisi DZ, et al. Geographical and behavioral risks associated with *Schistosoma haematobium* infection in an area of complex transmission. *Parasit Vectors*. 2018; 11(1):481. <https://doi.org/10.1186/s13071-018-3064-5> PMID: 30144807
39. Atalabi TE, Lawal U, Ipinlaye SJ. Prevalence and intensity of genito-urinary schistosomiasis and associated risk factors among junior high school students in two local government areas around Zobe Dam in Katsina State, Nigeria. *Parasit Vectors*. 2016; 9(1):388. <https://doi.org/10.1186/s13071-016-1672-5> PMID: 27388007
40. Geleta S, Alemu A, Getie S, Mekonnen Z, Erko B. Prevalence of urinary schistosomiasis and associated risk factors among Abobo Primary School children in Gambella Regional State, southwestern Ethiopia: a cross sectional study. *Parasit Vectors*. 2015; 8:215. <https://doi.org/10.1186/s13071-015-0822-5> PMID: 25886292
41. Kapito-Tembo AP, Mwapasa V, Meshnick SR, Samanyika Y, Banda D, Bowie C, et al. Prevalence distribution and risk factors for *Schistosoma haematobium* infection among school children in Blantyre, Malawi. *PLoS neglected tropical diseases*. 2009; 3(1):e361. <https://doi.org/10.1371/journal.pntd.0000361> PMID: 19156193
42. Slootweg R, Kooyman M, de Koning P, van Schooten M. Water contact studies for the assessment of schistosomiasis infection risks in an irrigation scheme in Cameroon. *Irrigation and Drainage Systems*. 1993; 7(2):113–30.
43. King CH, Keating CE, Muruka JF, Ouma JH, Houser H, Siogok TK, et al. Urinary tract morbidity in schistosomiasis haematobia: associations with age and intensity of infection in an endemic area of Coast Province, Kenya. *The American journal of tropical medicine and hygiene*. 1988; 39(4):361–8. <https://doi.org/10.4269/ajtmh.1988.39.361> PMID: 3142286
44. Downs JA, van Dam GJ, Changalucha JM, Corstjens PL, Peck RN, de Dood CJ, et al. Association of Schistosomiasis and HIV infection in Tanzania. *The American journal of tropical medicine and hygiene*. 2012; 87(5):868–73. <https://doi.org/10.4269/ajtmh.2012.12-0395> PMID: 23033399
45. Downs JA, Dupnik KM, van Dam GJ, Urassa M, Lutonja P, Cornelis D, et al. Effects of schistosomiasis on susceptibility to HIV-1 infection and HIV-1 viral load at HIV-1 seroconversion: A nested case-control study. *PLoS neglected tropical diseases*. 2017; 11(9):e0005968. <https://doi.org/10.1371/journal.pntd.0005968> PMID: 28945756
46. Kjetland EF, Ndhlovu PD, Gomo E, Mdluzwa T, Midzi N, Gwanzura L, et al. Association between genital schistosomiasis and HIV in rural Zimbabwean women. *AIDS*. 2006; 20(4):593–600. <https://doi.org/10.1097/01.aids.0000210614.45212.0a> PMID: 16470124

47. Colombe S, Lee MH, Masikini PJ, van Lieshout L, de Dood CJ, Hoekstra PT, et al. Decreased Sensitivity of *Schistosoma* sp. Egg Microscopy in Women and HIV-Infected Individuals. *The American journal of tropical medicine and hygiene*. 2018; 98(4):1159–64. <https://doi.org/10.4269/ajtmh.17-0790> PMID: 29405114
48. Mazigo HD, Heukelbach J. Diagnostic Performance of Kato Katz Technique and Point-of-Care Circulating Cathodic Antigen Rapid Test in Diagnosing *Schistosoma mansoni* Infection in HIV-1 Co-Infected Adults on the Shoreline of Lake Victoria, Tanzania. *Tropical medicine and infectious disease*. 2018; 3(2). <https://doi.org/10.3390/tropicalmed3020054> PMID: 30274450
49. N'Zoukoudi-N'Doundou MY, Dirat I, Akouala JJ, Penchenier L, Makuwa M, Rey JL. [Bilharziasis and human immunodeficiency virus infection in Congo]. *Med Trop (Mars)*. 1995; 55(3):249–51.
50. Karanja DM, Colley DG, Nahlen BL, Ouma JH, Secor WE. Studies on schistosomiasis in western Kenya: I. Evidence for immune-facilitated excretion of schistosome eggs from patients with *Schistosoma mansoni* and human immunodeficiency virus coinfections. *The American journal of tropical medicine and hygiene*. 1997; 56(5):515–21. <https://doi.org/10.4269/ajtmh.1997.56.515> PMID: 9180601
51. Fontanet AL, Woldemichael T, Sahlu T, van Dam GJ, Messele T, Rinke de Wit T, et al. Epidemiology of HIV and *Schistosoma mansoni* infections among sugar-estate residents in Ethiopia. *Ann Trop Med Parasitol*. 2000; 94(2):145–55. PMID: 10827869
52. Mwanakasale V, Vounatsou P, Sukwa TY, Ziba M, Ernest A, Tanner M. Interactions between *Schistosoma haematobium* and human immunodeficiency virus type 1: the effects of coinfection on treatment outcomes in rural Zambia. *The American journal of tropical medicine and hygiene*. 2003; 69(4):420–8. PMID: 14640503
53. Mazigo HD, Dunne DW, Wilson S, Kinung'hi SM, Pinot de Moira A, Jones FM, et al. Co-infection with *Schistosoma mansoni* and Human Immunodeficiency Virus-1 (HIV-1) among residents of fishing villages of north-western Tanzania. *Parasit Vectors*. 2014; 7:587. <https://doi.org/10.1186/s13071-014-0587-2> PMID: 25511298
54. Sanya RE, Muhangi L, Nampijja M, Nannozi V, Nakawungu PK, Abayo E, et al. *Schistosoma mansoni* and HIV infection in a Ugandan population with high HIV and helminth prevalence. *Tropical medicine & international health*. 2015; 20(9):1201–8.
55. Kallestrup P, Zinyama R, Gomo E, Butterworth AE, van Dam GJ, Erikstrup C, et al. Schistosomiasis and HIV-1 infection in rural Zimbabwe: implications of coinfection for excretion of eggs. *J Infect Dis*. 2005; 191(8):1311–20. <https://doi.org/10.1086/428907> PMID: 15776378
56. Ndhlovu PD, Mduluzi T, Kjetland EF, Midzi N, Nyanga L, Gundersen SG, et al. Prevalence of urinary schistosomiasis and HIV in females living in a rural community of Zimbabwe: does age matter? *Trans R Soc Trop Med Hyg*. 2007; 101(5):433–8. <https://doi.org/10.1016/j.trstmh.2006.08.008> PMID: 17064746
57. Kleppa E, Klinge KF, Galaphaththi-Arachchige HN, Holmen SD, Lillebo K, Onsrud M, et al. *Shistosoma haematobium* infection and CD4+ T-cell levels: a cross-sectional study of young South African women. *PLoS One*. 2015; 10(3):e0119326. <https://doi.org/10.1371/journal.pone.0119326> PMID: 25768005
58. Dunne DW, Hassounah O, Musallam R, Lucas S, Pepys MB, Baltz M, et al. Mechanisms of *Schistosoma mansoni* egg excretion: parasitological observations in immunosuppressed mice reconstituted with immune serum. *Parasite Immunol*. 1983; 5(1):47–60. <https://doi.org/10.1111/j.1365-3024.1983.tb00722.x> PMID: 6600833
59. Sady H, Al-Mekhlafi HM, Mahdy MA, Lim YA, Mahmud R, Surin J. Prevalence and associated factors of Schistosomiasis among children in Yemen: implications for an effective control programme. *PLoS neglected tropical diseases*. 2013; 7(8):e2377. <https://doi.org/10.1371/journal.pntd.0002377> PMID: 23991235
60. Useh MF, Ejezie GC. Modification of behaviour and attitude in the control of schistosomiasis. 1. Observations on water-contact patterns and perception of infection. *Ann Trop Med Parasitol*. 1999; 93(7):711–20. <https://doi.org/10.1080/00034989957961> PMID: 10715699
61. King CH. Parasites and poverty: the case of schistosomiasis. *Acta Trop*. 2010 Feb; 113(2):95–104. <https://doi.org/10.1016/j.actatropica.2009.11.012> PMID: 19962954
62. Chadeka EA, Nagi S, Sunahara T, Cheruiyot NB, Bahati F, Ozeki Y, et al. Spatial distribution and risk factors of *Schistosoma haematobium* and hookworm infections among schoolchildren in Kwale, Kenya. *PLoS neglected tropical diseases*. 2017; 11(9):e0005872. <https://doi.org/10.1371/journal.pntd.0005872> PMID: 28863133
63. Nagi S, Chadeka EA, Sunahara T, Mutungi F, Justin YKD, et al. Risk Factors and Spatial Distribution of *Schistosoma mansoni* Infection among Primary School Children in Mbita District, Western Kenya. *PLoS Negl Trop Dis* 2014; 8(7): e2991. <https://doi.org/10.1371/journal.pntd.0002991> PMID: 25058653
64. Koukounari A, Toure S, Donnelly CA, Ouedraogo A, Yoda B, Ky C, et al. Integrated monitoring and evaluation and environmental risk factors for urogenital schistosomiasis and active trachoma in Burkina

- Faso before preventative chemotherapy using sentinel sites. *BMC Infect Dis*. 2011; 11:191. <https://doi.org/10.1186/1471-2334-11-191> PMID: 21749703
65. Brooker S. Spatial epidemiology of human schistosomiasis in Africa: risk models, transmission dynamics and control. *Trans R Soc Trop Med Hyg*. 2007; 101(1):1–8. <https://doi.org/10.1016/j.trstmh.2006.08.004> PMID: 17055547
 66. Brooker S, Michael E. The potential of geographical information systems and remote sensing in the epidemiology and control of human helminth infections. *Adv Parasitol*. 2000; 47:245–88. [https://doi.org/10.1016/s0065-308x\(00\)47011-9](https://doi.org/10.1016/s0065-308x(00)47011-9) PMID: 10997209
 67. Clark NJ, Umulisa I, Ruberanziza E, Owada K, Colley DG, Ortu G, et al. Mapping *Schistosoma mansoni* endemicity in Rwanda: a critical assessment of geographical disparities arising from circulating cathodic antigen versus Kato-Katz diagnostics. *PLoS Negl Trop Dis* 2019; 13(9):e0007723. <https://doi.org/10.1371/journal.pntd.0007723> PMID: 31568504
 68. Woodhall DM, Wiegand RE, Wellman M, Matey E, Abudho B, et al. Use of Geospatial Modeling to Predict *Schistosoma mansoni* Prevalence in Nyanza Province, Kenya. *PLoS ONE* 2013; 8(8): e71635. <https://doi.org/10.1371/journal.pone.0071635> PMID: 23977096
 69. Kulinkina AV, Walz Y, Koch M, Biritwum N-K, Utzinger J, Naumova EN (2018) Improving spatial prediction of *Schistosoma haematobium* prevalence in southern Ghana through new remote sensors and local water access profiles. *PLoS Negl Trop Dis* 2018; 12(6): e0006517. <https://doi.org/10.1371/journal.pntd.0006517> PMID: 29864165
 70. Appleton CC. The influence of stream geology on the distribution of the bilharzia host snails, *Biomphalaria pfeifferi* and *Bulinus (physopsis) sp.* *Ann Trop Med Parasitol*. 1975; 69(2):241–55. <https://doi.org/10.1080/00034983.1975.11687007> PMID: 1155991
 71. Rabone M, Wiethase JH, Allan F, Gouvras AN, Pennance T, Hamidou AA, et al. Freshwater snails of biomedical importance in the Niger River Valley: evidence of temporal and spatial patterns in abundance, distribution and infection with *Schistosoma* spp. *Parasit Vectors*. 2019 Oct 22; 12(1):498. <https://doi.org/10.1186/s13071-019-3745-8> PMID: 31640811
 72. Rollinson D, Stothard JR, Southgate VR. Interactions between intermediate snail hosts of the genus *Bulinus* and schistosomes of the *Schistosoma haematobium* group. *Parasitology*. 2001; 123 Suppl: S245–60.
 73. Haddock KC. Control of schistosomiasis: the Puerto Rican experience. *Soc Sci Med D*. 1981; 15(4):501–14. [https://doi.org/10.1016/0160-8002\(81\)90045-9](https://doi.org/10.1016/0160-8002(81)90045-9) PMID: 7330692
 74. Alharbi MH, Condemine C, Christiansen R, LaCourse EJ, Makaula P, Stanton MC, et al. *Biomphalaria pfeifferi* Snails and Intestinal Schistosomiasis, Lake Malawi, Africa, 2017–2018. *Emerg Infect Dis*. 2019; 25(3):613–5. <https://doi.org/10.3201/eid2503.181601> PMID: 30602122
 75. Sturrock HJ, Pullan RL, Kihara JH, Mwandawiro C, Brooker SJ. The use of bivariate spatial modeling of questionnaire and parasitology data to predict the distribution of *Schistosoma haematobium* in Coastal Kenya. *PLoS neglected tropical diseases*. 2013; 7(1):e2016. <https://doi.org/10.1371/journal.pntd.0002016> PMID: 23359829
 76. Sousa-Figueiredo JC, Stanton MC, Katokele S, Arinaitwe M, Adriko M, Balfour L, et al. Mapping of Schistosomiasis and Soil-Transmitted Helminths in Namibia: The First Large-Scale Protocol to Formally Include Rapid Diagnostic Tests. *PLoS neglected tropical diseases*. 2015; 9(7):e0003831. <https://doi.org/10.1371/journal.pntd.0003831> PMID: 26196386
 77. Clements AC, Bosque-Oliva E, Sacko M, Landoure A, Dembele R, Traore M, et al. A comparative study of the spatial distribution of schistosomiasis in Mali in 1984–1989 and 2004–2006. *PLoS neglected tropical diseases*. 2009; 3(5):e431. <https://doi.org/10.1371/journal.pntd.0000431> PMID: 19415108
 78. Soares Magalhaes RJ, Biritwum NK, Gyapong JO, Brooker S, Zhang Y, Blair L, et al. Mapping helminth co-infection and co-intensity: geostatistical prediction in Ghana. *PLoS neglected tropical diseases*. 2011; 5(6):e1200. <https://doi.org/10.1371/journal.pntd.0001200> PMID: 21666800
 79. Wood CL, Sokolow SH, Jones IJ, Chamberlin AJ, Lafferty KD, Kuris AM, et al. Precision mapping of snail habitat provides a powerful indicator of human schistosomiasis transmission. *Proc Natl Acad Sci U S A*. 2019 Nov 12; 116(46):23182–23191. <https://doi.org/10.1073/pnas.1903698116> PMID: 31659025
 80. Weber MD, Blair DM, Clark VV. The pattern of schistosome egg distribution in a micturition flow. *Cent Afr J Med*. 1967; 13(4):75–88. PMID: 6068871
 81. Ibranke OA, Phillips AE, Garba A, Lamine SM, Shiff C. Diagnosis of *Schistosoma haematobium* by detection of specific DNA fragments from filtered urine samples. *The American journal of tropical medicine and hygiene*. 2011; 84(6):998–1001. <https://doi.org/10.4269/ajtmh.2011.10-0691> PMID: 21633040
 82. Dabiri Z, Blaschke T. Scale matters: a survey of the concepts of scale used in spatial disciplines. *Eur J Remote Sens*. 2019 Jun 10; 52(1):419–434. <https://doi.org/10.1080/22797254.2019.1626291> PMID: 31275899